

Index

Page numbers in *italics* denote figures, those in **bold** denote tables.

- ABCDE system 125–30
- Acute Illness Management (AIM) 122
- acute kidney injury **145**
- Acute Life-threatening Event Recognition and Treatment (ALERT) 122
- acutely ill patients 121–58
 - assessment 121–2, 124–30
 - airway 125–6, **126**
 - breathing 126–8, **127**
 - circulation 128–9, **129**
 - disability 129–30, **129**
 - exposure/examination 130
 - clinical scoring systems 122–3, **123**
 - critical care outreach team 123, **124**
 - deterioration in 131
 - emergency management 131–57
 - see also specific conditions*
 - further actions 130–1
 - initial approach to patient 125
 - receiving a call 123–4
 - recognition 121–2
- adrenaline (epinephrine) 52, 94
- Advanced Life Support (ALS) 122
- Advanced Paediatric Life Support (APLS) 122
- Advanced Trauma Life Support (ATLS) 122
- adverse drug reactions
 - anaphylaxis 93–4
 - suxamethonium 44
- air, medical 25
 - cylinder colour **24**
 - see also anaesthetic gases*
- Airtraq 23, 23
- airway 63–69, 63, 64
 - assessment 6–8, 7, 8, 125–6, **126**
 - nasopharyngeal 19, 19, 64–5, 65
 - noises **126**
 - obstruction *see* airway obstruction
 - oropharyngeal 19, 19, 63–4, 64
 - problems 65
 - see also* airway equipment; tracheal intubation
- airway equipment 18–24, 100
 - adjuncts 19
 - facemasks 18–19, 19, 65, 65
 - laryngoscopes 22–4, 67
 - see also* supraglottic airway devices
- airway obstruction 102, **126**
 - lower airway problems, acute 134, 137–41
 - asthma 134, 137
 - pneumonia 137
 - pneumothorax 138
 - pulmonary embolism 140–1, 140
 - pulmonary oedema 139–40, **139**, 140
 - upper airway 132–4
 - blocked tracheostomy 133–4
 - external compression post-surgery 133
 - reduced consciousness level 132
 - swelling/tumour 132–3
 - treatment 133
- albumin **55**
- alcohol intake 6
- alfentanil **47**
- allergies 5, 94
 - latex 94
 - see also* anaphylaxis
- alveolar hypoventilation 101–3, 102
- amethocaine 51
 - risk indicators 12–13, **12**
 - risks 11–12
 - safe delivery 24–31
 - total intravenous 44
- anaesthetic drugs
 - inhaled 42–3, **42**
 - intravenous 40–1, **41**
 - see also individual drugs*
- anaesthetic assessment 3–8
 - drug history and allergies 5
 - exercise tolerance 4–5
 - family history 5
 - investigations 8–11
 - medical history 3–4
 - physical examination 6–8
 - previous anaesthetics/operations 5
 - social history 5–6
- anaesthetic gases 24–5
 - cylinder colours **24**
 - flow control 25–6, 25
 - medical air 25
 - nitrous oxide 24–5, 42, 43
 - oxygen 24
 - pollution 29–30

178 Index

- anaesthetic machines 25, 25, 29, 29
 preoperative check 58
- anaesthetic record 76
- anaesthetic vapours 26, 26
- analgesia
 combined techniques 119
 infiltration 80–1, 80
 NSAIDs 40, 49, **49**
 opioids 46–9, **47**, 113, **115**
 paracetamol 40, 49
 patient-controlled 114–15
 postoperative pain 112–19
 premedication 40
 sites of action 114
see also regional anaesthesia
- analgesic drugs 46–9
- anaphylaxis 93–4
 causes 94
 investigations 94
 management 94
- angina 147
- anti-emetics 40, 50, **50**
- antibiotics
 allergy 94
 premedication 40
- anticholinesterases 45–6
- anticoagulants, prophylactic 40
- antidiuretic hormone (ADH) 111
- antihistamines 94
- anuria 106, 145
see also oliguria
- aorticaval compression 90–1
- Apfel score 50
- arrhythmias 149–50
 bradycardias 107, 150
 postoperative 106–7
 tachycardias 107, 149–50
 and tracheal intubation 69
- arterial blood gases
 pneumonia 137
 pulmonary embolism 140
- arterial cannulation 62
- ASA physical status scale **12**
- aspiration 69, 92–3
 at induction 92–3
 management 92–3
 reducing risks of 88–9, 89
 with supra-glottic airway 93
- aspiration pneumonia 137
- asthma 134
- treatment 134
- atracurium **45**
- AVPU **123**
- basilic vein 60
- bispectral index (BIS) 36
- blood glucose 8
- blood loss 36
- blood pressure **123**
 invasive/direct monitoring 34, **143**
 non-invasive monitoring 31–2
- blood/blood components 55–6
 predepositing 56
 risk of transfusion 55–6
- brachial plexus block 81
- bradycardias 150
 central neural blockade 84
 postoperative 107
- breathing 126–8, **127**
 abnormal 126–7
 rate **123**
- breathing systems 26–8
 circle system 27–8, 27
 components of 26–7
 mechanical ventilation 28
 monitoring 36
- breathlessness *see* dyspnoea
- bronchodilators 94
- Buccastem 109
- bupivacaine 52, **53**
- buprenorphine 48
- BURP manoeuvre 96
- caesarean section 89–91
- Calder test 7
- cannulas, intravascular 30, 30
- capillary refill time 129
- capnometry 33, **33**, 67
- carbon dioxide
 cylinder colour **24**
 PaCO₂ 102, 128
- cardiac arrest 150–3, 151, 152
- cardiac arrhythmias *see* arrhythmias
- cardiac output
 and hypotension 106–7
 low 84
 and mechanical ventilation 103
 oesophageal Doppler monitoring 34–5, 35
 pulse analysis monitoring 35, 35
- cardiogenic shock 144–5
 investigations 145
 treatment 144–5
- cardiopulmonary exercise (CPX) testing 8–9, **9**
- cardiopulmonary resuscitation (CPR) 152
- cardiovascular disease, medical referral 9
- cardiovascular system
 history 3–4, 4
 obese patients 10
 physical examination 6
- Care of the Critically Ill Surgical Patient (CCrISP) 122
- catheter over needle technique 61
- cell savers 31, 56
- central neural blockade 80
 complications 84–5, **84**
 hypotension and bradycardia 84
 nausea and vomiting 84–5

- post-dural puncture headache 85
 - epidural anaesthesia 81–3, 82, 84
 - spinal anaesthesia 83–4, 83
- central respiratory depression 102
- central venous cannulation 60–1, 61
 - equipment 61
 - internal jugular vein 60–1, **61**
 - subclavian vein 61
- central venous pressure 34, **34**, 111
- cephalic vein 60
- cerebral haemorrhage 102
- cerebral ischaemia 102
- cervical spine X-ray 8
- chest movement 125–6
- chest pain 147–9
 - angina 147
 - myocardial infarction 3, 148–9, 148
- chest X-ray 8
 - cardiogenic shock 145
 - pneumonia 137
 - pulmonary oedema 140
- chlorphenamine 94
- circle system 27–8, 27
- circulation
 - assessment 128–9, **129**
 - collapse 85–6, 93
- cisatracurium **45**
- clinical scoring systems 122–3, **123**
- coagulation screen 8
- coagulopathy 84
- cocaine 51
- colloids 54–5, **55**
- coma 129–30, **130**
 - see also* unconscious patients
- community-acquired pneumonia 137
- conscious level, decreased **129**, 154–7
- COX-1 inhibitors 49
- COX-2 inhibitors 49
- cricoid pressure 89, 89
- critical care outreach team 123, **124**
- cryoprecipitate 55
- crystalloids 53–4, **54**
- cyclizine **50**, 109

- dantrolene 95, 100
- desflurane **42**
- dexamethasone **50**, 109
- diabetes mellitus 5
- diaphragmatic splinting 102
- diffusion hypoxia 104
- disability 129–30, **129**
- dorsal metacarpal vein 60
- dose calculation 53
- drugs 39–57
 - dose calculation 53
 - history of use 5
 - see also specific drugs and classes*
- dyspnoea 132–41
 - Early Warning Score (EWS) 122, **123**
- ECG 8, 31
 - pulmonary embolism 141
 - pulmonary oedema 139
- echocardiography 9
 - transthoracic 143
- electrocardiogram *see* ECG
- electrolytes 8
- emergence from anaesthesia 76–7
- emergency surgery 88
- EMLA 40
- endocrine disorders, medical
 - referral 10
- entonox
 - cylinder colour **24**
 - postoperative analgesia 119
- epidural anaesthesia 81–3, 82
 - caesarean section 90
 - complications 117–19
 - post-dural puncture headache 85, 118–19
 - contraindications 84
 - inadequate 118
 - postoperative 116–17
- epinephrine *see* adrenaline
- Epworth sleepiness assessment **11**
- equipment
 - airway *see* airway equipment
 - cell savers 31
 - giving sets and fluid warmers 30
 - intravascular cannulas 30, 30
 - monitoring 36
 - patient warming 30–1
 - postanaesthesia care unit 100–1
 - syringe pumps 31
 - tracheal intubation 67
 - ultrasound 31
- etomidate **41**
- European Trauma Course (ETC) 122
- eutectic mixture of local anaesthetics
 - see* EMLA
- exercise tolerance 4–5

- facemasks 18–19, 19, 65, 65
 - oxygen delivery 104–5, 105
- failed intubation 96–8
- family history 5
- fentanyl **47**, 113
- fiberoptic bronchoscope 23, 23
- flowmeters 25–6, 25
- fluid therapy 108–11
 - clinical assessment 111
 - major surgery 110–11
 - minor surgery 108–10
 - monitoring 111
 - stress response 111
 - third space losses 110–11
- fluid warmers 30

180 Index

- fluids 53–6
 colloids 54–5, **55**
 crystalloids 53–4, **54**
 intraoperative 75–6
 accrued deficit 75
 requirements 75–6
see also blood/blood components
- fondiparinix 14
- fresh frozen plasma 55
- functional residual capacity 103
- gastric contents
 aspiration *see* aspiration
 full stomach 88
 pH/volume 39–40, 89
- gastric emptying, delayed 88
- gastrointestinal system, obese patients 11
- Gelofusine **55**
- general anaesthesia 58–78
 airway 63–70
 emergence from 76–7
 induction 62–3
 maintenance 70–2
 preoperative checks 58–9
 preparation for 59–62
 transfer to operating theatre 72–6
- giving sets 30
- Glasgow Coma Scale **130**
- glyceryl trinitrate (GTN), transdermal 40
- gum elastic bougie 24
- haemacel **55**
- haematological disorders, medical referral 10
- haemodilution, preoperative 56
- haemothorax 103
- Hartmann's solution **54**
- heart failure 3, 4
- helium/oxygen, cylinder colours **24**
- heparin 14
- hetastarch **55**
- hiatus hernia 5
- high airflow with oxygen enrichment 105, **105**
- hormone replacement therapy 5
- hospital-acquired pneumonia 137
- Hudson mask 105, **105**
- hydrocortisone 94
- hyoscine **50**
- hyperpyrexia, malignant 95
- hypertension 3–4
 postoperative 108
 and tracheal intubation 69
- hypoglycaemia 154–5
 investigations 154–155
 treatment 154
- hypopnoea syndrome 10–11
- hypotension 141–5
 causes **129**
 cardiogenic shock 144–5
 central neural blockade 84, 117
 hypovolaemic shock 84, 106, 142–3
 sepsis/septic shock 143–4
 postoperative 106–7
- hypothermia 103
- hypovolaemia/hypovolaemic shock 106,
 142–3, 157
 investigations 143
 treatment 142–3
- hypoxaemia 101–4, 157
 alveolar hypoventilation 101–3, **102**
 diffusion hypoxia 104
 management 104–6, **105**
 pulmonary diffusion defects 104
 reduced inspired oxygen concentration 104
 ventilation-perfusion mismatch 103–4, **104**
- hypoxia
 diffusion 104
 oesophageal intubation 69
 aspiration
 failed intubation
- induction of anaesthesia 62–3
 aspiration during 92–3
 rapid sequence 89
- infiltration analgesia 80–1, **80**
- informed consent 14–16
 definition 14
 evidence of 16
 obtaining 15–16
 patient information 14–15
 unconscious patients 16
- inhalational anaesthesia 42–3, **42**
 maintenance 70
 minimum alveolar concentration 43, **43**
 solubility 42–3
- internal jugular vein, cannulation 60–1, **61**
- intracranial haemorrhage 155–6, **155**
- intracranial pressure, raised 84
- intrathecal anaesthesia *see* spinal anaesthesia
- intravascular cannulas 30, **30**
- intravenous access 60–2
 arterial cannulation 62
 central venous cannulation 60–1, **61**
 complications **60**
- intravenous anaesthetic drugs 40–1, **41**
 total *see* total intravenous anaesthesia
- intubating laryngeal mask airway 20,
 20, **21**
- intubation *see* tracheal intubation
- investigations 2, 8–11
see also individual investigations and tests
- ipratropium 94
- isoflurane **42**
- jaundice 5
- ketamine **41**

- laryngeal mask airway (LMA) 19–20, 20
 - intubating 20, 20, 21
- laryngeal spasm 69
- laryngectomy 136
- laryngoscopes 22–4, 67
 - direct 22, 22
 - gum elastic bougie 24
 - indirect 22–3, 22, 23
- lateral position 73, 73
- latex allergy 94
- left ventricular dysfunction 106
- leg weakness 118
- levobupivacaine 53
- lidocaine 52, 53
 - infiltration analgesia 80–1
- lipid emulsions 100
- lithotomy position 73
- liver function tests 8
- Lloyd-Davies position 73
- local anaesthesia *see* regional anaesthesia
- local anaesthesia
 - role of 79–80
 - technique 80, 81
- local anaesthetic agents 50–3, 53
 - mechanism of action 50–1
 - overdose 85
 - toxicity 85–6
 - circulatory collapse 85–6
 - management of 85–86

see also individual drugs
- loss of consciousness *see* consciousness, reduced
- low urine output 145–147
 - investigations 146–147
 - treatment 146
- magnesium 94
- maintenance of anaesthesia 70–2
 - inhalational anaesthesia 70
 - total intravenous anaesthesia 70–1
- malignant hyperpyrexia
 - presentation 95
 - anaesthesia for susceptible patients 95
 - immediate management 95
 - investigation of family 95
- Mallampati criteria 7, 7
- mechanical ventilation 28, 71–2, 77
 - and cardiac output 103
- medical history 3–4
 - cardiovascular system 3–4, 4
 - respiratory system 4
- medical referral 9–10
- metabolic disorders, obese patients 11
- metoclopramide 40, 50
- midazolam 41
- minimum alveolar concentration (MAC) 43, 43
- Misuse of Drugs Act (1971) 48
- Misuse of Drugs Regulations (2001) 48
- mivacurium 45
- monitoring
 - equipment 36
 - fluid therapy 111
 - patient 31–6, 59–60
 - postanaesthesia care unit 100–1
 - potential hazards 59–60
 - regional anaesthesia 84

see also individual procedures
- morphine 47, 115
 - see also* opioids
- motor block 118
- multiple organ dysfunction syndrome (MODS) 144
- musculoskeletal system, physical examination 6
- myocardial infarction 3, 148–9
 - non-ST-segment-elevation (NSTEMI) 148
 - ST-segment-elevation (NSTEMI) 148–9, 148
- naloxone 113
- nasal cannulae 104–5, 105
- nasopharyngeal airway 19, 19, 64–5, 65
- National Confidential Enquiry into Perioperative Outcome and Death (NCEPOD) 13
- nausea and vomiting
 - central neural blockade 84–5
 - postoperative (PONV) 50, 108, 109
- needle cricothyroidotomy 97, 98
- nervous system, physical examination 6
- neuromuscular blocking drugs 44–6, 45
 - anticholinesterases 45–6
 - assessment of blockade 74–5
 - depolarizing 44–5
 - non-depolarizing 45

see also individual drugs
- neuromuscular disorders 5
- nitrous oxide 24–5, 42, 43
 - cylinder colour 24
 - systemic effects 43
- non-steroidal anti-inflammatory drugs *see* NSAIDs
- NSAIDs 49
 - contraindications 49
- nurses, preoperative assessment by 2
- obese patients 10–11, 10, 11
- obstetric patients 89–91
 - aorticaval compression 90–1
- obstructive sleep apnoea 10–11
- oesophageal Doppler cardiac output monitoring 34–5, 35
- oesophageal intubation, unrecognised 69
- oesophageal speech 134
- oliguria 106, 145–7
 - investigations 146–7
- omeprazole 40
- ondansetron 50, 109
- one-lung ventilation 91–2, 91
- opioids 46–9, 47, 113, 115
 - central and peripheral effects 47
 - long-term complications 113
 - narcosis 155

182 Index

- opioids (*Continued*)
 - overdose 113
 - pure agonists 46–8, 47
 - pure antagonists 48
 - regulation of 48–9
 - see also individual drugs*
- optical stylets 23, 23
- oral contraceptives 5
- oropharyngeal (Guedel) airway 19, 19, 63–4, 64
- oxygen 24
 - alveolar concentration 104
 - blood content 104
 - cylinder colour 24
 - inspired 123
 - reduced concentration 104
 - PaO₂ 102, 128
 - supply monitoring 36
- oxygen delivery 104–6, 105
 - fixed-performance devices 105–6, 105
 - variable-performance devices 104–5, 105
- oxygen masks 104–5, 105
- oxygen saturation 123
- PaCO₂ 102, 128
- PaO₂ 102, 128
- pain
 - acute 112–13, 113
 - experience of 112
 - and hypoxaemia 103
 - difficult problems 119
 - management *see* analgesia
- pancuronium 45
- paracetamol 40, 49
- parecoxib 49
- patient-controlled analgesia (PCA) 114–15
 - advantages and disadvantages 115
- patients
 - acutely ill *see* acutely ill patients
 - informed consent 14–15
 - monitoring 31–6, 59–60
 - see also individual procedures*
 - positioning 72–4
 - lateral 73, 73
 - prone 73–4, 74
 - supine 72–3, 72
 - preoperative check 58–9
 - sign in 59
 - warming 30–1, 70
- pericardial rub 128
- peripheral nerve blocks 115–16
- peripheral nerve stimulator 33
- pethidine 47
- physical examination
 - acutely ill patients 130
 - anaesthetic assessment 6–8
- platelet concentrates 55
- pneumonia 137
 - investigations 137
 - treatment 137
- pneumothorax 103, 138
 - investigations
 - monitoring
 - treatment
- pollution 29–30
- positive end-expiratory pressure (PEEP) 28
- positive pressure ventilation 71–2
- post dural puncture headache 85, 118–19
- post-anaesthesia care 100–21
 - analgesia 112–19
 - complications *see* postoperative complications
 - fluid therapy 108–11
- postanaesthesia care unit 100–1
 - discharge from 101, 101
- postoperative analgesia 113
- postoperative complications 101–8
 - hypertension 108
 - hypotension 106–7
 - hypoxaemia 101–4
- postoperative nausea and vomiting (PONV) 50, 108, 109
- pregnancy 6
- premedication 39–40
 - analgesia 40
 - anti-emetics 40
 - pH/volume of gastric contents 39–40
- preoperative assessment
 - anaesthetist-led 2–3
 - baseline investigations 2
 - investigations 2
 - nurse-led 2
- preoperative assessment clinic 2–3
- preoperative checks 58–9
 - anaesthetic machine 58
 - patient 58–9
 - sign in 59
- preoxygenation 62
- preparation for surgery 1–17
- pressure controlled ventilation (PCV) 28
- pressure support ventilation (PSV) 28
- prilocaine 52
- prochlorperazine 109
- prone position 73–4, 74
- propofol 41
 - total intravenous anaesthesia 70–1
- pruritus 118
- pseudocholinesterase deficiency 44–5
- pulmonary diffusion defects 104
- pulmonary embolism 140–1, 140
 - investigations
 - treatment
- pulmonary function tests 8
- pulmonary oedema 139–40
 - causes 139
 - monitoring
 - treatment
 - investigations 139–40, 140
- pulse 123

- pulse analysis cardiac output monitoring 35
- PiCCO[®]
- LiDCO[®]
- Flotrac[®]
- pulse oximetry 32, 127

- ranitidine 40
- rapid sequence induction 89
- recreational drugs 6
- red cell concentrates 55
- regional anaesthesia 79–86
 - awake vs. anaesthetised patients 86
 - caesarean section 89–90
 - monitoring 84
 - postoperative 115–19, 116
 - role of 79–80
 - techniques 80–4
 - see also specific blocks*
- remifentanyl 47
- renal disease
 - acute kidney injury 145
 - medical referral 9
- renal failure 5
 - pre-renal vs. intrinsic 146
- respiration rate 123
- respiratory depression 117–18
- respiratory disease
 - medical referral 9–10
 - signs of 127
 - see also individual conditions*
- respiratory system
 - history 4
 - obese patients 10–11, 11
 - physical examination 6
- rheumatoid disease 5
- Ringer's lactate 54
- risk indicators 12–13, 12
- rivaroxaban 14
- rocuronium 45
- ropivacaine 52, 53

- salbutamol 94
- saltatory conduction 51
- scavenging systems 29–30
- Schedule 2 drugs
 - supply and custody 48
- Seldinger technique 61
- Sellick's manoeuvre 89, 89
- sepsis/septic shock 143–4
 - treatment 144
- sevoflurane 42
- shock
 - cardiogenic 144–5
 - hypovolaemic 84, 106, 142–3
 - septic 143–4
- sickle-cell screen (sickledex) 8
- sign in 59
- sign out 76–7
- silent chest 134
- sinus bradycardia 107
- sinus tachycardia 107
- Situation Background Assessment Recommendation (SBAR) 122
- skin sepsis 84
- smoking 5–6
- social history 5–6
 - smoking
 - alcohol
 - drugs
 - pregnancy
- sodium chloride 0.9% 54
- sodium citrate, oral 40
- spinal anaesthesia 83, 83
 - caesarean section 90
 - contraindications 84
 - postoperative 119
- spontaneous ventilation 71, 76–7
- status epilepticus 156
 - treatment 156
 - investigations 156
- steroids 40, 94
- stress response 111
- stroke 155–6, 155
 - treatment 155
 - CT scan 156
- subclavian vein, cannulation 61
- sugammadex 46
- supine position 72–3, 72
- supraglottic airway devices 19–22, 20, 21, 66–7, 66, 76–7
 - aspiration with 93
 - insertion 66–7, 66
- operation, classification of 13
- suxamethonium 44
 - side effects 44
- syringe pumps 31
- systemic inflammatory response syndrome (SIRS) 143

- tachycardias 149–50
 - postoperative 107
- target controlled infusion 70
- temperature 123
 - hypothermia 103
 - malignant hyperpyrexia 95
 - monitoring 33–4
 - patient warming 30–1
- tension pneumothorax 138
 - treatment 138
- thiopental 41
- thoracotomy 91–2, 91
- thyromental distance 7, 8
- total intravenous anaesthesia (TIVA) 44
 - advantages 71
 - disadvantages 71
 - maintenance 70–1

184 Index

- tracheal intubation 67–9
 - complications 69
 - confirmation of position 68–9
 - difficult 95–6, 96
 - equipment 67
 - failed 96, 98
 - indications **67**
 - reflex activity 69
 - technique 67–8, 68
 - tracheal tubes 20–2, 21
 - tracheal tubes
 - reinforced
 - preformed
 - double lumen
 - uncuffed
- tracheostomy, blocked 133–4, 135
- track and trigger systems 122–3, **123**
- tramadol 47–8
- transfer to operating theatre 72–4
 - anaesthetic record 76
 - assessment of neuromuscular blockade 74–5
 - intraoperative fluids 75–6
 - patient positioning 72–4, 72, 73
 - time out 74
- transthoracic echocardiography 143
- transversus abdominis plane block 81, 81, 116
- trauma during tracheal intubation 69
- Trendelenburg position 73
- ultrasound 31
- unconscious patients 132
 - coma 129–30
 - Glasgow Coma Scale **130**
 - hypoglycaemia 154–5
 - hypoxaemia/hypovolaemia 157
 - informed consent 16
 - opioid narcosis 155
 - status epilepticus 156
 - stroke/intracranial haemorrhage 155–6, 155
- urea 8
- urinalysis 147
- urinary retention 118
- urine output
 - absent *see* anuria
 - low *see* oliguria
- vacuum 25
- vaporizers 26, 26
- vapour concentration analysis 33
- vasodilatation 107
- vecuronium **45**
- venous thromboembolism, prevention 13–14
- ventilation 100, 128
 - impaired mechanics 102
 - mechanical *see* mechanical ventilation
 - modes of 28
 - one-lung 91–2, 91
 - positive pressure 71–2
 - spontaneous 71, 76–7
 - see also* breathing systems
- ventilation-perfusion mismatch 103–4, **104**
- videolaryngoscopes 23
- Volulyte **55**
- Voluven **55**
- vomiting *see* nausea and vomiting
- warming of patients 30–1, 70
- Wilson score 7