

Index

• Symbols •

α (alpha)
controlling across important hypotheses, 75
defined, 42, 44
limiting chances for, 45
power affected by, 45
for sample-size calculation, 226
scaling sample size for, 373
spending strategy, 76
Type I error inflation, 74–75
usual level of, 46

* (asterisk)
multiplication indicated by, 21
powers indicated by (**), 22

β (beta), 42, 44

[] (brackets), multiplication indicated by, 21

^ (caret), powers indicated by, 22

{ } (curly braces), multiplication indicated by, 21

= (equal sign), equations indicated by, 26

γ (gamma) as skewness coefficient, 113

κ (kappa)
for Cohen's Kappa, 202
for Pearson kurtosis index, 114

– (minus sign), 20–21

() (parentheses), multiplication indicated by, 21

π (pi, lowercase) constant, defined, 19

Π (pi, uppercase) as array symbol, 29

+ (plus sign), addition indicated by, 20

· (raised dot), multiplication indicated by, 21

/ (slash), division indicated by, 22

|| (vertical bars), absolute value indicated by, 25

1-dimensional arrays, 26–27

2-dimensional arrays, 27, 28

3D charts, avoiding, 105

• A •

absence of harmful effects, 211–212

absolute values, indicating, 25

accuracy
defined, 38, 121
improving for measurements, 126–127
improving for sampling, 126
overall, 198, 281
precision versus, 38–39, 121
of sample statistic, 123
systematic errors affecting, 124–125

actuarial table. *See* life table

addition
mathematical operator for, 20
sums of array elements, 28–29

addresses, recording, 96

administrative details for a study, 70

adverse events, handling, 69

aims of a study, 61, 62

Akaike's Information Criterion (AIC), 277, 295, 352

Algebra I For Dummies (Sterling), 17, 30

Algebra II For Dummies (Sterling), 17

alignment charts (nomograms), 59–60, 170–171, 208

alpha (α)
controlling across important hypotheses, 75
defined, 42, 44
power affected by, 45
for sample-size calculation, 226
scaling sample size for, 373
setting a value for, 45
spending strategy, 76
Type I error inflation, 74–75
usual level of, 46

alternate hypothesis (H_1 or H_{Alt}), 41

alternative models, comparing, 297–298

analysis of covariance (ANCOVA), 158, 160–161

analysis of variance (ANOVA)

- basic idea of, 164–165
- for data grouped on different variables, 158
- for dissimilar standard deviations, 158
- interpreting output of, 166–168
- one-way, 157
- one-way, three-level, 157
- one-way, two-level, 157
- post-hoc tests, 165–166
- running, 166
- on summary data, 168–169
- for three or more groups, 157
- three-way, 158
- variance table, 166–167
- Welch unequal-variance, 158

analytical populations, 67, 69

analyzing your data. *See* data analysis and rule for probability, 32–33

animal studies, 79

antilogarithm (antilog), 24

apps, 48, 58

arithmetic mean, 107

arrays, 26–29

assay sensitivity, 214

asterisk (*)

- multiplication indicated by, 21
- powers indicated by (**), 22

attrition, sample-size estimation allowing for, 374

average value, 41. *See also* comparing averages

• B •

background information for a study, 68

balanced confidence interval, 135

bar charts, 105, 118–119. *See also* histograms

baseline hazard function, 342

baseline survival function, 342–344

beta (β), 42, 44

between-group difference, 200–201

bimodal distribution, 106, 108

binary logarithms, 23

binary variables, 189, 193–194, 254

binomial distribution, 37, 362

bio-creep, 218

bioequivalence, 211

bioequivalence studies (BE), 88, 212, 213, 214, 215

blinding, 65, 69

Bonferroni adjustment, 75

Bonferroni test, 165

Box plots, 119–120

box-and-whiskers (B&W) charts, 119–120

Box-Cox transformation, 117–118

brackets ([]), multiplication indicated by, 21

• C •

calculators

- cross-tabulated data for, 174
- for error propagation, 150–151
- for fourfold tables, 190
- for power, 48
- printed, 59–60
- scientific and programmable, 57
- web-based, 48, 58–59, 150–151, 190

calibrating instruments, 126–127

caret (^), powers indicated by, 22

Case Report Form (CRF), 63, 72

case-sensitivity, 20, 25

categorical data

- coding categories carefully, 96–97
- in data dictionary, 102
- graphing, 105
- multi-level categories, 97–98
- odds ratios for predictors, 286–287
- recording, 96–98
- summarizing, 104
- tabulating to check errors, 101

categorical variables

- cross-tabulation by, 104
- defined, 35
- for multiple regression, 254–255
- recoding as numerical, 255–256
- reference level, 255

cells and tissues, studies on, 79

censored data. *See also* estimating censored data

- example, 315–316
- reasons for, 315

- recording censoring information, 327–328
- techniques for handling, 316–317
- techniques not applicable to, 317–318
- Centers for Disease Control, 56
- centiles, 112, 329
- central limit theorem (CLT), 127, 137
- central tendency of data, 107–110
- certification, 72
- charts. *See* graphs and charts; tables
- chemical studies, 79
- chi-square distribution, 366–367
- chi-square test. *See* Pearson chi-square test
- CI. *See* confidence interval
- CL. *See* confidence interval (CI)
- clinical research. *See also* drug development
 - analyzing your data, 73–76
 - collecting and validating data, 72–73
 - conducting a study, 70–73
 - data collection and validation, 72–73
 - designing a study, 61–70
 - as focus of this book, 1, 11
 - protecting your subjects, 70–72
 - validating data, 72–73
- cloud-based software, 59
- clustered events, Poisson regression
 - with, 298
- coding data
 - categorical variables as numerical, 255–256
 - for gender, 96–97
 - for missing values, 98, 99
 - for multi-level categories, 97–98
 - numerical data, 99
- coefficient of determination, 247
- coefficient of variation, 111
- Cohen's Kappa (κ), 202
- collecting data, 72–73
- collinearity, 263–264, 285
- common logarithms, 23
- comparing averages
 - for data grouped on different variables, 158
 - for within-group changes between groups, 160–161
 - for matched sets, 159–160
 - mean to hypothesized value, 156
 - nuisance variable adjustment for, 158
 - situations requiring, 155–156
 - for three or more groups, 157
 - for two groups, 156–157
- comparing survival between groups.
 - See also* survival regression
 - in complicated situations, 337
 - Kaplan-Meier method for, 330
 - life-table method for, 330
 - log-rank test for, 332–336
 - sample-size estimation, 337–338, 371–372
- complete separation problem, 287–288
- computer software. *See also* software
 - case-level data with, 174
 - case-sensitivity of, 25
 - commercial, 52–55
 - demo versions, 52
 - development of, 51–52
 - free, 55–57
 - indicating factorials in, 24
 - for logistic regression, 274–275
 - for log-rank test, 333
 - for multiple regression, 254–258
 - for power calculation, 48
 - spreadsheet programs, 54
- concordance, 352
- confidence interval (CI)
 - around means, 137–139, 142
 - around proportions, 139
 - around regression coefficients, 141
 - assessing significance using, 141–142
 - balanced versus unbalanced, 135
 - calculating for rate ratio, 207, 208
 - for Cohen's Kappa, 202
 - described, 12, 40, 134
 - for equivalence and noninferiority, 215–217
 - formulas for large samples, 136–137
 - keeping as narrow as possible, 135
 - no-effect values, 141
 - normal-based, multipliers for, 137
 - one-sided, 136
 - p value compared to, 142
 - for precision of incidence rate, 205–206
 - as range of values, 134
 - standard error versus, 134
 - upper and lower limits of, 195

- confidence limit (CL). *See* confidence interval (CI)
- confidentiality, 70
- confounders (nuisance variables), 158, 187–188
- confounding, 65
- constants. *See also specific types*
 explicit versus symbolic representation of, 19
 in linear functions, 231
 standard error multiplied or divided by, 147
 standard error not changed by adding or subtracting, 146
- continuous variables, 35
- co-primary endpoints, 85
- correlation coefficient
 described, 193
 Pearson, 222–226
 straight-line regression, 247
 tetrachoric, 193–194
- correlation coefficient analysis
 precision of r value, 224–225
 r significantly different from zero, 223–224
 regression analysis versus, 239
 sample size required for test, 226
 significant difference in two r values, 225
- correlation, defined, 221, 222
- covariates for survival, 330
- Cox PH regression, 341–345
- CRF (Case Report Form), 63, 72
- crossover structure of a study, 64
- cross-tabulation. *See also* fourfold tables
 for comparing proportions, 174
 described, 173
 Fisher Exact test, 181–185
 fourfold tables, 173, 174, 189–202
 Kendall test, 185–187
 Mantel-Haenszel chi-square test, 187–188
 marginal totals or marginals, 173
 Pearson chi-square test, 174–181, 183–185, 186
 by two categorical variables, 104
- curly braces ($\{ \}$), multiplication indicated by, 21
- D •
- data
 collecting, 72–73
 file describing, 102
 interval, 94
 levels of measurement, 94–95
 numerical versus non-numerical, 4
 ordinal, 94
 privacy and confidentiality of, 70
 ratio, 94
 recording, 63, 95–101
 validating, 73, 101
- data analysis
 handling missing data, 74
 handling multiplicity, 74–75
 interim analyses, 76
 selecting analyses, 66–67
 Statistical Analysis Plan (SAP), 69
- data dictionary, 102
- data safety monitoring board or committee (DSMB or DSMC), 72
- dates and times, 99–101, 102
- deciles, 112
- degrees of freedom (df), 162, 179
- demo versions of software, 52
- designing a clinical study
 aims, objectives, hypotheses, and variables, 61–63
 analytical populations, 67, 69
 parallel versus crossover structure, 64
 protocol components, 68–70
 randomization, 64–66
 sample size, 67–68
 sample subject criteria, 63–64
 selecting analyses, 66–67
- diagnostic procedures, evaluating, 197–199
- dichotomous (binary) variables, 189, 193–194, 254
- difference table, 176–177
- dispersion, 110–112
- distributions
 bimodal, 106, 108
 binomial, 37, 362
 central tendency of, 107–110
 characteristics of, 106
 chi-square, 366–367

- for common test statistics, 359
 - dispersion of, 110–112
 - equations for, 37
 - exponential, 363
 - Fisher F, 367–368
 - kurtosis of, 113–114
 - log-normal, 37, 116–117, 157, 361
 - normal, 37, 106, 360–361
 - normalizing, 117–118
 - parametric functions for, 49
 - pointy-topped, 106, 114
 - Poisson, 37, 298, 362–363
 - population, 108
 - for random fluctuations in data, 359
 - representation of, 37
 - skewed, 106, 117–118
 - skewness of, 113
 - for statistical tests, 38
 - Student t distribution, 364–365
 - Student t test assumption about, 157
 - summary statistics showing, 103
 - uniform, 360
 - Weibull, 364
 - division, mathematical operators for, 22
 - dose-finding trials, 82–84
 - dose-limiting toxicities (DLTs), 80, 81
 - dose-response behavior, 83–84
 - dot, raised (\cdot), multiplication indicated by, 21
 - double-blinding, 65
 - double-indexing for arrays, 27
 - drug development. *See also* clinical research
 - bioequivalence studies, 88
 - pharmacokinetics/pharmacodynamics studies, 86–88
 - Phase I: Maximum tolerated dose, 80–82
 - Phase II: Safety and efficacy, 82–84
 - Phase III: Proving the drug works, 84–85
 - Phase IV: Monitoring the marketed drug, 85–86
 - preclinical studies, 78–79
 - reasons for looking at, 77
 - regulatory agencies, 80
 - steps in, 77–78
 - thorough QT trials, 88–90
 - DSMB or DSMC (data safety monitoring board or committee), 72
 - dummy variables, 255–256
 - Dunnett's test, 166
 - Dupont, W. D. (software developer), 56
- E •**
- e* constant, 2, 19, 22–23
 - effect size
 - for bioequivalence studies, 213
 - power affected by, 45, 46–47
 - for QT safety studies, 213
 - for sample-size calculation, 226
 - sample size relationship to, 47–48, 170–171
 - for therapeutic noninferiority studies, 213
 - effective doses on logistic curve, 280
 - effectiveness, 78
 - efficacy
 - co-primary endpoints, 85
 - defined, 78
 - drug testing for, 82–84
 - effectiveness versus, 78
 - noninferiority testing precaution, 217
 - popular usage of, 78
 - efficacy endpoints, 83
 - efficacy objectives, 62
 - elements of arrays, 28–29
 - environmental factors in precision, 125
 - Epi Info software, 56
 - epidemiology, 203
 - equal sign (=), equations indicated by, 26
 - equations, 17, 26. *See also* formulas
 - equivalence testing, 13, 214–217
 - equivalent functions, 305–306
 - error propagation
 - applying simple rules consecutively, 150
 - defined, 144
 - formulas for simple expressions, 146–149
 - online calculators for, 150–151
 - simulating, 152
 - error-checking for data, 73, 101
 - estimating censored data
 - interpreting survival curves, 328
 - Kaplan-Meier method for, 324–325
 - life-table method for, 318–324
 - recording censoring information correctly, 327–328
 - recording survival times correctly, 327

event counts or rates
 calculating observed and expected counts, 175–177
 comparing two counts with identical exposure, 209
 comparing two rates, 107–109
 confidence interval around, 140
 Poisson regression for, 291–298
 standard error of, 130–131
ex vivo studies, 79
 Excel software
 data-recording cautions, 97, 99
 error-checking techniques, 101
 PopTools add-in for, 56
 randomization using, 66
 uses for biostatistics, 54
 excess kurtosis, 114
 exclusion criteria, 64, 126
 explicit representation of constants, 19
 exploratory objectives, 62
 exponential distribution, 363
 exponentiation, 22–23, 30
 exposure, 204
 expressions, 17. *See also* formulas

● F ●

F statistic, 247–248, 260
 F test for equal variances, 164
 factorials, 24
 false discovery rate (FDR), 75
 false negative results, 197–198, 282–283
 false positive results, 197–198, 282–283
 FDA Adverse Event Reporting System (FAERS), 86
 feedback to the author, 1
 Fisher Exact test, 181–185
 Fisher F distribution, 367–368
 Fisher, R. A. (statistician), 181
 Fisher z transformation, 224, 225
 Fisher's LSD test, 166
 fitted logistic formula, 278–280
 Food and Drug Administration (FDA), 71, 80, 84–85, 86
 formulas
 arrays in, 28
 defined, 17

hierarchy rules for, 25–26
 typeset versus plain text format, 18
 usage in this book, 17
 fourfold tables. *See also* cross-tabulation
 for assessing risk factors, 194–197
 for comparing proportions, 174
 conventions in this book for, 190–191
 defined, 173, 189
 for evaluating diagnostic procedures, 197–199
 Fisher Exact test, 181–185
 fundamentals of, 190–191
 for inter- and intra-rater reliability, 201–202
 for investigating treatments, 199–201
 Mantel-Haenszel chi-square test, 187–188
 Pearson chi-square test, 174–181, 183–185, 186
 sample statistic from, 190
 sampling strategies for, 191–192
 scenarios leading to, 192
 for tetrachoric correlation coefficient, 193–194
 free software, 48, 55–59
 free-text data, recording, 95
 frequency bar charts, 105
 frequency distributions. *See* distributions
 Friedman tests, 158, 159
 functions
 defined, 25
 equivalent, 305–306
 linear, 231
 nonlinear, 231, 299
 parametric distribution, 49
 S-shaped, for logistic regression, 270–273
 using in formulas, 25

● G ●

Galton, Francis (statistician), 222
 gamma (γ) as skewness coefficient, 113
 gender, 2, 96–97
 generalized linear model (GLM), 292–293
 Generalized Savage's test. *See* log-rank test
 geometric mean (GM), formulas for, 109–110
 gestational age, recording, 99
 goals of a study, 61, 62
 good clinical practice (GCP), 72

good doses, defined, 84
 goodness-of-fit indicators, 247–248, 351–352
 G*Power software, 57
 GraphPad InStat software, 54
 GraphPad Prism software, 53–54
 graphs and charts. *See also* scatter plots;
 tables
 bar charts, 105, 118–119
 box-and-whiskers charts, 119–120
 of categorical data, 105
 hazard rates and survival probabilities, 324
 histograms, 35, 116–118
 logistic regression data, 269–270
 nomograms (alignment charts), 59–60,
 170–171, 208
 of numerical data, 115–120
 pie charts, 105
 of probability distributions, 37
 Receiver Operator Characteristics,
 283–285
 of relationships between numerical
 variables, 120
 residuals versus fitted, 242–243, 249
 S-shaped function for logistic regression,
 270–273
 3D charts, avoiding, 105

• H •

H_0 (null hypothesis), 41, 43–45
 H_1 or H_{Alt} (alternate hypothesis), 41
 hazard rate
 defined, 316
 estimating censored data, 318–328
 graphing from a life table, 324
 interpreting life-table output, 323
 hazard ratios, 345, 350–351
 hierarchical testing strategy, 75
 hierarchy rules for formulas, 25–26
 histograms, 35, 116–118
 historical control, comparing results to, 156
 Hood, Greg (ecologist), 56
 Hosmer-Lemeshow (H-L) test, 277
 human error, 124
 hypotheses, identifying, 62
 hypothesis testing, 41–43

• I •

icons in this book, explained, 5–6
 identification (ID) numbers, recording,
 95–96
 imprecision. *See* precision
 imputing values, 74, 317
in silico studies, 79
in vitro studies, 79
in vivo studies, 79
 inaccuracy. *See* accuracy
 incidence, 204
 incidence rate (R), 204–207
 inclusion criteria, 64, 126, 128
 index of an array, 27
 Informed Consent Form (ICF), 71–72
 inner mean, 109
 Institutional Review Boards (IRBs), 71, 72, 76
 intent-to-treat (ITT) population, 67
 inter- and intra-rater reliability, 201–202
 intercept, 231, 244
 interim analyses, 76
 inter-quartile range (IQR), 112
 interval data, 94, 100–101, 314
 interval of uncertainty (IOU), 145

• K •

Kaplan-Meier (K-M) method, 324–328
 kappa (κ)
 for Cohen's Kappa, 202
 for Pearson kurtosis index, 114
 Kendall test, 185–187
 Krukall-Wallis test, 158
 kurtosis, 113–114

• L •

Last Observation Carried Forward
 (LOCF), 74
 LazStats software, 55
 least-squares regression, 291–292. *See also*
 nonlinear regression
 leptokurtic distribution, 114
 lethal doses on logistic curve, 280

life table
 basic idea of, 318–319
 creating, 319–323
 graphing hazard rates and survival
 probabilities from, 324
 guidelines for, 326–328
 interpreting, 319–323
 Likert agreement scale, 94, 97
 linear, defined, 252
 linear functions, 231
 linear regression. *See* multiple regression;
 straight-line regression
 logarithms, 23–24
 logistic regression
 assessing model adequacy, 276–277
 calculations for, 273
 collinearity in, 285
 complete separation problem, 287–288
 described, 14, 230
 fitted logistic formula, 278–280
 gathering and graphing data for, 268–270
 interpreting output of, 275–285
 issues to beware of, 285–288
 with multiple predictors, 274
 for nonlogistic data, avoiding, 285
 odds ratios for categorical predictors,
 286–287
 odds ratios for numerical predictors, 286
 Receiver Operator Characteristics curve,
 283–285
 reverse-coding of outcome variable, 286
 running software for, 274–275
 sample-size estimation, 288–289
 S-shaped function for data, 270–273
 steps for, 274–275
 summary information about variables, 276
 table of regression coefficients, 278
 typical categorical outcomes of, 267
 typical output of, 275, 276
 uses for, 267–268
 for yes or no predictions, 280–285
 log-normal distribution, 37, 116–117, 361
 log-normal transformation, 117
 log-rank test
 assumptions about data in, 336
 basic idea of, 333

calculations for, 334–336
 data for, 332
 extended versions of, 331
 limitations of, 340
 other names for, 331
 running on software, 333
 LOWESS curve-fitting, 14
 LOWESS regression, 306–310

• M •

Mann-Whitney (M-W) test, 157
 Mantel-Cox test. *See* log-rank test
 Mantel-Haenszel chi-square test, 187–188
 Mantel-Haenszel test for survival data. *See*
 log-rank test
 margin of error, 137, 144
 marginal totals or marginals, 173, 176
 masking (blinding), 65
 matched numbers, comparing, 159–160
 matched-pair data, comparing, 159
 mathematical operations, 20–25
 matrices, 28. *See also* arrays
 maximum tolerated dose (MTD), 81–82
 mean
 arithmetic, 107
 comparing to hypothesized value, 156
 confidence interval around, 137–139, 142
 geometric, 109–110
 inner, 109
 root-mean-square, 110
 of sample versus population, 36
 standard error of, 129–130
 measurement
 calibrating instruments for, 126–127
 imprecision sources in, 125
 improving accuracy of, 126–127
 improving precision of, 128
 indirect, 143
 as kind of sampling, 123–124
 systematic errors in, 124
 median, 107–108
 metadata, 102
 Miller, Bill (software developer), 55
 Minitab software, 55
 minus sign (–), 20–21

- missing data
 - censored data versus, 317–318
 - coding, 98, 99
 - in data dictionary, 102
 - dates, 100
 - methods for handling, 74
 - mixed-model repeated measures (MMRM), 74
 - mobile devices, software for, 58
 - mode, sample versus population, 108
 - model building problem, 264
 - Monte-Carlo analysis, 152
 - MTD (maximum tolerated dose), 81–82
 - multicollinearity, 263–264, 285
 - multiple regression
 - basic idea of, 251–252
 - categorical variables for, 254–256
 - collinearity in, 263–264
 - dummy variables for, 255–256
 - fit of model to data, 262
 - interpreting output of, 258–262
 - matrix notation for, 252
 - for model building problem, 264
 - optional output for, 260
 - recoding categorical variables as numerical, 255–256
 - reference level for, 255
 - running software for, 254–258
 - sample-size estimation, 265
 - scatter plots for, 256–258
 - solving simultaneous equations for, 253
 - steps for, 258
 - suitability of data for, 261
 - synergistic effect in, 263
 - terms and notation used in, 252–253
 - typical output of, 259–260
 - uses for, 253
 - multiplication
 - mathematical operators for, 21
 - products of array elements, 28–29
 - standard error for products, 148
 - multiplicity, 74–75
 - multivariable linear regression, 14
 - multivariable regression, 230
 - multivariate regression, 229
 - M-W (Mann-Whitney) test, 157
- N •
- names, recording, 96
 - National Institutes of Health (NIH), 71
 - natural logarithms, 23, 24
 - negative numbers, indicating, 21
 - negative predictive value (NPV), 199
 - NNT (number needed to treat), 201
 - nominal variables, 94
 - nomograms (alignment charts), 59–60, 170–171, 208
 - noninferiority creep, 218
 - noninferiority testing, 13, 214–218
 - nonlinear functions, 231, 299
 - nonlinear least-squares regression, 14
 - nonlinear regression
 - drug research example, 300–302
 - interpreting output of, 304–305
 - as least-squares regression, 291
 - nonlinear functions, 299
 - other regressions versus, 299–300
 - running, 302–303
 - using equivalent functions to fit parameters, 305–306
 - nonlinear trends, Poisson regression for, 296–297
 - nonparametric regression, 306
 - nonparametric tests, 49, 50, 169
 - nonsuperiority, confidence intervals for, 217
 - normal distribution, 37, 106, 157, 360–361
 - normality assumption, 157
 - not* rule for probability, 32
 - NPV (negative predictive value), 199
 - nuisance variables, 158, 187–188
 - null hypothesis (H_0), 41, 43–45
 - null model, 248
 - number needed to treat (NNT), 201
 - numerical data
 - in data dictionary, 102
 - in Excel, 99
 - graphing, 115–120
 - grouping in intervals, avoiding, 98
 - recording, 98–99
 - structuring summaries in tables, 114–115
 - summarizing, 106–114
 - numerical variables, 98–99, 255–256

• 0 •

objectives of a study, 62
 odds, 33–34, 195
 odds ratio (OR), 188, 195–197, 286–287
 Office for Human Research Protections (OHRP), 71
 one-dimensional arrays, 26–27
 one-sided confidence interval, 136
 OpenStat software, 55
 operator variability, 125
 or rule for probability, 33
 order of precedence for formulas, 25–26
 ordinal data, 94, 185–187. *See also*
 categorical data
 ordinary, defined, 252
 ordinary multiple linear regression. *See*
 multiple regression
 ordinary regression, 230
 outliers, 111–112, 249
 overall accuracy, 198, 281

• p •

p value
 chi-square test for determining, 178–179
 confidence interval compared to, 142
 defined, 42
 Fisher Exact test for, 181–183
 logistic regression, 277
 multiple regression, 260
 significance in reference to, 2, 190
 straight-line regression, 246
 survival regression, 350
 as a test result, 43
 paper calculators, 59–60
 parallel structure of a study, 64
 parameter error-correlations matrix, 260
 parametric distribution functions, 49
 parametric tests, 49, 50
 parentheses [()], multiplication indicated
 by, 21
 Pearson chi-square test
 basic formulas for, 180
 calculating observed and expected
 counts, 175–177

cell naming and notation, 179
 degrees of freedom for, 179
 determining the p value, 178–179
 development of, 174
 Kendall test versus, 186
 power and sample size calculation for,
 183–185
 pros and cons of, 180
 summarizing and combining scaled
 differences, 177–178
 Yates continuity correction for, 181, 184
 Pearson correlation coefficient
 described, 222
 precision of r value, 224–225
 r significantly different from zero, 223–224
 sample size required for test, 226
 significant difference in two r values, 225
 in straight-line regression, 247
 for straight-line relationships only, 223
 Pearson, Karl (statistician), 174, 222
 Pearson kurtosis index, 114
 percentages, 32, 104. *See also* proportions
 per-protocol (PP) population, 67
 person-time data. *See* event counts or rates
 pharmacokinetics/pharmacodynamics
 (PK/PD) studies, 86–88, 300–302
 physical factors in precision, 125
 pi, lowercase (π) constant, defined, 19
 pi, uppercase (Π) as array symbol, 29
 pie charts, 105
 pivotal Phase III studies, 85
 placebo effect, 65
 plain text format
 absolute values in, 25
 arrays in, 27, 29
 division in, 22
 logarithms in, 23
 multiplication in, 21
 powers in, 22
 roots in, 23
 scientific notation in, 30
 typeset format compared to, 18
 platykurtic distribution, 114
 Plummer, W. D. (software developer), 56
 plus sign (+), addition indicated by, 20
 pointy-topped distribution, 106, 114
 Poisson distribution, 37, 298, 362–363

- Poisson regression
 with clustered events, 298
 for comparing alternative models, 297–298
 described, 14, 230
 interpreting output of, 295–296
 for nonlinear trends, 296–297
 running, 293–294
 with unequal observation intervals, 298
- PopTools software, 56
- population parameter, 122
- populations
 analytical, 67, 69
 as confidence interval focus, 134
 defined, 36
 mode for distribution of, 108
- positive control, 214, 217
- positive predictive value (PPV), 199
- post-hoc tests for ANOVA, 165–166
- power of a test
 calculating for chi-square and Fisher
 Exact tests, 183–185
 defined, 42
 effect size relationship to, 45, 46–48
 factors affecting, 45
 methods for calculating, 48
 for sample-size calculation, 226
 sample size relationship to, 45, 46, 47–48
 scaling sample size for, 372
- powers, 22–23, 149
- PP (per-protocol) population, 67
- precedence order for formulas, 25–26
- precision
 accuracy versus, 38–39, 121
 defined, 38, 121
 improving for measurements, 128
 improving for sampling, 127–128
 of incidence rate, 205–206
 measurement imprecision, 125
 of numerical data, 98–99
 of *r* value, 224–225
 random errors affecting, 125
 of sample statistic, 123
 sample-size estimation for, 132
 sampling imprecision, 125
- preclinical studies, 78–79
- predictors, 230, 274, 286–287
- prevalence, 204, 205
- primary objectives, 62
- printed calculators, 59–60
- privacy considerations, 70
- probability
 and rule for, 32–33
 defined, 31
 not rule for, 32
 numbers between 0 and 1 for, 32
 odds compared to, 33–34
 or rule for, 33
 percentages for, 32
- probability distributions. *See* distributions
- procedural descriptions, 69
- product description for a study, 69
- programmable calculators, 57
- programs. *See* software
- propagation of errors. *See* error propagation
- proportional-hazards regression, 347–348, 353–356
- proportions
 confidence interval around, 139
 fourfold table for comparing, 174
 between-group difference, 200–201
 prevalence, 204, 205
 standard error of, 130
- protocol components of a study, 68–70
- PS software, 56–57
- pseudo R-square values, 277
- **Q** •
- QT interval, 89
- QT safety studies, 212, 213, 214, 215
- QT/QTc (thorough QT) trials, 88–90
- quartiles, 112
- quintiles, 112
- **R** •
- r*. *See* Pearson correlation coefficient
- R (incidence rate), 204–207
- R software, 56, 295
- raised dot (\cdot), multiplication indicated by, 21

- randomization
 - advantages of, 65
 - of data versus numbers, 34
 - imprecision from errors in, 125
 - prediction not possible with, 35
 - in study protocol, 69
 - using in a study, 64–66
 - randomized controlled trials (RCTs), 64–66
 - range of a set of values, 111–112
 - rate ratio (RR), 206–207, 208
 - ratio data, 94
 - rationale for a study, 68
 - ratios, standard error for, 148
 - Receiver Operator Characteristics (ROC)
 - curve, 283–285
 - recording data
 - Case Report Form (CRF) for, 63
 - categorical data, 96–98
 - censoring information, 327–328
 - dates and times, 99–101
 - in Excel, 97, 99
 - free-text data, 95
 - ID numbers, 95–96
 - names and addresses, 96
 - numerical data, 98–99
 - simple method for, 63
 - survival times, 327
 - reference level for multiple regression, 255
 - regression analysis. *See also specific types*
 - correlation coefficient analysis versus, 239
 - generalized linear model, 292–293
 - linear versus nonlinear functions for, 231
 - logistic regression, 14, 230, 267–289
 - LOWESS regression, 306–310
 - multiple regression, 251–265
 - nonlinear regression, 298–306
 - number of outcomes, 229
 - number of predictors, 230
 - outcome variable's type of data, 230–231
 - Poisson regression, 14, 230, 291–298
 - purpose of, 227–228
 - software needed for calculations, 236
 - straight-line regression, 233–250
 - survival regression, 339–356
 - terms and notation used in, 228–229
 - types of, 14, 229–232
 - univariable versus multivariable, 230
 - univariate versus multivariate, 229
 - regression coefficients
 - confidence interval around, 141
 - standard error of, 131–132, 244–245, 350
 - survival regression, 350
 - regression, defined, 221
 - regression table
 - importance of, 243
 - for logistic regression, 278
 - for multiple regression, 260
 - for straight-line regression, 244–246
 - regulatory agencies, 71, 80
 - relative risk (RR), 194–195
 - Remember icon, 5
 - repeated-measures analysis of variance (RM-ANOVA), 159–160
 - residual standard error, 242, 260
 - residuals, 241–243, 259, 260
 - residuals versus fitted graph, 242–243, 249
 - risk evaluation mitigation strategy (REMS), 86
 - risk factor assessment
 - odds ratio, 195–197
 - relative risk (RR), 194–195
 - risk ratio, assessing, 194–195
 - ROC (Receiver Operator Characteristics)
 - curve, 283–285
 - root-mean-square (RMS), 110
 - root-mean-square (RMS) error, 242
 - roots, 23, 149
 - RR (rate ratio), 206–207
 - RR (relative risk), 194–195
 - Rumsey, Deborah J. (*Statistics For Dummies*), 3
- S ●
- s. *See* standard deviation
 - S Plus software, 55
 - safety considerations
 - certification in human subjects
 - protection, 72
 - indicators for drugs, 83
 - Informed Consent Form, 71–72
 - Institutional Review Boards, 71
 - monitoring board or committee, 72
 - Phase II in drug testing, 82–84
 - protecting your subjects, 70–72
 - regulatory agencies, 71

- safety objective, 62
- safety population, 67, 69
- in study protocol, 69
- sample
 - defined, 36
 - determining suitable subjects, 63–64
 - exclusion criteria, 64, 126
 - fourfold table strategies, 191–192
 - imprecision sources in, 125
 - improving accuracy of, 126
 - improving precision of, 127–128
 - inaccuracy in, 124
 - inclusion criteria, 64, 126, 128
 - measurement as, 123–124
 - perfection not possible for, 36
 - randomization of, 64–66
 - selection bias, 65
 - as standard error focus, 134
- sample size
 - effect size relationship to, 47–48, 170–171
 - power affected by, 45, 46
 - standard error affected by, 127
 - in study protocol, 69
- sample statistic, 122, 123, 190
- sample-size estimation
 - to achieve precision desired, 132
 - adjusting for unequal group sizes, 373–374
 - allowing for attrition, 374
 - Cheat Sheet for, 48
 - for clinical study, 67–68
 - for comparing averages, 169–171
 - for comparing means, 370
 - for comparing paired values, 370
 - for comparing proportions, 371
 - for comparing survival, 337–338, 371–372
 - for correlation test, 226
 - for Fisher Exact test, 183–185
 - free software for, 56–57
 - importance of, 15
 - for logistic regression, 288–289
 - for multiple regression, 265
 - nomograms for, 170–171
 - for noninferiority testing, 218
 - for Pearson chi-square test, 183–185
 - for rate comparisons, 209–210
 - for scaling from 0.05 to another alpha level, 373
 - scaling from 80% to another power, 372
 - for significant correlation, 371
 - simple formulas for, 169
 - software for, 170
 - for straight-line regression, 249–250
 - for survival regression, 356
 - web-based calculators for, 170
- SAP (Statistical Analysis Plan), 69
- SAS software, 52–53
- scatter plots
 - LOWESS regression, 306–310
 - multiple regression, 256–258, 260
 - straight-line regression, 234, 235, 238–239
- Scheffe's test, 166
- scientific calculators, 57
- scientific notation, 30
- Score Test. *See* log-rank test
- SD or sd. *See* standard deviation
- SE. *See* standard error
- secondary objectives, 62
- selection bias, 65
- sensitivity of a test, 198, 282
- significance
 - defined, 42
 - with multicollinearity, 263–264, 285
 - in reference to p value, 2, 190
 - statistical, 43
 - testing for, 42–43
- significance tests, 42, 214–215
- simple linear regression. *See* straight-line regression
- simulating error propagation, 152
- single-blinding, 65
- skewed distribution, 106, 113, 117–118
- skewness coefficients, 113
- slash (/), division indicated by, 22
- slope, 131–132, 244
- smoothing fraction for LOWESS, 309–310
- software. *See also specific types*
 - apps, 48, 58
 - case-level data with, 174
 - case-sensitivity of, 25
 - cloud-based systems, 59
 - computer, 48, 51–57
 - field-length limitations of, 95
 - indicating factorials in, 24
 - for logistic regression, 274–275

- software (*continued*)
- for log-rank test, 333
 - for multiple regression, 254–258
 - for power calculation, 48
 - required for regression, 236
 - for sample-size estimation, 170
 - sources of, 10
 - web-based, 48, 58–59
- specificity of a test, 198–199, 282
- sphericity, RM-ANOVA issue with, 160
- spreadsheet programs, 54, 101. *See also*
- Excel software
 - SPSS software, 53
- square-root transformation, 117
- standard deviation (SD, sd, or s)
- formula for, 110
 - of Poisson distribution, 298
 - of population distributions, 111
 - of sample versus population, 36
 - sensitivity to outliers, 111
- Student t test assumption about, 157
- standard error (SE). *See also* error propagation
- of average of N equally precise numbers, 148
 - confidence interval versus, 134
 - described, 12, 40, 125
 - of differences, 177
 - estimating for single-variable expressions, 144–145
 - of event counts and rates, 130–131
 - of means, 129–130
 - multiplying or dividing by a constant, 147
 - not changed by adding or subtracting a constant, 146
 - in Poisson regression, 295
 - for powers and roots, 149
 - for products and ratios, 148
 - of proportions, 130
 - of regression coefficients, 131–132, 244–245, 350
 - of residuals, 242, 260
 - sample size affecting, 127
 - sample-size estimation using, 132
 - Student t test calculation of, 162
 - when adding or subtracting two measurements, 147
- Stata software, 55
- statistic, defined, 42
- Statistical Analysis Plan (SAP), 69
- statistical decision theory, 12–14, 40–41
- statistical estimation theory, 12, 38–40
- statistical inference, 38
- Statistics For Dummies* (Rumsey), 3
- Sterling, Mary Jane
- Algebra I For Dummies*, 17, 30
 - Algebra II For Dummies*, 17
- straight-line regression
- conditions suitable for, 234–235
 - correlation coefficient, 247
 - error sources in, 249
 - example, 237–239
 - F statistic, 247–248
 - formulas for, 235–236
 - gathering data for, 237–238
 - goodness-of-fit indicators, 247–248
 - intercept, 244
 - interpreting output of, 239–248
 - other names for, 233
 - p value, 246
 - prediction formula, 248
 - regression table for, 243–246
 - residuals from, 241–243
 - sample-size estimation for, 249–250
 - scatter plot creation, 238–239
 - slope, 244
 - software needed for calculations, 236
 - standard error of coefficients, 244–245
 - steps for, 237
 - Student t value, 246
 - t value, 246
 - typical output of, 240
 - variables for, 233
- stratification, 187–188
- stratified Cochran-Mantel-Haenszel test.
- See* log-rank test
- Student t distribution, 364–365
- Student t tests
- assumptions about data in, 157
 - basic idea of, 161–162
 - calculation of difference, standard error, and degrees of freedom by, 162
 - for comparing matched pairs, 159

- for comparing mean to hypothesized value, 156
 - for comparing two groups, 157
 - F test for equal variances, 164
 - interpreting output of, 163–164
 - for matched-pair data, 159
 - one-group, 156, 162
 - paired, 159, 163
 - running, 162–163
 - on summary data, 168–169
 - unpaired or independent sample, 157, 163
 - Welch, 157, 162–163
 - subject-to-subject variability, 125
 - subtraction, 20
 - summary statistics
 - for categorical data, 104
 - defined, 103
 - for numerical data, 106–114
 - for residuals, 241–242
 - structuring in tables, 114–115
 - Student t and ANOVA tests with, 168–169
 - superscripting, powers indicated by, 22
 - survival analysis
 - applicability to other events, 313, 331, 339
 - comparing survival times, 330, 331–338
 - custom prognosis chart, 330
 - determining factors affecting survival, 330
 - estimating censored data, 318–328
 - five-year (or other time) survival rate, 329
 - median (or centile) survival time, 329
 - survival regression, 231, 339–356
 - survival data. *See also* estimating censored data
 - censoring of, 315–318
 - estimating censored data, 318–328
 - interpreting survival curves, 328
 - as interval data, 314
 - non-normality of, 314–315
 - overview, 14–15
 - techniques not applicable to, 314–315, 317–318
 - survival rate
 - defined, 316
 - estimating censored data, 318–328
 - five-year (or other time), 329
 - graphing from a life table, 324
 - interpreting life-table output, 323
 - survival regression
 - baseline hazard function, 342
 - baseline survival function, 342
 - with censored data, 341
 - concepts behind, 340–345
 - constructing prognosis curves, 353–356
 - Cox PH regression, 341–345
 - described, 231
 - goodness-of-fit indicators, 351–352
 - hazard ratios, 345, 350–351
 - interpreting output of, 347–352
 - linear combination for, 340–341
 - with nonparametric versus parametric survival curves, 341
 - proportional-hazards regression, 347–348, 353–356
 - running, 346–347
 - sample-size estimation, 356
 - table regression coefficients, 350
 - testing validity of assumptions, 349
 - uses for, 340
 - survival times
 - comparing between groups, 330, 331–337
 - median (or centile), 329
 - recording correctly, 327
 - symbolic representation of constants, 19
 - synergistic effect, 263
 - systematic errors, 124–125
- T ●
- t value, 246, 260
 - tables. *See also* fourfold tables; graphs and charts
 - life table, 318–328
 - for power calculation, 48
 - of probability distributions, 37
 - of regression coefficients, 60, 243–246, 278
 - structuring numerical summaries in, 114–115
 - Technical Stuff icon, 2, 5
 - terachoric correlation coefficient, 193–194
 - test statistic, 42–43
 - tetrachoric correlation coefficient, 193–194
 - theoretical molecular studies, 79
 - therapeutic noninferiority, 211

therapeutic noninferiority studies, 212, 213, 214, 215
 therapeutic range for a drug, 84
 thorough QT (TQT or QT/QTc) trials, 88–90
 3D charts, avoiding, 105
 tied values, 50
 times and dates, 99–101, 102
 time-to-event data. *See* survival data
 Tip icon, 6
 title of a study, 68
 Torsades de Points (TdP), 88
 trimmed mean, 109
 Tukey-Kramer test, 166
 Tukey's HSD test, 166
 two-dimensional arrays, 27, 28
 two-way tables. *See* cross-tabulation
 Type I error, 42, 44, 45, 74–75
 Type II error, 42, 44

• U •

unary operator, 21
 unbalanced confidence interval, 135
 unequal observation intervals, Poisson regression with, 298
 unequal-variance t (Welch t) test, 157, 163–164
 uniform distribution, 360
 univariable regression, 230
 univariate regression, 229

• V •

validating data, 73, 101
 variables
 case of, 20
 categorical, 35, 104, 254–256
 continuous, 35
 in data dictionary, 102
 for dates and times, 99–100
 defined, in different fields, 19–20

free-text, 95
 identifying for a study, 63
 for names and addresses, 96
 nominal, 94
 numerical, 98–99, 255–256
 sorting values to check errors, 101
 unary operator for, 21
 variance, 111
 variance table from ANOVA, 166–167
 vectors, 27. *See also* arrays
 vertical bars (| |), absolute value indicated by, 25

• W •

Warning! icon, 6
 web resources, 59
 web-based software
 calculators for error propagation, 150–151
 calculators for power, 48
 defined, 58
 for fourfold tables, 190
 for sample-size estimation, 170
 statistics packages, 59
 Weibull distribution, 364
 Welch t test, 157, 163–164
 Welch unequal-variance ANOVA, 158
 Wilcoxon Signed-Ranks (WSR) test, 156, 159
 Wilcoxon Sum-of-Ranks (WSOR) test, 157
 withdrawal criteria, 64
 within-subject variability, 125

• Y •

Yates continuity correction, 181, 184
 yes or no predictions, 280–285

• Z •

zero point, 94