

Index

| abortion, spontaneous see miscarriage | allium 260 |
|--|---|
| absorption see digestion and absorption; malabsorption | alveolar cells (breast) 124 |
| acetaldehyde and drinking 210, 211, 253 | Alzheimer's disease xvi |
| acute phase (inflammatory) response 5 | anaemia and 293, 294 |
| additives (food) and hyperactivity 161–2 | ApoE gene and 13 |
| adiponectin xvi, 39, 234 | Ames test xvi, 254–5 |
| adiposity rebound xvii, 134, 174 | amino acids 304–5 |
| adolescence (teenage years) 189–218 | essential amino xix, 305 |
| nutritional intake 199–202 | formulas feeds 148 |
| nutritional problems 203–14 | structure 305 |
| nutritional requirements 197–9 | anaemia 290–2 |
| physical development 189–96 | elderly 290–4 |
| growth see growth | iron deficiency see iron deficiency anaemia |
| pregnancy 209–10 | pernicious xxiii, 56, 292 |
| high-risk 74 | android pattern of body fat distribution 223 |
| psychosocial development 196 | anencephaly xvi, 54 |
| adrenal glands xvi | animals |
| adrenarche xvi, 191, 192, 193 | experimental studies |
| adulthood 219–67 | caloric restriction and lifespan 273, 274 |
| fetal nutrition related to disease in see programming | maternal nutrition and fetal programming 103–4, 112–13 |
| guidelines for healthy nutrition 220–3 | protein availability from, by world region 15, 16 |
| infant/preschool child nutrient intake and transition | anorexia nervosa (AN) xvii, 204, 206–7 |
| to patterns in 156–67 | bone growth and 195 |
| nutritional requirements/needs 11, 219–20 | drug abuse and 214 |
| compared to adolescents/11-18 year olds 197 | anovulation xvi, 37, 39 |
| compared to infant/perschool children 152 | anthropometry xvi, 14, 223 |
| Adventists, cancer 247 | birth, and disease in later life 100, 116 |
| advertising, food and beverage 165–7 | antibodies see immunoglobulins |
| aflatoxins xvi, 255 | antihypertensive drugs in pregnancy and hypertension in adult |
| age, maternal, risks relating to 84 | offspring 115 |
| preterm delivery 62 | anti-inflammatory cytokines 5 |
| ageing 268–98 | anti-inflammatory eicosanoids and pregnancy 67 |
| population 268–9 | antioncogenes (tumour suppressor genes) xxv, 243 |
| process 269–76 | antioridants xvi |
| nutritional modulation 273–6 | cancer and 257–9 |
| see also senescence | cardiovascular disease and 239 |
| | fertility and |
| albumin, serum, measurements 20 | females 41–3 |
| alcohol (ethanol) 210–12, 253 | males 41–5 |
| adolescence 210–12 | |
| binge drinking 73, 88, 210, 213 | pre-eclampsia and 78 |
| cancer and 253 | supplements see supplementation |
| elderly and osteoporosis and 283–4 | antiretroviral drugs and breastfeeding 142–3 |
| fertility and | Apolipoprotein E (ApoE) gene 13 |
| females 43–4 | apoptosis (programmed cell death) xvi, 270, 272 |
| males 47–8 | appetite |
| pregnancy and 73–4, 85–6 | adolescents 202 |
| miscarriage and 71 | melanocortin and 177 |
| allergy xvi | areola 123–4 |
| breastfeeding and 135 | ARF 273 |
| food see foods | arterial atheroma see atherosclerosis |

Nutrition, Health and Disease: A Lifespan Approach, Second Edition. Simon Langley-Evans. © 2015 John Wiley & Sons, Ltd. Published 2015 by John Wiley & Sons, Ltd. Companion website: www.wiley.com/go/langleyevans/lifespan





5/23/2015 3:03:59 AM



| artificial feeds see formula feeds | high see hypertension |
|---|---|
| ascorbic acid see vitamin C | reference levels (normal vs high) 241 |
| ATBC trial, Finnish 238, 246 | sodium and see sodium |
| atherosclerosis xvi, 230–2 | body composition |
| antioxidants and 239 | adolescent 190–1 |
| cerebrovascular disease and 232 | anthropometric measures 14, 15 |
| coronary disease and 232, 233 | infant/perschool children 151 |
| diabetes type 2 and 234 | multiple pregnancy and 86 |
| dietary fat and 235, 236 | pre-pregnancy 63 |
| homocysteine and 237, 238 | pre-eclampsia and 76 |
| obesity and 233, 234 | trauma impacting on 5 |
| Atkins 225 | see also fat |
| atopy and breastfeeding 135 | body height see height |
| attention deficit hyperactivity disorder xvii, 162 | body image (about shape/size/fatness), adolescent 203-4 |
| Australia | anorexia nervosa 207 |
| children | body mass index (BMI) xvii |
| food and drink advertising 165 | assessment 14 |
| food intake of school-age children 173–4 | cancer and 250 |
| dietary reference values 31, 32 | overweight/obesity and 223 |
| pregnancy and excess weigh management 74 | classification of 15, 223 |
| aversions and cravings in pregnancy 82–4 | school-age children 170-1, 178 |
| | pregnancy 63 |
| B cells xvii | sperm quantity and quality and 46–7 |
| ageing and 288 | body proportions, infant/perschool children 151 |
| babies see infants; newborns | body weight see weight |
| Baby Friendly Hospital Initiative (UNICEF) 140–1 | bone |
| bacterial overgrowth of small intestine, elderly 289 | growth in adolescence/puberty 193–6, 198–9 |
| balance (nutritional) 2–10 | mineralization and remodelling 281–2 |
| classical studies/measurements 8–10 | bone health (incl. mineral density) |
| BAPEN Nutrition Screening Survey 277 | adolescence and |
| Bardet–Biedl syndrome 175, 176 | female athletes and 206 |
| bariatric surgery xvii, 226, 227 | obesity and 196 |
| basal (resting) metabolic rate xvii, 5 | substance use and 213 |
| adolescent 198 | breastfeeding and 132 |
| adults 220 | definition of bone mineral density xvii |
| pregnancy 66 | elderly 276, 281–6 |
| Bax and apoptosis 272 | disorders 281–6 |
| Bcl2 and apoptosis 272 | minerals and 285–6 |
| behaviour land land land land land land land land | protein and 286 |
| breastfeeding and outcomes related to 133–4 | bottle-feeds see formula feeds |
| food-related, infant/perschool children 152 | bowel see intestine |
| health, adolescent 205 | boys |
| beverages (drinks) | genitalia see genitalia |
| advertising 165–7 | nutritional requirements |
| sugar-sweetened 175 bias 21–2 | adolescence 197 |
| | young children 152 |
| publication xxiv | puberty and sexual maturation 191, 192, 193 |
| binge drinking 73, 88, 210, 213 binge eating | brain growth and development breastfeeding and 133 |
| | infant/perschool children 151 |
| anorexia nervosa 206, 207 bulimia nervosa 204 | iron deficiency and 161 |
| bioimpedance xvii, 15 | BRCA1/BRCA2 gene 243 |
| biomarkers of nutritional status xvii, 18–20 | breakfast, children 165, 169–70 |
| biotin, sources/functions/deficiency symptoms 307 | missing 174, 202 |
| birth anthropometry, and disease in later life 100, 116 | breakfast cereals, fortification 169, 307 |
| birth weight | breast |
| fetal programming and 96, 97, 98, 100–1, 102, 103, 104, | anatomy 124–5 |
| 112, 116 | breastfeeding and the |
| multiple pregnancy 86 | breast cancer risk 131–2 |
| bladder cancer 244, 247 | problems arising in breast 139–40 |
| blood in pregnancy | cancer 243, 246, 247, 248, 249, 250, 251, 253 |
| composition 64 | breastfeeding and risk of 131–2 |
| volume expansion 63–4 | dietary factors reducing risk 255–6, 258, 259, 260 |
| blood pressure xvii, 240–2 | development 191 |
| | - |







| in pregnancy 127–8 | cardiovascular disease xvii, 230–42 |
|---|--|
| weaning and 128–9 | ApoE gene and 13 |
| breastfeeding 11, 123, 130–47 | definition and pathogenesis 230–2 |
| to 6 months 135-6 | maternal nutrition and risk to child in later life 100-1 |
| advantages 130-6 | obesity and 233-4 |
| alternatives to see formula feeds | childhood 176, 177, 178 |
| contraindications 141–2 | risk factors 233–42 |
| promoting 140–1 | Cardiovascular Health Study (US) 290 |
| reasons for not trying or giving up 138–40 | CARET trial 248, 258 |
| successful 139 | caries (dental) caries and fluoride, infant/perschool children 154, |
| ten steps 140 | 155 |
| trends 136–41 | |
| | carotenoids (incl. β-carotene) xvii, 307 cancer and 246, 248, 258 |
| weaning from see complementary feeding | |
| see also lactation | cardiovascular disease and 239 |
| British Association for Parenteral and Enteral Nutrition (BAPEN), | male fertility and 48–9 |
| Nutrition Screening Survey 277 | sources and functions 307 |
| bulimia nervosa (BN) xvii, 204, 208 | status determination 19 |
| drug abuse and 214 | symptoms of deficiency 307 |
| Bumps and Beyond intervention (UK) 74 | casein |
| | breast milk 127 |
| caffeine | cow's milk 144 |
| elderly 286 | casein-rich formula 145 |
| females | case-control studies xvii, 24, 25, 26 |
| fertility and 43–4 | cancer 247-8, 251, 255-6, 258, 259, 260 |
| pregnancy and 71–2, 75 | cause and effect 20–1 |
| calciferol see vitamin D | randomized controlled trials and evidence of 27 |
| calcium 306 | cells |
| adolescents 198–9 | cycle 270 |
| bone growth and 195-6 | p53 and 271–2 |
| blood pressure and 242 | telomeres and 272 |
| elderly, supplements 284–5 | fetal programming and number and type of 108–9 |
| infant/preschool children requirements 154 | programmed death xvi, 270, 272 |
| pregnancy 69 | senescence/ageing 270–3 |
| fetal programming and 104–5 | immune system 288 |
| pre-eclampsia and 77–8 | cereals (breakfast), fortification 169, 307 |
| prostate cancer and 257 | cerebrovascular disease (incl. stroke) xxiv, 232 |
| • | |
| sources/functions/deficiency symptoms 306 | cervical incompetence and preterm delivery 72 |
| caloric restriction and lifespan 273–5 | CHAOS study 239 |
| Canada, dietary reference values 31–2 | children 150–88 |
| cancer (malignant tumours) xvii, 242–61 | malnutrition see malnutrition |
| breastfeeding and 131–2 | nutrient intake see nutrient intake |
| diet as modifiable determinant of risk 244–5, 248 | nutrient requirements see nutrient requirements |
| dietary factors increasing risk 249–55 | obesity see obesity |
| dietary factors reducing risk 255–61 | preschool see infants and preschool children; newborns |
| nutritional epidemiology 245–9 | school-age 167–70 |
| see also specific sites | see also adolescence |
| cannabinoid receptors | China, GENSALT trial 242 |
| antagonist 226 | Chinese school-age children, food intake 173 |
| bone 213 | cholesterol 304 |
| cannabis 213, 214 | cardiovascular disease and 235 |
| capsaicin as carcinogen 255 | high-density lipoprotein see high-density lipoprotein-cholesterol |
| carbohydrates (sugars) 300–2 | low-density lipoprotein see low-density lipoprotein-cholesterol |
| adult intake 221 | uptake and transport 304 |
| breastmilk, synthesis 125 | chorionic gonadotrophin, human, and nausea and vomiting in |
| complex see complex carbohydrates | pregnancy 80 |
| digestion and absorption 301–2 | chronic disease |
| functions 300 | anaemia of 291 |
| preterm formulas 145–6 | elderly 279 |
| structure and classification 300–1 | ÷ · |
| | clay consumption in pregnancy 83, 84 |
| weight loss by restriction of 225–6 | clinical examination 20 |
| carbon dioxide transport and pregnancy 64 | cobalamin see vitamin B12 |
| carcinogenesis xvii, 243–4 | cocaine 214 |
| carcinogenic agents (carcinogens) xvii, 243–4, 245, 252, 253 | coffee, elderly 286 |
| specific ones in food 244, 253–5 | see also caffeine |







| cognitive development | cell, programmed xvi, 270, 272 |
|--|--|
| adolescent 196 | fetal, relating to diet 71 |
| breakfast and 169 | maternal teenage 209 |
| breastfeeding and 133 cognitive impairment, elderly 293–4 | deficiency, nutritional children |
| cohort studies xvii, 24, 25, 26 | infants/preschool 159–62 |
| cancer 248, 249, 251, 252, 253, 255, 256, 257, 259 | school-age 167 |
| maternal nutrition affecting offspring disease in later life 100–2 | defining 6 |
| Collaborative Health Behaviours in School-Aged Children | pica in pregnancy and 83 |
| Study 203 | see also specific nutrients |
| colorectal cancers (colon and rectal cancer; CRC) 247, 249, 251, | dehydration, elderly 276–7 |
| 252, 253, 259 | demand see supply and demand |
| complex carbohydrates and 256-7 | dementia xviii |
| milk/dairy produce and 257 | anaemia and 293–4 |
| see also large intestine disorders | dental caries and fluoride, infant/perschool |
| colostrum xvii, 125, 128 | children 154, 155, 155 |
| composition 124 | Department of Agriculture (US) and School Breakfast |
| commercially-available weaning foods 159 | Program 169–70 |
| Committee on Medical Aspects of Food Policy (COMA) 28, 29, | Department of Health's Public Health |
| 30, 66 | Responsibility Deal 182 |
| community, elderly nutritional support in the 280, 281 | developed (industrialized/Westernized) countries xviii |
| complementary feeding (weaning) 128–9, 156–9 | adolescent/teenager 189 |
| definition xvii, 137 | drug abuse 213–14 |
| early 158 | intakes 200 |
| late 158 | pregnancy 209, 210 |
| complex carbohydrates (fibre; non-starch polysaccharides) xvii, | vegetarianism 204 |
| xx, xxii | breastfeeding trends 138–9 |
| cancer and 255–6 | children |
| conception 10, 61 | complementary feeding 157 faddy eating 162 |
| see also preconceptual and periconceptual period conditionally-essential amino acids 305 | physical activity 172 |
| confounding factors xviii, 22 | weaning of infants 157 |
| constipation xviii | diabetes type 2 in ethnic minorities 228 |
| elderly 289–90 | male fertility problems 46 |
| pregnancy 84 | malnutrition 4, 7–8 |
| copper, sources/functions/deficiency symptoms 306 | migrant studies see migrant studies |
| coronary (ischaemic) heart disease (CHD) xviii, 232 | pregnancy |
| fetal programming and risk in later life of 99, 100-1 | adolescent 209, 210 |
| risk factors 233, 235, 236, 238, 239 | nutrient requirements 66, 69, 70 |
| obesity/overweight 179, 233-4 | developing countries xviii |
| corpus luteum xviii, 36, 37, 41, 62 | breastfeeding 132, 135, 136 |
| cosmetics, endocrine-disrupting chemicals 49 | alternatives 143 |
| cow's milk | children |
| cancer and 257 | complementary feeding 156–7, 158 |
| young children 155 | nutrient requirements 153, 154 |
| formula feeds 143–6 | poverty 163–4 |
| cravings and aversions in pregnancy 82–4 | zinc deficiency 160 |
| cross-sectional study xviii, 24, 25–6 | iron status 292 |
| cruciferous vegetables xviii | pregnancy 68, 69 |
| cancer and 261 | iron supplementation 69 |
| cryptorchidism xviii, 39, 50 cultural factors see social | malnutrition 4, 7, 8 pregnancy |
| (and socioeconomic/sociocultural) factors | adolescent 209, 210 |
| cytochrome P450 system xviii | iron status 68, 69 |
| CYP2E1 and nitrosamines 244 | protein requirements 67 |
| cytokines xviii, 5 | tuberculosis 143 |
| anti-inflammatory 5 | undernutrition (adults) 220 |
| elderly 287 | development |
| pro-inflammatory see pro-inflammatory cytokines | breastfeeding and 133–4 |
| i e e e e e e e e e e e e e e e e e e e | fetal programming see programming |
| daidzein 50, 51, 286 | infant/perschool children |
| dairy products and cancer 257 | iron deficiency and 161 |
| dark adaptation test 19 | milestones 150–1 |
| db/db mouse 40, 195 | psychosocial 152 |
| death (mortality) | physical see physiological (incl. physical) changes |







| Ji-h 227 0 | 1 1 212.14 |
|--|--|
| diabetes 226–9 | drug abuse 213–14 |
| childhood obesity and 176 | dyslipidaemia (hyperlipidaemia) xvii, 235, 236 |
| gestational 84–6 | diabetes and 234 |
| maturity onset, of the young xxi, 227–8 | dyspepsia in pregnancy 84 |
| type 1 107, 176, 226 | |
| type 2 176, 226–9 | Eat Well and Keep Moving 182 |
| and fetal programming 112, 116 | eating behaviour and adolescent food choice 200, 201 |
| diaphysis and adolescent bone growth 193 | eating disorders, adolescent 204, 206–8 |
| diet | drug abuse and 214 |
| adolescent bone growth and 195 | see also feeding; food; meals |
| disease and see disease | Eatwell plate 221, 222 |
| intake see nutrient intake | eclampsia xviiI, 76 |
| pregnancy outcome relating to 71–9 | ecological studies xvii, 24, 25 |
| recall methods 16–17 | cancer 245–6, 251 |
| reference values (DRVs) 28–31 | education, elderly 279 |
| UK 28-31 | egg (ovum) 36, 37 |
| other countries 31–2 | eicosanoids and pregnancy 67 |
| requirements see nutrient requirements | elderly 268–98 |
| source of minerals 306 | barriers to healthy nutrition 277–81 |
| source of vitamins 307 | health problems/disease related to nutrition 279, 281–94 |
| Dietary Approaches to Stop Hypertension (DASH) study 242 | nutrient requirements 276–7 |
| dietary interventions (incl. dietary modification and diversification) | vitamin B12 status and folate fortification in 56 |
| cancer risk and 244–5, 248 | embryo (and embryogenesis) xviii, 10 |
| obesity 225–6 | programming 94–5 |
| childhood 179–80 | emotional bond in breastfeeding 131 |
| in osteoporosis 284–6 | emotional challenges in adolescence 207 |
| vitamin A deficiency prevention 8 | endocrine system (and its regulation incl. hormones) |
| see also dieting; restriction | adolescence and puberty 192, 193, 195 |
| dieting xvii | bone mineralization and 283 |
| adolescent 202, 203–4 | chemicals interfering with (endocrine disruptors) xviii, 49–51 |
| digestion and absorption | fertility/reproductive function and |
| carbohydrates 301–2 | females 36–7, 44–5 |
| lipids 303–4 | males, obesity and 47 |
| proteins 306 | lactation and 127–8 |
| see also malabsorption | pregnancy and 62 |
| digestive tract see gastrointestinal tract | fetal programming and 109 |
| disaccharides 300, 301 | nausea and vomiting in 80 |
| discordance hypothesis of obesity 171–2 | endothelial dysfunction xix |
| disease 223–61 | atherosclerosis and 231 |
| diet/nutrition and, relationship between 20–8, 28–9, 223–61 | energy demands/requirements 276 |
| elderly 279, 281–94 | elderly 276 |
| quantifying 22–4 | infant/preschool children 152–4 |
| epidemiology see epidemiology | injury and infection impacting 5 |
| obesity and see obesity | lactation and 130 |
| diverticula/diverticulitis/diverticular disease, elderly xvii, 290 | pregnancy 65–6, 66–8 |
| DNA | multiple 86–7 |
| cancer and | school-age children 167 |
| Ames test for mutagenesis xvi, 254–5 | adolescents 197–8 |
| folate and 259 | see also protein–energy malnutrition |
| methylation see methylation | energy intake/consumption |
| oxidative damage 260, 271 | doubly-labelled water method of estimation 18, 19 |
| telomere see telomere | maternal, fetal programming and 105–6 |
| uracil misincorporation xxv, 259 | school-age children 174–5 |
| docosahexaenoic acid (DHA) | adolescents 200, 202 |
| breastfeeding and 134 | in treatment of obesity 179–80 |
| cardiovascular disease and 236 | energy source |
| pregnancy and 67 | carbohydrate as 300 |
| doubly-labelled water method of energy intake estimation 18, 19 | lipid as 302 |
| drinks see alcohol; beverages | environmental factors |
| drug(s), (pharmacological/prescription/therapeutic) | adolescent food choice 200 |
| breastfeeding and 141 | fetal programming through exposure |
| in diabetes type 2 229 | to see programming |
| in hyperlipidaemia 236–7 | obesity, combined with genetic factors 171–2 |
| in obesity treatment 226 | periconceptual period and 52 |
| in pregnancy regarding fetal programming 115 | environmental oestrogens 49–51 |







| epidemiology 20–8 | breastfeeding and its advantages for 130-2 |
|--|--|
| cancer 245–9 | coronary heart disease deaths 232 |
| malnutrition 7–8 | fertility and reproductive function 36–44 |
| maternal nutrition affecting offspring disease in later life | menarche see menarche |
| 99–103 | nutritional requirements of adults |
| study designs 24–8 | compared with adolescent children 197 |
| see also prevalence | compared with infant/preschool children 152 |
| epigenetics xix | obesity see obesity |
| fetal programming and 113–15 | young see girls |
| and transgenerational transmission of disease risk 117–18 see also methylation | see also pregnancy female athlete triad 206 |
| epiphyseal plate and adolescent bone growth 193 | fertility (and reproductive function) |
| erythrocytes (red blood cells), examination in anaemia 290, 291 | female 36–44 |
| erythropoiesis, disordered 291 | male 44–51 |
| essential amino acids xix, 305 | vegetarian adolescents 205 |
| essential fatty acids xix | see also infertility and subfertility |
| dietary reference values 30 | fetus (foetus) 10–11 |
| infant/preschool children requirements 154 | alcohol and (fetal alcohol syndrome and fetal alcohol spectrum |
| ethanol see alcohol | disorders) 73, 87–8 |
| ethnic groups | death relating to diet 71 |
| cardiovascular disease risk related to 233 | developmental programming (and disease in later life) see |
| minority, in Westernized countries, type 2 diabetes 228 | programming |
| EURAMIC study 19, 239 | folate demand 54 |
| Europe and European Union (EU) | gestational diabetes and 85 |
| breastfeeding trends 137 | growth see growth |
| dietary reference values 31 | placenta and 62 |
| European Prospective Investigation into Cancer (EPIC) 241, 248, | see also teratogens |
| 249, 251, 256 | fibre (dietary) see complex carbohydrates |
| evolution and obesity 171–2 | Finnish ATBC trial 238, 246 |
| examination, clinical 20 | first-stage milk 144 |
| exercise see physical activity experimental studies see animals | flavonoids and cancer 260–1 see also isoflavones |
| experimental studies see animals | fluoride (fluorine) 306 |
| faddy eating xix, 162–3 | infant/preschool children 154, 155 |
| faecal bulking by fibre and colorectal cancer 257 | sources/functions/deficiency symptoms 306 |
| falls and vitamin D, elderly 285 | fluorosis 155 |
| family in childhood obesity prevention 181 | foam cells xix |
| fast food 165, 166, 167, 174–5, 201–2 | and atherosclerosis 231–2 |
| fat, body 223 | folic acid/folate (vitamin B9) 259, 307 |
| female | cancer and 246, 257, 259 |
| distribution 223 | cardiovascular disease and 237–9 |
| fertility and 39 | elderly, deficiency 293 |
| pregnancy and 66 | pregnancy and 54–6 |
| male, distribution 223 | sources/functions/deficiency symptoms 307 |
| fat, dietary see fatty acids; lipid | follicle-stimulating hormone (FSH) xix |
| fat-soluble vitamins 307 | females 36, 37 |
| fatty acids xix, 235–6, 302, 303 | males 44, 45 |
| breast milk 126 | follicular phase xix, 36, 37 |
| essential see essential fatty acids | follow-on milk 144–5 |
| infant/preschool children requirements 154 polyunsaturated <i>see</i> polyunsaturated fatty acids | food(s) |
| pregnancy and supply of 67–8 | accounts 16 allergy xix |
| saturated 235, 302, 303 | children 167 |
| trans- xxv, 235–6, 302 | balance sheets 14–16 |
| unsaturated <i>see</i> monounsaturated fatty acids; polyunsaturated | carcinogens in 244, 253–5 |
| fatty acids; unsaturated fatty acids | choice in adolescence, factors influencing 200–1 |
| uptake and transport 304 | consumption out of home by adolescent 201–2 |
| fatty streak xix, 231, 234 | cravings and aversions in pregnancy 82–4 |
| feeding | fast 165, 166, 167, 174–5, 201–2 |
| elderly 281 | fortification see fortification |
| infant see infants | frequency questionnaire 16, 17–18 |
| see also eating; food; meals | functional see functional foods |
| female(s) (women) 11 | groups xix, 221 |
| basal metabolic rate 220 | infant/preschool children and |
| body fat see fat | advertising 165–7 |







| food additives and hyperactivity 161–2 | bone growth 194–5 |
|--|--|
| food neophobia xix, 162–3 | fetal programming 96, 106–7, 111–13, 116 |
| food-related behaviour 152 | transgenerational transmission of disease risk |
| restriction 167 | 117–18 |
| intake see nutrient intake | polymorphisms see polymorphisms |
| intolerance xix | see also epigenetics; inborn errors of metabolism |
| lactose xxi, 146, 279 | genistein 50 |
| labelling 28, 223 | genitalia, boys |
| adolescents and 201 | defects xviii, 39, 50 |
| symbols 222 | development in adolescence 191 |
| records/diaries 16, 17 | genomic imprinting xix, 117, 118 |
| snack see snacking | genotype, thrifty 106, 107, 228 |
| see also eating; feeding; meals | GENSALT trial (China) 242 |
| food chain, endocrine-disrupting chemicals 49 | germ cell (male) 44 |
| Food Circle, Swedish 221 | gestational diabetes 84–6 |
| food industry and obesity 182 | girls |
| Food Pyramid 221 | adolescent |
| foodstuffs | body fat 39 |
| non-nutrient components, cancer and 260-1 | nutritional requirements 197 |
| traffic light classification 179, 223 | puberty and sexual maturation 191, 192, 193 |
| foremilk 125 | sport and the female athlete triad 206 |
| formula/artificial/bottle feeds 143–6 | young, nutritional requirements 152 |
| trends/preferences 136, 137, 138–40 | global/worldwide dimensions |
| fortification xix | obesity 223–4 |
| breakfast cereals 169, 307 | undernutrition 104, 106 |
| folic acid 55–6 | glucocorticoids and fetal programming 109–11 |
| iodine 69, 70 | gluconeogenesis xx, 5 |
| vitamin A 8 | glucose |
| frail elderly, nutritional support in the community 280, | blood |
| 281 | control (by diabetic patient) xx, 228–9 |
| free radicals xix | transport in pregnancy 65 |
| elderly 271 | homeostasis xx |
| fertility and | glucose tolerance |
| females 41 males 48 | impaired 101, 106, 230 |
| see also reactive oxygen species | oral glucose tolerance test xxii, 227 glucosinolates xx, 261 |
| fruit and vegetable consumption | GLUT4 and pregnancy 65 |
| cancer and 258, 260–1 | glutathione xx |
| children 168–9 | glycaemic control (by diabetic patient) xx, 228–9 |
| FTO gene 175, 176 | glycaemic index xx |
| functional foods xix | and dietary interventions 225–6 |
| cancer and 257 | glycosylated haemoglobin xx, 228 |
| functional measures of biomarkers 19 | goat's milk (formula feed) 148 |
| fungal toxins, carcinogenic 255 | gonadotrophin, human chorionic, and nausea and vomiting |
| | in pregnancy 80 |
| galactosaemia xix | gonadotrophin-releasing hormone (GnRH) xx |
| breastfeeding 141 | females 36, 37 |
| Gambia, maternal nutrition and fetal programming 96–7 | males 44, 45 |
| gastric see stomach | government and obesity prevention 182 |
| gastrointestinal (digestive) tract | growth |
| ageing 269 | adolescent/puberty (incl. growth spurt) 11, 39, 189–90, 191, |
| disorders 289–90 | 192, 197, 198 |
| breastfeeding and protection from 133 | bone 193-6, 198-9 |
| in pregnancy 65 | fetal/neonatal 11, 95–9 |
| disturbances 84 | constraints 96–7 |
| generations, transmission of disease across 117-18 | health and disease and 97–9 |
| genetic factors | maternal growth and, competition between 209-10 |
| diabetes type 2 230 | infant/preschool children 151 |
| eating disorders | gustation (taste), elderly 279 |
| anorexia nervosa 207 | gynoid pattern of body fat distribution 223 |
| bulimia nervosa 208 | |
| obesity 175–6, 227–8 | haemoglobin |
| environmental combined with 171–2 | glycosylated xx, 228 |
| genetic syndromes, premature ageing 271 | in pregnancy 64 |
| genetics 12–14 | elevated 68 |







| hair | incidence rates 22, 23 |
|---|---|
| micronutrient determination 19 | individual (person) |
| pubic, development 191, 193 | adolescent food choice influenced by personal factors |
| see also hirsutism | 200-1 |
| Hayflick limit xx, 242, 270, 272 | advice on nutrition tailored to xxiii |
| hazard ratio 22–4 | in pregnancy regarding fetal programming 115 |
| Healthy Lifestyle in Europe by Nutrition in Adolescence | overall nutritional status 9–10 |
| (HELENA) Study 200, 201 | response to nutrition 10–13 |
| Healthy Start Service and vouchers 156, 165 heart disease, ischaemic <i>see</i> coronary heart disease | indoles and cancer 261 industrialized countries see developed countries |
| heartburn in pregnancy 84 | infants (babies) and preschool children |
| height/stature of children | advertising susceptibility 165, 166 |
| gain | barriers to healthy nutrition 162–7 |
| adolescent 190 | development see development |
| infant/preschool children 151 | feeding 123–47 |
| stunting 153 | advantages of breastfeeding for infant 130, 132–5 |
| HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) | newborn see newborns |
| Study 200, 201 | nutrient intake 154–6 |
| Helicobacter pylori xx, 245 | transition to adult pattern 156-67 |
| elderly 289 | nutrient requirements 151–4 |
| herbs, carcinogens 255 | nutritional problems 159–62 |
| high-density lipoprotein-cholesterol (HDL-cholesterol) xx | obesity 170–1 |
| childhood obesity and 179 | premature 72–6 |
| hindmilk 125 | sugar-sweetened beverages 175 |
| hirsutism xx, 38 | infection |
| history-taking 20 | breastfeeding and |
| HIV transmission risk in breastfeeding 142–3 | protective role 125, 132–3 |
| homeostasis model assessment – insulin resistance (HOMA-IR) | risk of breast infection 139–40 |
| scale 229 homocysteine and cardiovascular disease 237–9 | risk of transmission of infection 142–3 |
| HOPE 2 trial 238 | elderly 286–7 inflammatory response see inflammatory response |
| hormones see endocrine system | infertility and subfertility xxi |
| human chorionic gonadotrophin and nausea and vomiting in | females 36–9, 43 |
| pregnancy 80 | males 44–6 |
| human immunodeficiency virus transmission risk in | inflammatory response (to trauma or infection) 5 |
| breastfeeding 142–3 | acute phase 5 |
| hydrolysed protein in formulas feeds 148 | in pregnancy 74 |
| hydroxyproline, adolescents 199 | pre-eclampsia and 77, 78 |
| 11-hydroxysteroid dehydrogenase-2 109–10, 111 | see also entries under anti-inflammatory; pro-inflammatory |
| 5-hydroxytryptamine (serotonin) and anorexia nervosa 207 | information bias 21–2 |
| hyperactivity and food additives 161–2 | inhibin 37 |
| hyperemesis gravidarum xx, 80, 81–2 | initiation phase of carcinogenesis 243 |
| hyperhomocysteinaemia and cardiovascular disease xx, | INK4a 273 |
| 237–8 | Innocenti Declaration 135 |
| hyperlipidaemia see dyslipidaemia | institutionalized (incl. nursing home) elderly persons 276–80, 28 |
| hypermetabolic response to trauma 4–6 | 285, 287, 292 |
| hypertension (high BP) xx, 232–3 | insulin 229–30 |
| in adult offspring (and fetal programming) interventions 115 | pregnancy 65 |
| maternal sodium intake and 105 | insulin resistance xxi, 229, 230 |
| pregnancy 76–8 | atherosclerosis and 234 fetal nutrition and, in later life 100, 101, 105 |
| hypospadias xx, 39, 50 | obesity and 38, 47, 251 |
| hypothalamus xx | pregnancy 65, 84–5, 87 |
| leptin and 40 | insulin resistance syndrome (syndrome X; metabolic |
| menstrual cycle and 36 | syndrome) xxi, 100, 176, 229–30 |
| spermatogenesis and 44 | intelligence and breastfeeding 133 |
| | intergenerational transmission of disease risk 117–18 |
| immunity | internet, food and drink advertising aimed |
| ageing and 269–70, 286–9 | at children 165, 166 |
| breastfeeding and 125, 132–3 | INTERSALT 241, 242 |
| zinc and 117, 160 | intestine (bowel) |
| immunoglobulins (antibodies) xxi | elderly, disorders 289–90 |
| IgA, breast milk 126 | in pregnancy 65 |
| imprinting, genomic xix, 117, 118 | invasion phase of carcinogenesis 243 |
| inborn errors of metabolism and breastfeeding 141–2 | involution of breast 129 |







| iodine 306 | congenital deficiency 175 |
|---|---|
| deficiency (and its prevention) 8, 306 | fetal programming and 112 |
| pregnancy 69–70 | miscarriage and 71 |
| sources/functions 306 | receptor defects 175, 176 |
| IQ and breastfeeding 133 | letdown reflex xxi, 128, 131, 140 |
| iron 306 | leukocyte telomere length 272–3 |
| excess (consumption or overload) 3 | Leydig cells xxi, 44 |
| maternal requirements 68 | life expectancy xxi, 268 |
| sources/functions 306 | lifespan |
| iron deficiency 104 | caloric restriction and 273–5 |
| adolescents 199 | fetal programming 275 |
| in pregnancy 210 | introduction to lifespan nutrition 1-34 |
| anaemia see anaemia | lifestyle factors (and their modification) 223-61 |
| children 161 | adolescent bone growth 195 |
| newborn infants 136 | adulthood disease states 223–61 |
| elderly 290, 291, 292 | bone growth 195 |
| pregnancy 7, 64, 68, 69, 83, 84, 104, 210 | obesity 174, 179, 180, 181 |
| stages 7 | periconceptual period and 52 |
| symptoms 306 | LIMIT trial (Australia) 74 |
| iron deficiency anaemia 6, 7, 104 | linoleic acid 67, 236, 303 |
| adolescence 199, 210 | deficiency in children 30 |
| children 161 | linolenic acid 67, 302 |
| elderly 290, 292 | lipid, dietary (fats) 235–9, 302–4 |
| newborn infants 136 | adult intake 221 |
| pregnancy 7, 63, 69, 83, 84, 104, 210 | breast milk 126 |
| ischaemic heart disease see coronary heart disease | cardiovascular disease and 235–9 |
| islet cells and fetal programming 108–9 | digestion and absorption 303-4 |
| isoflavones 50, 51, 260, 286 | functions 302 |
| isothiocyanates and cancer 261 | infant/preschool children requirements 153-4 |
| | liver, accumulation in alcohol abusers 211 |
| JAK-STAT3 40–1 | pregnancy, demands/requirements 67-8 |
| Jamaican boys, maternal nutrition and disease in later life 101 | structure and classification 302–3 |
| Japan | see also dyslipidaemia |
| migrants in US from, cancer 246–7 | lipoproteins xxi |
| Okinawan diet 274–5, 275 | see also Apolipoprotein E; high-density lipoprotein-cholesterol; |
| | low-density lipoprotein-cholesterol |
| kidney (renal) | liquorice 111 |
| pregnancy-related changes 64 | liver |
| remodelling (in fetal programming) 108 | cancer 250, 255, 258 |
| Korean migrants in US, cancer 247 | lipid accumulation in alcohol abusers 211 |
| kwashiorkor xxi, 7 | pregnancy and consumption of 54 |
| | Livsmedelsverket Keyhole 223 |
| labelling, food see food | long chain polyunsaturated fatty acids (LCPUFA) in pregnancy 67, 68 |
| labour | losartan in pregnancy and hypertension in adult offspring 115 |
| abnormal 79 | low-density lipoprotein-cholesterol (LDL-cholesterol) xxi, 231, |
| premature, and delivery 72–6 | 235, 236 |
| α-lactalbumin in whey-rich formula milks 144–5 | childhood obesity and 178, 179 |
| lactation 123–30 | hyperhomocysteinaemia and 238 |
| maintenance 129–30 | lower reference nutrient intake (LRNI) xxi |
| nutritional demands 129–30 | luteal phase xxi, 36, 37 |
| lactiferous ducts and sinuses 124 | luteinizing hormone (LH) xxi |
| lactoferrin | females 37 |
| breast milk 126–7, 145 | males 44, 45 |
| whey-rich formula milks 145 | lymphocytes xxi |
| lactoglobulin in whey-rich formula milks 144 | and ageing 287 |
| lactose | |
| breast milk 125 | macronutrients 300 |
| intolerance xxi, 146, 279 | breast milk vs. other feeds 144 |
| `large baby syndrome' (macrosomia) xxi, 85 | child requirements and intake |
| large intestine disorders, elderly 289–90 | adolescence 197–8 |
| see also colorectal cancers | infants/preschool 152–4 |
| laryngeal cancer 250, 258 | school-age 173, 174 |
| latching on 124, 139 | definition 300 |
| leptin 39–41 | elderly requirements 276 |
| adolescent bone growth and 195 | maternal intake and fetal programming 105–6 |
| = | |







| macrophages xxi | trauma effects 4–6 |
|---|---|
| ageing and 287 | xenobiotic xxiii, 214, 244, 261 |
| atherosclerosis and 231–2 | see also basal metabolic rate |
| see also foam cells | methionine cycle 237 |
| macrosomia xxi, 85 | methylation of DNA xvii |
| magnesium 306 | cancer and 259 |
| blood pressure and 242 | fetal programming and 114 |
| pregnancy and 69 | see also epigenetics |
| sources/functions/deficiency symptoms 306 | methylenetetrahydrofolate reductase (MTHFR), C677T single |
| malabsorption, elderly 289 | nucleotide polymorphism 12, 13, 237–8, 259 |
| males (men) | microcytic anaemia 291 |
| basal metabolic rate 220 | micronutrients (trace nutrients) 300, 306–8 |
| body fat distribution 223 | adolescence 198–9, 199 |
| coronary heart disease deaths 232 | sport and physical activity and 205 |
| fertility 44–51 | breastfeeding and 129–30 |
| obesity 46–7, 224 | cancer and 246 |
| young see boys | deficiencies xxii |
| malignant tumours see cancer | infant/preschool children 159–61 |
| malnutrition xxi, 6–8, 220 | school-age children 167 |
| causes 7 | definition 300 |
| children 7, 153 | elderly requirements 276 |
| poverty and 163 | hair, determination 19 |
| elderly 277–81 | infant/preschool children |
| combating 279–81 | deficiencies 159–61 |
| vicious cycle of disease and 286–7 | requirements 153, 154 |
| epidemiology 7–8 | supplements 155–6 |
| prevention see prevention | pregnancy 68–9 |
| protein–energy see protein–energy malnutrition | fetal programming and 104–5 |
| marasmus xxi, 7 | pre-eclampsia and 77 |
| marginal nutrition 2 | see also antioxidants; minerals; vitamins |
| mastitis and breastfeeding 139–40 | migrant studies (in Westernized countries) |
| maturity onset diabetes of the young xxi, 227–8 | cancer 246–7 |
| meals | diabetes type 2 228 |
| elderly 280–1 | milk |
| missing/skipped 174, 202 | breast |
| school 168–9, 181–2 | expression, and breast pumps xix, 129 |
| see also feeding; food; meals | synthesis/composition 124–7 |
| measurement bias 21–2 | non-human |
| meat consumption and cancer 247, 252–3 | cancer and consumption of 257 |
| media | for formulae feeds see formula feeds |
| body image and 203–4 | see also cow's milk |
| healthy nutrition and 221 | Millennium cohort 103, 158 |
| see also internet | Millennium Development Goal (UN) 164 |
| Medical Research Council (MRC) Vitamin Study Group 11, | Millennium Project (UN) 164 |
| 54, 55 | minerals (incl. trace elements) 306–8 |
| megaloblastic anaemia 291, 292, 293 | adolescence 198–9 |
| melanocortin 177 | port and physical activity and 205 |
| receptors 177 | blood pressure and 240–2 |
| mutations/defects 175, 176, 177 | infant/preschool children 154 |
| men see boys; males | supplements 155–6 |
| menarche xxi, 36, 39, 193 | osteoporosis 284–6 |
| earlier 39, 208, 251 | pregnancy 68–70 |
| menstrual cycle 36–7 | pre-eclampsia and 77–8 |
| body fat and 39 | sources/functions/deficiency symptoms 306–8 |
| disorders 37–8, 39 | miscarriage (spontaneous abortion) 71–2 |
| follicular phase xix, 36, 37 | teenagers 209 |
| luteal phase xxi, 36, 37 | mismatched environments 108, 117 |
| meta-analysis xxi, 27–8 | monkey studies of caloric restriction and lifespan 273 |
| metabolic syndrome (insulin resistance syndrome; syndrome X) xxi, | monosaccharides 300, 300–1 |
| 100, 176, 229–30 | monounsaturated fatty acids (MUFA) 235, 251 |
| metabolism | structure and classification 302, 303 |
| inborn errors of, and breastfeeding 141–2 | mortality see death |
| pregnancy-related changes 65–6 | mouth |
| thrifty see thrift | cancer 250 |







| antioxidants and 258 | nutrient requirements |
|--|---|
| elderly, disorders 289 | adults see adulthood |
| MRC Vitamin Study Group 11, 54, 55 | children |
| Mullerian ducts xxii, 45 | adolescents 197–9 |
| multiple pregnancy/birth 86–8 | infants/preschool 151-4 |
| preterm delivery 72, 86 | school-age 167–8 |
| multivitamin supplements | elderly 276–7 |
| elderly 287 | pregnancy 66–71 |
| male infertility 49 | nutrient types 300–8 |
| pregnancy preparation 53, 54 | nutritional balance/deficiency/status etc. see balance; deficiency; |
| muscle (skeletal) mass and ageing 269 | status etc. |
| MUST (malnutrition universal screening tool) 280 | |
| mutagens, Ames test xvi, 254–5 | ob/ob mouse 40, 195 |
| My Pyramid 221–2 | obesity and overweight xxii, 170-82, 223-6, 249-51 |
| | adulthood 223–6 |
| n-3 (omega-3) fatty acids xxii, 67, 68, 302 | causes 224 |
| cancer risk and 251 | prevalence 223–4 |
| cardiovascular disease risk and 236 | treatment/interventions 224–6, 234 |
| n-6 (omega-6) fatty acids xxii, 67, 68, 302, 303 | atherosclerosis and 233, 234 |
| National Diet and Nutrition Surveys (NDNS in UK) 199, 200 | breastfeeding and 134, 135 |
| adolescence 199 | children 170–82 |
| preschool children 154, 160–1 | adolescent bone growth and health and 195, 196 |
| National Health and Nutrition Examination Study (NHANES) 28, | causes 171–6 |
| 70, 171, 174, 203, 224, 293 | consequences 176–8 |
| National School Fruit and Vegetable Scheme (UK) 168 | prevalence 170–2 |
| natural killer cells xxii | prevention 180–2 |
| and ageing 287 | tracking 176–8 |
| nausea and vomiting in pregnancy (NVP) 79–82 | treatment 177, 178–80 |
| food cravings and aversions and 82–3 | weaning and risk of 158 |
| negative stereotypes in breastfeeding 139 | classification 15, 223 |
| neonates see newborns | disease and 220 |
| neophobia, food xix, 162–3 | cancer 249–51 |
| Netherlands Cohort Study 256 | cardiovascular see cardiovascular disease |
| neural tube defects xxii | childhood obesity 176, 178 |
| and folic acid 54–6 | female 224 |
| neurodevelopmental disorders and fetal alcohol exposure 87 | polycystic ovary syndrome and 38, 71 |
| neutrophils and ageing 287 | pregnancy and see subheading below |
| New Zealand, dietary reference values 31, 32 | insulin resistance and 38, 47, 251 |
| newborns (neonates) 11 | male 46–7, 224 |
| breastfeeding 140 | pregnancy and 73, 74, 102 |
| growth see growth | hypertension and 78–9 |
| Newcastle Thousand Families Study 177–8, 178 | management 74 |
| NHANES (National Health and Nutrition Examination Study) 28, | miscarriage and 71 |
| 70, 171, 174, 203, 224, 293 | neural tube defects and 55 |
| niacin, sources/functions/deficiency symptoms 307 | offspring risks 102 |
| nicotinic acid (niacin), sources/functions/deficiency symptoms 307 | ochratoxins 255 |
| nipple 123–4 | odds ratio xxii, 22–4 |
| nitrosamines and nitroso compounds as carcinogens 244, 245, | oesophageal disorders |
| 247, 252, 253 | cancer 20, 21, 22, 244, 249, 250, 251, 253, 254, 255, 257–8 |
| normocytic anaemia 291 | elderly 289 |
| Northampton, breastfeeding trends 138–9 | oestrogen(s) |
| nuns (English), cancer 247 | anorexia nervosa and 207 |
| Nurses Health Study (US) 72, 100, 131, 248, 251, 255, 256 | breast cancer and 251 |
| nursing homes and other institutional case system for elderly | environmental 49–51 |
| persons 276–80, 281, 285, 287, 292 | menstrual cycle and 36, 37 |
| nutrient intake (dietary/food intake) | plant-derived (phytoestrogens) xxiii, 50-1, 260-1, 286 |
| ageing process and 273–6 | in pregnancy |
| children | breast and 127 |
| adolescent 199–202 | nausea and vomiting and 80 |
| infants/preschool see infants | Okinawan diet 274–5, 275 |
| school-age 173–5 | olfaction (smell), elderly 279 |
| estimation 14–18 | oligosaccharides 300, 301 |
| reference see reference nutrient | omega-3 and -6 fatty acids see n-3 fatty acids; n-6 fatty acids |
| see also diet | oncogenes xxii, 243 |







| oocytes 36, 43, 48 | pesticides 51 |
|---|---|
| | • |
| oral glucose tolerance test xxii, 227 | pharmacotherapy see drugs |
| oral health in pregnancy 74–6 | pharyngeal cancer 250, 258 |
| organ(s) | phencyclidine 214 |
| aging 269 | phenolic compounds (incl. polyphenolics) xxiii |
| formation (organogenesis) 61, 62 | cancer and 260, 261 |
| remodelling (in fetal programming) 108–9 | cardiovascular disease and 240 |
| organic farming xxii | phenotype, thrifty 106–7, 108, 116 |
| semen quality and 51 | phenylketonuria xxiii |
| orlistat 226 | and breastfeeding 142 |
| osteoblasts xxii, 281, 282 | phospholipids xxiii, 303 |
| osteoclasts xxii, 281 | phosphorus, sources/functions/deficiency symptoms 306 |
| osteopenia xxii, 206 | phthalates 50 |
| of prematurity 145 | physical activity, exercise (and its lack/sedentary activity), children |
| osteoporosis xxiii, 112, 196, 282–6 | (5-13) 172–3 |
| childbearing/breastfeeding and 132 | adolescents and 204–6 |
| dietary interventions 284–6 | bone growth and 195 |
| fetal programming and 112 | energy requirements and 198 |
| pathology and prevalence 282–3 | obesity and |
| risk factors 283–4 | risk of 172–3 |
| ovaries | treatment of 179 |
| cancer | physical activity/exercise (and its lack/sedentary activity), adults |
| breastfeeding and 131 | elderly, and bone mineralization 283 |
| complex carbohydrates and 255 | obesity and 224 |
| polycystic ovary syndrome (PCOS) xxiii, 38-9, 71 | treatment of 226 |
| overnutrition xxiii, 2–4, 220 | physical changes see body composition; physiological changes |
| fetal programming and 94, 103, 112, 115, 117, 118 | physical examination 20 |
| see also obesity | physiological (incl. physical) changes |
| overweight see obesity and overweight | adolescents 189–96 |
| ovulation xxiii, 36, 37, 41 | ageing/the elderly 269-70, 279 |
| absence (anovulation) xvi, 37, 39 | infant/preschool children 151 |
| ovum (egg) 36, 37 | pregnancy 62–6 |
| oxidative stress and damage xxiii | phytic acid 160 |
| ageing and 271, 272, 273 | phytoestrogens xxiii, 50–1, 260–1, 286 |
| DNA 260, 271 | pica xxiii |
| female fertility and 41, 43 | in pregnancy 84 |
| oxygen, reactive species <i>see</i> reactive oxygen species | pituitary gland |
| oxytocin 128, 131 | anterior xvi |
| 0.1,10cm 120,131 | leptin and 40 |
| p16INK4a 273 | posterior xxiii |
| p53 and ageing 271–2 | placenta 61–2 |
| Paget's disease of bone xxiii, 286 | fetal programming and 109–10, 111 |
| pancreas | hypertensive disease and formation of 77 |
| cancer 250, 257, 259 | plants |
| islets and fetal programming 108–9 | oestrogens derived from (phytoestrogens) xxiii, 50–1, |
| pantothenic acid, sources/functions/deficiency symptoms 307 | 260–1, 286 |
| parents | |
| • | protein |
| faddy eating/food neophobia and 163 | availability by world region 15, 16 |
| food/food group restriction by 167 | quality 304 |
| in obesity | stanols and stanol esters 236 |
| in prevention 181 | see also fruit and vegetable consumption; herbs |
| in treatment 180 | plaques, atherosclerotic 231 |
| peer support in breastfeeding 141 | polycystic ovary syndrome (PCOS) xxiii, 38–9, 71 |
| peptic ulcers 289 | polymorphisms xxiii |
| periodontal disease xxiii | fetal programming and 111–12 |
| pregnancy 74–6 | type 2 diabetes and 112, 116 |
| peripheral artery disease 232 | single nucleotide see single nucleotide polymorphisms |
| peristalsis (age-related decline) | polyphenolics see phenolic compounds |
| large intestine 290 | polysaccharides 300, 301 |
| oesophagus 289 | non-starch polysaccharides see complex carbohydrates |
| pernicious anaemia xxiii, 56, 292 | polyunsaturated fatty acids (PUFA) 235, 236 |
| peroxisome proliferator-activated receptor γ2 (PPAR-γ2) | breastfeeding and 134 |
| 111–12, 116 | long chain (LCPUFA), in pregnancy 67, 68 |
| person see individual | structure and classification 302, 303 |







| ponderal index xxiii, 100, 101 | pro-inflammatory cytokines 5 |
|---|---|
| Portsmouth, maternal nutrition and fetal programming 97 | obesity and cardiovascular disease and 234 |
| potassium 306 | pregnancy 74, 76 |
| blood pressure and intake of 242 | pre-eclampsia and 77 |
| infant/preschool children requirements 153 | pro-inflammatory eicosanoids and pregnancy 67 |
| sources/functions/deficiency symptoms 306 | Project Viva 101, 102, 103 |
| poverty | prolactin 127–8 |
| children 163–5 | proliferation phase of carcinogenesis 243 |
| elderly 278 | promotion (health and healthy eating) |
| see also social (and socioeconomic/sociocultural) factors | breastfeeding 140-1 |
| Prader–Willi syndrome 175, 176 | children |
| preconceptual and periconceptual period 2, 52–7 | infants/preschool 163 |
| folic acid supplementation 55 | school-age and in school 168–9 |
| premature labour relating to 72–3 | prospective cohort study 25, 26 |
| pre-eclampsia xxiii, 75, 76, 77–9 | Prospective Diabetes Study (UK) 234–5 |
| periodontal disease and 75 | prostate cancer 247, 250, 257, 258 |
| preterm delivery 72 | protein 304–6 |
| pregnancy 52–7, 61–93 | animal and plant sources, availability by world region |
| 1st trimester 61–2 | 15, 16 |
| 2nd trimester 62 | breast milk 126–7, 130 |
| 3rd trimester 62 | demands/requirements 276 |
| adolescent see adolescence | adolescents 197–8 |
| breast in 127–8 | determination 9 |
| cravings and aversions in 82–4 | infant/preschool children 153 |
| fetal programming and interventions in 115 | injury and infection impacting 5 |
| high-risk 84–8 | pregnancy 65–6, 66–8 |
| iron deficiency 7, 64, 68, 69, 83, 84, 104, 210 | digestion and absorption 306 |
| multiple see multiple pregnancy | formulas feeds, hydrolysed 148 |
| nausea and vomiting see nausea and vomiting | functions 304 |
| nutrient requirements 66–71 | maternal restriction, fetal programming and 105 |
| outcome relating to diet 71–9 | osteoporosis and 285–6 |
| physiological demands of 62–6 preparing for <i>see</i> preconceptual and periconceptual period | structure 305–6 |
| see also breastfeeding; conception; fetus; lactation; teratogens | weight-loss diets and 225–6 protein–energy malnutrition xxi, 4, 7, 8 |
| premature ageing syndromes 271 | children 153 |
| premature ageing syndromes 271 premature infant (preterm) infant, formulas for 145–6 | elderly 278 |
| premature labour (preterm labour) labour and delivery 72–6 | proto-oncogenes 243 |
| multiple pregnancy 72, 86 | psychological morbidity of childhood obesity 176 |
| preschool children see infants and preschool children | psychosocial development |
| prevalence (and prevalence rates) xxiii, 22, 23 | adolescent 196 |
| obesity see obesity | infant/preschool children 152 |
| osteoporosis 282–3 | puberty xxiii, 191–2 |
| see also epidemiology | females, and leptin 40 |
| prevention | growth see growth |
| malnutrition 8 | pubic hair development 191, 193 |
| elderly 279–81 | public health nutrition/interventions xxiv |
| obesity in children 180–2 | and fetal programming 115 |
| primate studies of caloric restriction and lifespan 273 | Public Health Responsibility Deal (DoH) 182 |
| processed meat and cancer 252 | publication bias xxiv |
| progenitor cells, embryonic 95, 108 | pyridoxine see vitamin B6 |
| progesterone | |
| menstrual cycle 37 | quantitative insulin sensitivity check index (QUICKI) 229 |
| in pregnancy | quercetin 260, 261 |
| breast and 127 | |
| nausea and vomiting and 80 | randomized controlled trials 24, 25, 26-7 |
| programmed cell death xvi, 270, 272 | reactive oxygen species (ROS) xxiv, 41, 42 |
| programming, fetal developmental (and disease in later life) xxiii, | ageing and 271 |
| 94–122 | and fertility |
| alcohol and 86 | females 41, 41-3, 42 |
| concept of programming 94–5 | males 48 |
| criticisms of hypothesis 102–3 | see also free radicals |
| implications of hypothesis 115–18 | recommended daily amounts (RDIs) 28 |
| lifespan and 275 | red blood cells (erythrocytes), examination in anaemia 290, 291 |
| mechanisms 106–15 | red meat and cancer 252–3 |







| reference nutrient intake (RNI in UK) xxiv, 29, 30, 31 | FTO 175, 176 |
|---|--|
| elderly, energy and protein 276 | skeletal growth in adolescence/puberty 193–6, 198–9 |
| lactating women 129 | skeletal muscle mass and ageing 269 |
| lower (LRNI) xxi | skinfold thicknesses xxiv, 15 |
| relative risk 22–4 | small intestine disorders, elderly 289 |
| renal see kidney | SMART lunchbox intervention 181–2 |
| reproductive function and system see fertility; genitalia | smell, sense of, elderly 279 |
| respiratory changes in pregnancy 64–5 | smoking (tobacco) 212–13 |
| resting metabolic rate see basal metabolic rate | adolescent 212–13 |
| restriction (diet/food/food groups) | bone health and 213 |
| adolescent 203 | cancer and 244 |
| caloric, and lifespan 273–5 | cardiovascular disease and 233 |
| in obesity treatment 225–6 | periodontal disease in pregnancy and 75 |
| by parents 167 | snacking (children) |
| retinol 307 | adolescent 202, 203 |
| activity equivalents xxiv, 52 | drug use and 214 |
| alcohol and 211 | advertising and 166 |
| pregnancy and 52, 53, 54 | social (and socioeconomic/sociocultural) factors |
| sources and functions 307 | bottle-feeding rather than breastfeeding 138–9 |
| symptoms of deficiency 307 | fetal growth 96 |
| retrospective cohort study 25, 26 | periconceptual period and 52 |
| rhesus monkey, caloric restriction and lifespan 273 | pica 83 |
| riboflavin, sources/functions/deficiency symptoms 307 | see also poverty |
| rimonabant 226 | social isolation, elderly 278–9 |
| rodent studies of caloric restriction and lifespan 273 | sodium 240–2, 308 |
| rooting reflex xxiv, 124 | hypertension and 240–2 |
| | adult offspring hypertension and maternal sodium intake of 105 |
| safrole 244, 255 | infant/preschool children 154 |
| salt 221 | sources/functions/deficiency symptoms 306 |
| infant/preschool children 154 | see also salt |
| sensitivity vs. resistance 240–1 | soil consumption in pregnancy 83, 84 |
| stomach cancer and 27 | SOLINUT study 279 |
| see also sodium | South Korean migrants in US, cancer 247 |
| saturated fats/fatty acids 235-6, 302, 303 | Southampton Women's Study 103, 158, 164 |
| Scandinavia | soy formulas 148 |
| breastfeeding trends 137 | spermatogenesis 44–5 |
| Livsmedelsverket Keyhole 223 | obesity and 47 |
| school | see also semen quality |
| breakfast 169–70 | spina bifida xxiv, 54, 56, 293 |
| lunches/meals 168-9, 181-2 | sport, adolescents 204–6 |
| obesity prevention and 181–2 | stanols and stanol esters, plant 236 |
| school-age children 167–70 | STAT3 40-1 |
| screening for malnutrition, elderly 278, 280 | statins xxiv, 236–7 |
| second-stage milk 144 | stature see height |
| sedentary activity see physical activity | status (nutritional) xxii, 13–20 |
| selection bias 21 | adolescent 205 |
| selenium | pregnancy and 209–10 |
| elderly, supplementation 287 | assessment 13–20 |
| male fertility and 49 | biomarkers xvii, 18–20 |
| self-esteem and bulimia nervosa 208 | cytokines impacting on 5 |
| semen quality | definition 2 |
| antioxidants and 48-9 | overall (of individual) 9–10 |
| organic food vs. pesticide exposure 51 | stem cells 270 |
| see also spermatogenesis | embryonic 95 |
| senescence xxiv | Steno-2 study 234 |
| cellular see cells | stereotypes in breastfeeding, negative 139 |
| see also ageing | stillbirth 71–2 |
| serotonin and anorexia nervosa 207 | stomach (gastric) |
| Sertoli cells xxiv, 44, 45 | cancer 245, 247, 255, 257, 258, 260 |
| Seventh Day Adventists, cancer 247 | salt and 27 |
| sexual maturation 191–3 | elderly, disorders 289 |
| sheep milk (formula feed) 148 | in pregnancy 65 |
| single nucleotide polymorphisms (SNPs) xxiv, 12–14 | surgery in obesity (bariatric surgery) xvii, 226, 227 |
| ApoF 14 | stool bulking by fibre and colorectal cancer, 257 |







| stress (psychological) and preterm delivery 62 | tooth caries and fluoride, infant/preschool children 154, 155, 155 |
|--|--|
| stroke (cerebrovascular accident) xxiv, 232 | trace nutrients see micronutrients |
| stunting 153 | traffic light classification of foodstuffs 179, 223 |
| subfertility see infertility and subfertility | trans-fatty acids xxv, 235–6, 302 |
| substance use/abuse see alcohol; drug abuse; smoking | transgenerational transmission of disease risk 117–18 |
| suckling 123, 124, 128, 129 | trauma xxv |
| sugar-sweetened beverages 175 | inflammatory response see inflammatory response |
| see also carbohydrates | metabolic response 4–6 |
| sulphopharane and cancer 261 | triacylglycerides see triglycerides |
| superoxide dismutase 43, 48, 238, 271 | Trials of Hypertension Prevention study 242 |
| supplementation xxiv | triglycerides (triacylglycerides) 303 |
| antioxidants | breast milk 126 |
| ageing and lifespan and 275–6 | tuberculosis, countries xxv, 143 |
| male fertility and 48–9 | tumour see cancer |
| minerals | tumour suppressor genes xxv, 243 |
| calcium, elderly 284–5 | twin pregnancy see multiple pregnancy |
| fluoride, children 154 | twin studies and thrifty phenotype hypothesis 107 |
| infant/preschool children 155–6 | |
| iron in pregnancy 69 | UK see United Kingdom |
| selenium, elderly 287 | ulcers, peptic 289 |
| zinc, elderly 287–8 | UN see United Nations |
| selenium, elderly, minerals 287 | undernutrition 4–8 |
| vitamins | adolescent drug use and 214 |
| folic acid 54, 55 | fetal programming and 97, 99, 100, 101, 103, 107, 108, 109, |
| infant/preschool children 155–6 | 110, 112, 113, 114, 115, 117, 118 |
| vitamin A 8 | global dimensions 104, 106 |
| vitamin B6, and homocysteine 238 | United Kingdom (UK) 28–31 |
| vitamin B12 see vitamin B12 | breastfeeding trends 136–7, 138–9 |
| vitamin C, and pre-eclampsia prevention 78 | DRVs see diet, reference values |
| vitamin D see vitamin D | elderly malnutrition 277 |
| vitamin E see vitamin E | healthy nutrition advice (adults) 221–2 |
| see also multivitamin supplements | infant/preschool children |
| supply and demand 2 | poverty and 165 |
| compromised 6 | supplements 156 |
| increased demand 4 | vitamin D deficiency 160–1 |
| surgery, bariatric xvii, 226, 227 | weaning 159 |
| sweat gland number 85 | maternal nutrition and fetal programming 97, 103 |
| Swedish Food Circle 221 | National Diet and Nutrition Survey see National Diet and |
| syndrome X (insulin resistance syndrome; metabolic | Nutrition Survey |
| syndrome) xxi, 100, 176, 229-30 | National Health and Nutrition Examination Study |
| systematic reviews xxiv-xxv, 27–9 | (NHANES) 28, 70, 171, 174, 203, 224, 293 |
| | National School Fruit and Vegetable Scheme 168 |
| Γ cells and ageing 287, 288 | obesity |
| aste, elderly 279 | childhood interventions 181–2, 182 |
| ea and cardiovascular disease 239, 240 | studies from childhood to adulthood 177–8 |
| eenagers see adolescence | pregnancy and excess weigh management 74 |
| elevision | Prospective Diabetes Study (UKPDS) 234–5 |
| advertising 165, 166 | RNI see reference nutrient intake |
| watching by children as sedentary activity 173 | School Breakfast Program 165 |
| elomere xxv | United Nations (UN) |
| shortening 271, 272–3 | Children's Fund and breastfeeding (UNICEF) 135, 140 |
| eratogens xxv | Millennium Development Goal 164 |
| vitamin A as 53, 54 | Millennium Project 164 |
| esticular non-descent (cryptorchidism) xviii, 39, 50 | United States |
| estosterone 44–5 | cancer and migrant studies in 246–7 |
| hermogenesis xxv | Cardiovascular Health Study 290 |
| hiamin, sources/functions/deficiency symptoms 307 | dietary reference values 31–2 |
| Thousand Families Study (Newcastle) 177–8, 178 | folate fortification 55 |
| hrift, metabolic (concept) 228 | infants/preschool/young children |
| thrifty genotype 106, 107, 228 | food and drink advertising 165 |
| thrifty phenotype 106–7, 108, 116 | supplements 156 |
| issue remodelling (in fetal programming) 108–9 | Nurses Health Study 72, 100, 131, 248, 251, 255, 256 |
| obacco smoking see smoking | obesity prevention in children 182 |
| ocopherols see vitamin E | School Breakfast Program 169–70 |







| Universal Salt Iodization (USI) program 69 | elderly 276 |
|---|---|
| unsaturated fatty acids 302 | falls and 285 |
| see also monounsaturated fatty acids; polyunsaturated fatty | supplements 284–5 |
| acids | excess consumption 3 |
| UPBEAT trial (UK) 74 | infant/preschool children |
| uracil misincorporation xxv, 259 | deficiency 160–1 |
| USA passim see United States | supplements 156 |
| | pregnancy 70–1 |
| vascular dementia 294 | receptor, and osteoporosis 112 |
| vascular endothelium and atherosclerosis 231 | sources/functions/deficiency symptoms 307 |
| vegans xxv, 167, 204 | vitamin E (tocopherols) 239, 307 |
| vegetables see cruciferous vegetables; fruit and vegetable | ageing and supplementation with 275-6, 288 |
| consumption | cancer prevention and 258 |
| vegetarianism xxv | cardiovascular disease prevention and, and its |
| adolescent 199, 204, 205 | supplementation 239 |
| very-low-calorie diet (VLCD) 226 | male fertility and 48–9 |
| vitamin(s) 308 | pre-eclampsia management and supplementation with 78 |
| fat-soluble 307 | sources/functions/deficiency symptoms 307 |
| infant/preschool children 154, 155-6 | status determination 19 |
| lactation and 129–30 | vitamin H (biotin), sources/functions/deficiency symptoms |
| sources/functions/deficiency symptoms 308 | 307 |
| supplements see supplementation | vitamin K, sources/functions/deficiency symptoms 307 |
| water-soluble 307 | Vitamin Study Group, MRC 11, 54, 55 |
| vitamin A 307 | VITATOPS trial 238 |
| cancer and 248 | vomiting in pregnancy see nausea and vomiting |
| deficiency | |
| adolescent 199 | waist circumference xxv, 15 |
| combating 8 | obesity classification 223 |
| infant/preschool children 159 | waist–hip ratio xxv, 223 |
| symptoms 307 | cancer and 249–50 |
| excess consumption 3 | water-soluble vitamins 307 |
| lactation and 130 | water supply |
| pregnancy and 52–4 | endocrine-disrupting chemicals 49 |
| sources and functions 307 | fluoridation 155 |
| status determination 19 | WCRF (World Cancer Research Fund) 249, 251, 252, 253, 255 |
| see also carotenoids; retinol | 256, 257, 258, 259, 260 |
| vitamin B1 (thiamin), sources/functions/deficiency symptoms 307 | weaning see complementary feeding |
| vitamin B2 (riboflavin), sources/functions/deficiency | weight |
| symptoms 307 | adolescents |
| vitamin B3 (niacin), sources/functions/deficiency symptoms 307 | control 203–4 |
| vitamin B5 (pantothenic acid), sources/functions/deficiency | energy requirements and 198 |
| symptoms 307 | gain 190 |
| vitamin B6 (pyridoxine) 307 | birth see birth weight |
| elderly 276 | dietary approaches to losing 225–6, 234 |
| and Alzheimer's disease 294 | infant/preschool children, rate of gain 151 |
| sources/functions/deficiency symptoms 307 | in polycystic ovary syndrome |
| supplementation, homocysteine and 238 | losing, effectiveness 38 |
| vitamin B9 see folic acid | maternal, management 52 |
| vitamin B12 (cobalamin) | pre-pregnancy gain, and premature birth 72 |
| elderly 276 | pregnancy gain 63 |
| cognitive problems and 294 | management 74 |
| deficiency 292, 294 | multiple pregnancy and 86 |
| folate fortification and status of 56 | premature birth and 72 |
| supplementation 276 | see also obesity |
| homocysteine and supplementation with 238 | Westernized countries see developed countries |
| sources/functions/deficiency symptoms 307 | wheat flours/products, folic acid fortification 55–6 |
| vitamin C (ascorbic acid) 307 | whey-rich formula milks 144–5 |
| cancer and 258 | WHO see World Health Organization |
| elderly 276 | women see females; girls |
| male fertility and 48–9 | World Cancer Research Fund (WCRF) 249, 251, 252, 253, 255 |
| pre-eclampsia and 78 | 256, 257, 258, 259, 260 |
| sources/functions/deficiency symptoms 307 | World Health Organization (WHO) |
| vitamin D (calciferol) 307 | on breastfeeding 135, 136, 137, 140 |
| adolescent bone growth and 195 | and HIV disease 142–3 |







Collaborative Health Behaviours in School-Aged Children Study 203 on food and drink advertising viewed by children 166 on healthy nutrition 221 metabolic syndrome diagnostic criteria 230 on obesity 223–4 prevention in childhood 180 worldwide dimensions see global/worldwide dimensions

xenobiotic metabolism xxiii, 214, 244, 261

zinc 306
adolescent (incl. deficiency) 198
determination (of status) 19
elderly, supplementation 287–8
immunity and 117, 160
infant/preschool children, deficiency 160
male fertility and 48
pregnancy 69
sources/functions/symptoms of deficiency 306



