

Index

Page numbers in *italics* refer to illustrations; those in **bold** refer to tables

- abdominal contour assessment 52, 52
- abdominal jiggling 321–322, 321
- abdominal lifting 320–321, 320
 - advantages and disadvantages 321
 - effects of 321
 - for malposition, cephalopelvic disproportion and macrosomia 142–143, 143
 - for prolonged prelabor and latent phase 117–119, 118
- abdominal palpation
 - fetal descent assessment 58, 58
 - Leopold's maneuvers 55–57, 56, 57, 70
- abuse
 - childhood abuse survivors 27–28
 - emotional dystocia during labor 159–161
 - during childbirth 31
 - sexual abuse 27–28
- Active Management of Third Stage Labor (AMTS) 218–219
- active phase
 - first stage labor 88, 126–127
 - second stage labor 171–175
 - see also* prolonged active phase of first stage labor; second stage dystocia
- acupressure 343–344, 343
 - contraction stimulation 36, 151
 - effectiveness 344
 - indications and contraindications 344
 - labor induction **103**
 - mechanisms of action 344
- acupuncture 344
 - contraction enhancement 151
 - intrapartum pain relief 251–252, 251
 - labor induction **105**
- anxiety 160
 - effects during labor 19–20, 19
 - psycho-emotional measures 24–31

The Labor Progress Handbook: Early Interventions to Prevent and Treat Dystocia, Fourth Edition.
Penny Simkin, Lisa Hanson and Ruth Ancheta.
© 2017 John Wiley & Sons, Inc. Published 2017 by John Wiley & Sons, Inc.

- anxiety (*cont'd*)
 - before labor 24–26
 - during labor 26–27
 - trauma-informed care 27–31, **29–30**
 - see also* emotional dystocia
- aortic compression 220
- artificial rupture of membranes (AROM) 234
 - with malpositioned fetus 134–135
- assessments during labor 66–89
 - contraction quality 68–70, 80
 - hydration 66–67
 - see also* labor, progress assessment
- internal assessments 75–77
 - bony pelvis 87
 - caput formation 86
 - cervix 79–81
 - fetal position 83–85, *84*, 87
 - presenting part 81
 - stations of descent 81–83, *81*
 - vaginal examinations 77–79, 78, 87
- nourishment 66–67
- psychology 67
- vital signs 69
 - see also* auscultation; fetal monitoring
- asymmetrical positions 278, 299, 300
 - effects of 299
 - indications and contraindications 299
 - prolonged active phase of first stage 140, *141*
 - second stage dystocia 179–182, *181*
 - see also* lunges
- asynclitism 85, *85*, 87, 116
 - anterior *116*, *131*, *190*
 - correction techniques 116–119, *117*, *118*
 - persistent 130, *130*, *131*, 187–192
 - posterior *116*, *130*, *191*
 - second stage 187–192, *192*
 - see also* fetal malposition
- auscultation 36, 37, 71–74
 - candidates for intermittent auscultation 72
 - counting method 73
 - fetal stethoscope use 37, 53, 72
 - frequency of 73
 - handheld Doppler device use 72
 - point of maximum fetal heart tone intensity (PMI) location 53–54, 53, 54
 - technique 73
 - timing of 73
 - see also* fetal heart rate (FHR)
- baby *see* newborn
- back massage 339, 339

- back pain management 183–186, 186–189, 350–364
 - birth ball 360–362, 361
 - cold and heat 357–359, 358
 - counterpressure 350–351, 350
 - Cook's technique No. 1 (ischial tuberosities) 354–355, 355
 - Cook's technique No. 2 (perilabial pressure) 355–357, 356
 - double hip squeeze 351–352, 351
 - hydrotherapy 189, 359, 359
 - knee press 353–354, 353
 - movement 359, 360
 - sterile water injections 189, 252–253, 364
 - TENS 189, 362–364, 362
 - see also* pain
- bath use during labor 334, 335
 - see also* hydrotherapy
- bearing down *see* pushing
- Belly Mapping 59–62, 59–61
- bimanual compression 220, 220
- birth ball 360–362, 361
- birth environment 21
- birth plan 25–26
- birth positions
 - hands and knees position 245, 245, 249
 - perineal management 241
 - versus pushing positions 179
 - see also* maternal positions and movements
- birth sling 303–304, 303
- birthing rope 305, 305
- Bishop scoring system 65–66, 66, 100, 101
- bladder distention 266
- body temperature during labor 69, 264–265
 - epidural effects 264
- breast crawl 222
- breast stimulation, contraction enhancement 150, 233
 - see also* nipple stimulation
- breastfeeding
 - encouragement practices 222–223, 223, 225
 - microbiota exposure 224
- breath-holding physiological effects
 - on fetus 172
 - on woman 172
- breathing techniques 364–366

- Canadian maternity care practice 5
- caput formation 86
- carboprost 220
- castor oil, labor induction 106
- catecholamines 18–19
 - effects during labor 19–20, 20
 - epidural effects on production 263–264

- cephalopelvic disproportion (CPD) 129–130
 - maternal positions and movements for 135–145, 194–199
 - abdominal lifting 142–143, 143
 - asymmetrical positions and movements 140, 141
 - forward-leaning positions 136, 137
 - lunges 199
 - side-lying positions 138–139, 138, 139, 140, 195, 196
 - squatting positions 196, 197
 - second stage dystocia and 193–199
 - time influence on 194
- cervix 64
 - assessment 79–80
 - prior to labor 64–65
 - dilation 80, 98–100
 - prenatal preparation for 102
 - factors that may prolong prelabor and latent phase 112
 - scar tissue 112
 - persistent cervical lip 81, 153
 - manual reduction 155, 238
 - maternal positions for 153–155, 154
 - rigid os 80–81
 - ripening 64–65
 - techniques to hasten ripening 102, **103–106**
 - stenosis management 233–234
 - swollen 153
 - maternal positions for 153–155, 154
 - reducing 238
 - zipper cervix 80
 - management 233–234
- cesarean delivery, neonatal microbiome impacts 224–225
- childhood abuse survivors
 - emotional dystocia during labor 159–161
 - sexual abuse (CSA) 27–28
- closed knee–chest position 120, 120, 298, 298–299
 - effects of 298
 - indications and contraindications 299
- cold therapy 331–333, 332
 - back pain 357–359, 358
- complementary alternative medicine (CAM), cervical ripening techniques 102, **103–106**
- continuous electronic fetal monitoring (CEFM) *see* electronic fetal monitoring (EFM)
- contractions
 - non-progressing contractions of prelabor 98
 - painful early contraction management 115–121
 - progressing contractions of labor 98
 - quality assessment 68–70, 80
 - see also* inadequate contractions
- controlled cord traction (CCT) 218

- Cook's counterpressure techniques
 - No. 1 (ischial tuberosities) 354–355, 355
 - No. 2 (perilabial pressure) 355–357, 356
- coping
 - assessment 155–156
 - with pain 23–24, 23
 - second stage 204
- Coping Algorithm 24
- cord *see* umbilical cord
- counterpressure 186, 350–351, 350
 - Cook's technique No. 1 (ischial tuberosities) 354–355, 355
 - Cook's technique No. 2 (perilabial pressure) 355–357, 356
- criss-cross back massage 339, 339
- crowning of the fetal head
 - guidance through 241–242
 - hand skills to protect the perineum 242–243, 242
- dangle position 183, 184, 196, 302–304, 303
 - effects of 304
 - indications and contraindications 304
- dehydration 147–148
- delayed clamping (DC) of the umbilical cord 214–216
- diffuse pushing 174–175, 174, 206
- digital rotation of the fetal head 235–236, 236
- directed pushing 367
- Doppler device *see* handheld Doppler device
- double hip squeeze 183–184, 184, 187, 351–352, 351
- doula presence 329, 345–346, 345
 - effectiveness of 346
- drinking *see* hydration during labor
- electronic fetal monitoring (EFM) 36
 - continuous (CEFM) 37–39, 71, 74
 - enhancing maternal mobility 37–42, 38, 41
 - in bath/shower 39–40, 39, 334–336, 335
 - intermittent 39–40
 - wireless telemetry 40–42, 41, 42
 - see also* fetal heart rate (FHR); fetal monitoring
- emotional dystocia 155–162, 160
 - addressing concerns 157–158
 - childhood abuse survivors 159–161
 - coping assessment 155–156
 - indicators of 156–157, 205
 - management of 158–159
 - poor relationship with staff 161
 - predisposing factors 157
 - second stage 204–207
 - unidentified cause 161–162
 - see also* anxiety

- emotional state
 - assessment of 348
 - hormonal influences 17–20, 19
 - painful/prolonged prelabor or latent phase relationship 113–114
 - wellbeing versus distress 21–24
 - see also* emotional dystocia; psycho-emotional measures
- endorphins 18
 - following birth 213
- environment for birth 21
- epidural analgesia 261–262
 - management of deleterious effects 266–273
 - attending to emotional needs 272–273
 - delayed pushing 269, 271
 - encouraging behavior resembling non-anesthetized woman 267–272, 268
 - guided physiologic pushing 271–272
 - information provision to woman 266–267
 - shortening duration of exposure 267
 - multisystem effects on labor progress 263–266
 - endocrine system 263–264
 - genitourinary system 266
 - musculoskeletal system 265–266
 - nervous system 264–265
 - pushing positions with 176, 177, 269–271, 270
- evening primrose oil, labor induction **106**
- exaggerated lithotomy position 247–248, 248, 278, 310–311, 311
 - effects of 310
 - indications and contraindications 311
 - prolonged second stage 202–203, 203
- exaggerated Sims' position *see* semiprone position
- exhaustion, maternal 149
 - positions for 149
- fetal head descent 194
- fetal heart rate (FHR) 71–75
 - accelerations defined 74–75
 - baseline definitions 74
 - counting method 73
 - decelerations defined 75
 - point of maximum heart tone intensity (PMI) location 53–54, 53, 54
 - three-tiered FHR interpretation system 75, 76
 - variability defined 74
 - see also* auscultation; electronic fetal monitoring (EFM)
- fetal malposition 62, 85, 129–143
 - artificial rupture of membranes and 134–135
 - identifying fetus likely to persist in malposition 63
 - maternal positions and movements for 135–145, 179–183, 192
 - abdominal lifting 141–143, 143
 - asymmetrical positions and movements 140, 141, 179–182, 181
 - forward-leaning positions 136, 137, 179, 180

- side-lying positions 138–139, 138, 139, 140, 182–183, 182, 195, 196
- squatting positions 179, 181, 183, 184, 196, 197
- supine and semisitting positions 142, 142
- nuchal hand 193, 193
- prolonged active phase and 129–132, 136
 - mechanisms of progress delay 132
- second stage dystocia and 178–187
- see also* asynclitism; fetal presentation and position; occiput posterior (OP) fetal position; occiput transverse (OT) fetal position
- fetal monitoring 70–71
 - enhancing maternal mobility during 36–42, 38, 41
 - see also* auscultation; electronic fetal monitoring (EFM); fetal heart rate (FHR)
- fetal presentation and position 50, 51, 86
 - assessment techniques 52, 129–130
 - abdominal contour 52, 52
 - abdominal palpation using Leopold's maneuvers 55–57, 56, 57
 - Belly Mapping 59–62, 59–61
 - internal assessment 83–85, 84, 87
 - point of maximal fetal heart tone intensity (PMI) location 53–54, 53, 54
 - problems of diagnosis during labor 133–134
 - ultrasound 129–130, 133–134
 - engagement assessment 58, 58
 - influencing position during labor 63
 - influencing position prior to labor 62–63
 - optimal fetal positioning (OFP) 107–108, 108
 - see also* fetal malposition
- fetal skull, digital assessment 84–85
 - fontanelles 85
 - molding 85
 - sagittal suture location 84–85, 84
- fetal stethoscope use 37, 53, 72
- fetal weight estimation 64
- fetus to newborn transition 211–212
 - see also* newborn
- fight-or-flight response 19–20
- first stage of labor
 - active and latent phase features 88, 126–127
 - onset 96–99
 - progress assessment 87–88, 126–127
 - see also* prolonged active phase of first stage labor; prolonged latent phase
- foot massage 341–343, 341, 342
- forward-leaning positions 278, 292–293, 292
 - effects of 293
 - indications and contraindications 293
 - kneeling 293–295, 294
 - prolonged active phase of first stage 136, 137
 - prolonged prelabor or latent phase 116, 117
 - second phase dystocia 179, 180
 - sitting 108, 179, 180, 197, 290–292, 291

fourth stage of labor 212–213, 221–222
 keeping mother and baby together 221–222
 Friedman, E. A. 14, 97

Gaskin maneuver 249
 geographical variation in maternity care practice 5
 gestational age 71
 guarding of the uterus 217, 218

half-squatting position 304–306, 305
 effects of 306
 indications and contraindications 306

hand massage 340–341, 340

handheld Doppler device 72

waterproof, for use with hydrotherapy 39–40, 39
 hands and knees position 245, 245, 249, 295–296, 295
 effects of 295
 flexion and rocking of hips and knees 315–316, 315
 indications and contraindications 296

hands-on approach to delivery 242–243
 versus hands-poised approach 243

heat therapy 330–331, 330
 back pain 357–359, 358

herbal tea, labor induction 106

hip sifting 314, 314

homeopathy, labor induction 105

hormonal influences on labor 17–20, 19

hydration during labor 66–67
 dehydration 147–148
 overhydration 148

hydrotherapy 329, 333–337, 333, 335
 back pain 189, 359, 359
 contraction enhancement 151–153, 152
 effectiveness 336–337
 fetal monitoring and 39–40, 39, 40, 333, 334–336, 335
 indications and contraindications 336
 mechanisms of action 334
 second stage dystocia 198

inadequate contractions 145, 203–204
 causes 145–150

dehydration 147–148
 exhaustion 149, 149
 immobility 145–147
 medication 147
 overhydration 148
 uterine lactic acidosis 149–150

techniques for stronger contractions 35–36, 35, 150–153, 203–204
 acupressure or acupuncture 36, 151
 breast stimulation 150, 233
 hydrotherapy 151–153, 152

- rollover sequence 145, 146
 - walking and position changes 151
- intermittent auscultation *see* auscultation
- intrauterine pressure catheter (IUPC) 39
- kangaroo care (KC) 222
- ketonuria 67
- knee press 188, 353–354, 353
- kneeling positions 117, 293–295, 294
 - asymmetrical kneeling 141, 181, 300
 - contraindications 295
 - indications 294
 - kneeling lunges 141, 186, 199, 316, 360
 - prolonged active phase of first stage 137, 144, 149, 154
 - prolonged second stage 177, 179, 180, 189
- labor
 - hormonal influences 17–20, 19
 - normal 10, 11–13
 - onset 96–97
 - Bishop scoring system 65–66, 65, 100, 101
 - definition 97
 - progress assessment 50, 66
 - active phase features 88, 126–127
 - first stage 87–88
 - latent phase features 88, 126
 - second stage 88–89
 - see also* assessments during labor
 - six ways to progress 99–100
 - versus prelabor 96–101
 - see also* first stage of labor; second stage of labor
- labor dystocia 14–15
 - care plan 3
 - causes 1–2, 15, 16
 - definition 14–15
 - see also* emotional dystocia; prolonged active phase of first stage labor; prolonged latent phase; second stage dystocia
- labor induction 102
 - expert non-medical techniques 105–106
 - self-administered techniques 103–104
- lactic acidosis, uterine 149–150
- lap squatting 306–308, 307
 - contraindications 308
 - effects of 306–307
 - indications 307
- latent phase
 - first stage labor 88, 126
 - second stage labor 169–170, 169, 244
 - persistence of 171
 - see also* prolonged latent phase
- lateral positions *see* side-lying positions

- Leopold's maneuvers 55, 70
 - abdominal palpation technique 55–57, 56, 57
- lidocaine, perineal application 254
- low-birth-weight neonates, kangaroo care (KC) 222
- lunges 140, 316–317
 - direction of 317
 - indications and contraindications 285, 317
 - kneeling 141, 186, 199, 316, 360
 - side-prone/semiprone 139, 139, 199, 284, 284, 285
 - standing 141, 185, 199, 316
- McRoberts' maneuver 247–248, 248
- macrosomia 129–130, 193–194
 - maternal positions and movements for 135–145
 - see also* cephalopelvic disproportion (CPD)
- malposition *see* fetal malposition
- manual rotation of the fetal head 61, 235–238, 237
- massage 337–343
 - criss-cross back massage 339, 339
 - effectiveness 338
 - hand massage 340–341, 340
 - indications and contraindications 337
 - perineal 243
 - prenatal 239–240, 239
 - shoulder mini-massage 338–339
 - three-part foot massage 341–343, 341, 342
 - uterine 218–219
- maternal mobility, fetal monitoring and 36–42, 38, 41
 - see also* maternal positions and movements
- maternal musculoskeletal variations 109
- maternal positions and movements 278–335
 - back pain management 359, 360
 - contraction enhancement 151
 - effects on labor 278–311
 - side-lying positions 279–288
 - exhausted women 149
 - for prolonged active phase of first stage 135–145
 - abdominal lifting 142–143, 143
 - asymmetrical positions and movements 140, 141
 - forward-leaning movements 136, 137
 - lunges 139, 141
 - side-lying positions 138–139, 138, 139, 140
 - supine and semisitting positions 142, 142
 - for prolonged prelabor and latent phase 117–121
 - closed knee–chest position 120, 120
 - open knee–chest position 119–120, 119
 - side-lying release 120–121
 - for second stage dystocia 176–187
 - asymmetrical positions 179–182, 181
 - cephalopelvic disproportion and 194–199

- dangle position 183, 184, 196
- exaggerated lithotomy position 202–203, 203, 247–248, 248
- forward leaning positions 179, 180
- lunges 185, 186, 199
- side-lying positions 182–183, 182, 195, 196
- squatting positions 179, 181, 196, 197
- supine positions 200–202, 201, 202
- frequent position changes 329
- importance of 33–35
- persistent cervical lip management 153–155, 154
- pushing positions versus birth positions 179
- rhythmic movements, benefits of 323–325, 325, 329
- swollen cervix management 153–155, 154
- uncontrollable premature urge to push and 143–145, 144, 145
- see also* asymmetrical positions; kneeling positions; side-lying positions; sitting positions; squatting positions; supine positions
- maternity care practice, geographical variation 5
- meconium 71
- Methergine 219
- microbiome 224
 - infant microbial health support 223–225, 224
- midwife as labor support provider 346–347
- misoprostol 219
- molding, fetal skull 85
- musculoskeletal variations, maternal 109
- nasopharynx suctioning, neonate 213–214
- neocortex overstimulation 17–18
- neuraxial analgesia 261–262
- newborn
 - breastfeeding encouragement 222–223, 223, 225
 - care of 213–216
 - delayed clamping of the umbilical cord 214–216
 - oral ad nasopharynx suctioning 213–214
 - delayed bathing 223
 - keeping mother and baby together 221–222
 - kangaroo care (KC) 222
 - skin-to-skin contact (SSC) 221–222, 225
 - microbial health support 223–225, 224
 - routine assessments 225–226
 - transition from fetus 211–212
- nipple stimulation
 - contraction enhancement 36, 150, 233
 - labor induction 103
- nitrous oxide 254
- normal labor 10, 11–13
- nourishment during labor 66–67, 148
- nuchal cord, tight 216
- nuchal hand 193, 193
- nurse as labor support provider 346–347

- occiput posterior (OP) fetal position 62, 130–131, 132, 178
 - identifying fetus likely to persist in OP position 63
 - influencing position during labor 63, 281
 - influencing position prior to labor 62–63
 - manual or digital rotation 187, 235–238, 236, 237
 - see also* fetal malposition
- occiput transverse (OT) fetal position 131, 132, 178
 - see also* fetal malposition
- open knee–chest position 119–120, 119, 296–298, 297
 - effects of 296
 - indications and contraindications 298
- optimal fetal positioning (OFP) 107–108, 108
- oral suctioning, neonate 213–214
- Ottawa Hospital Second State Protocol 175, 176
- overhydration 148
- oxytocin 18
 - epidural effects on production 263
 - levels following birth 213
 - synthetic 219
- pain
 - assessment of 22, 22
 - coping ability assessment 23–24, 23
 - management approaches 22
 - acupuncture 251–252, 251
 - cold and heat 357–359, 358
 - nitrous oxide 254
 - non-pharmacologic method guidelines 328–329
 - sterile water injections 252–254, 253, 364
 - topical perineal anesthetic 254
 - see also* back pain management; epidural analgesia
 - painful prolonged prelabor/latent phase 112–114
 - cervical factors 112
 - early painful contraction management 115–121
 - emotional factors 113–114
 - iatrogenic factors 112
 - soft tissue factors 113
 - troubleshooting measures 114–115
 - versus suffering 21–22
 - see also* back pain management
- pelvic press 183–184, 185, 323, 324
- pelvic rocking 185, 200, 312–313, 312, 313, 360
 - advantages and disadvantages 313
- perineal management 239–243
 - perineal massage 243
 - prenatal 239–240, 239
 - second stage 240–243
 - guidance through crowning of the fetal head 241–242
 - hand skills 242–243

- maternal birth positions 241
- verbal support for spontaneous bearing down 240–241
- topical anesthetic application 254
- persistent asynclitism 130, 130, 131
 - early interventions 187–192
- persistent cervical lip 81, 153
 - manual reduction 155, 238
- paternal positions for 153–155, 154
- perspiration, fluid loss 67
- physical comfort measures during labor 32
- physiologic measures during labor 32–33
- Pinard stethoscope use 37, 53
- placenta delivery 212, 216
 - active management 218–221
 - physiologic (expectant) management 217–218, 217
- positions, maternal *see* maternal positions and movements
- post-term fetuses 71
- postpartum hemorrhage 219–221
- prelabor 96, 98
 - prolonged 101, 109, 110
 - prenatal preventive actions 102
 - support measures 109–111
 - therapeutic rest 232–233
- prolonged and painful 112–114
 - cervical factors 112
 - early painful contraction management 115–121
 - emotional factors 113–114
 - iatrogenic factors 112
 - soft tissue factors 113
 - troubleshooting measures 114–115
- versus labor 96–101
- preterm fetuses 71
- prolactin 19, 213
- prolonged active phase of first stage labor 127, 128, 234–238
 - artificial rupture of membranes 234
 - causes 128–129
 - digital or manual rotation of the fetal head 235–238, 236, 237
 - emotional dystocia 155–162
 - fetal and fetopelvic factors 129–143
 - malposition and cephalopelvic disproportion 129–135
 - maternal positions and movements for 135–143
- inadequate contractions 145–153
 - causes 145–150
 - management techniques 150–153
- observable characteristics of 127–128
- persistent cervical lip 153–155
 - manual reduction 155, 238
- swollen cervix 153–155
 - reduction of swelling 238
- see also* second stage dystocia

- prolonged latent phase 109, 110
 - cervical factors 109
 - early support measures 109–111
 - fetal factors 107
- painful and prolonged latent phase 112–114
 - cervical factors 112
 - early painful contraction management 115–121
 - emotional factors 113–114
 - fetal factors 107
 - iatrogenic factors 112
 - soft tissue factors 113
 - troubleshooting measures 114–115
- therapeutic rest 232–233
 - see also* inadequate contractions
- prostaglandins 18
- psycho-emotional measures 24–31
 - before labor 24–26
 - during labor 26–27
 - trauma-informed care 27–31, **29–30**
- psychological distress 21–24, **160**
 - assessment during labor 67
 - effects during labor 19–20, 19
 - see also* emotional dystocia
- psychosocial comfort measures 347–350
 - comforting sensory stimuli 348
 - fear reduction 349–350
 - reassurance and praise 348–349
- pushing 366–367
 - avoidance during latent phase 170
 - diffuse pushing 174–175, 174, 206
 - directed pushing 367
 - maximal maternal bearing-down efforts 171
 - physiological effects of 172
 - premature urge to push 144
 - maternal positions for 143–145, 144, 145
 - pushing positions 269–271, 270
 - versus birth positions 179
 - self-directed pushing 367
 - spontaneous bearing-down efforts 172–173, 173, 366
 - verbal support 240–241
 - with epidural blocks 176, 177, 269–271, 270
 - delayed pushing 269, 271
 - guided physiologic pushing 271–272
- rebozo sifting 321–322, 321
- rhythm, coping and 155–156
- rhythmic movements, benefits of 323–325, 325, 329
- rollover sequence 145, 146, 283
 - use with epidural 267–269, 268, 283

- sacrospinous ligament 81–82, 82
- scar tissue, cervix 112
- second stage dystocia 176
 - cephalopelvic disproportion and 193–199
 - maternal positions for 194–199, 195–200
 - emotional dystocia 204–207
 - exaggerated lithotomy position use 202–203, 203, 247–248, 248
 - fetal malposition strategies 178–193
 - asymmetrical positions 179–182, 181
 - dangle position 183, 184
 - forward leaning positions 179, 180
 - lateral positions 182–183, 182
 - persistent asynclitism 192
 - squatting positions 179, 181
 - inadequate contractions 203–204
 - supine position use 200–202, 201, 202
 - see also* shoulder dystocia
- second stage of labor 168
 - duration 243–245
 - time limits 175–176
 - manual rotation of the fetal head 63
 - onset 243–244
 - Ottawa Hospital Protocol 175, 176
 - perineal management 240–243
 - phases of 168–171
 - active phase 171–175
 - latent phase 169–171, 169, 244
 - progress assessment 88–89
 - prolonged *see* second stage dystocia
- self-directed pushing 367
- semiprone position 138, 279–280, 280, 282, 282
 - benefits of 280
 - for vaginal examination 78, 78
 - prolonged active phase of first stage 139, 144
 - with inadequate contractions 145, 149
 - with persistent cervical lip 153, 154
 - prolonged prelabor 108
 - prolonged second stage 182–183, 182, 196
 - semiprone lunge 139, 139, 284, 284, 285
 - indications and contraindications 285
 - see also* side-lying positions
- semisitting positions 278, 288, 289
 - prolonged active phase of first stage 142, 142
 - second stage dystocia 177, 200–202, 201, 202
- sexual abuse 27–28
- sheet “pull-to-push” 309–310, 309
- shoulder dystocia 203, 244–245
 - maneuvers 246–250, 246
 - Gaskin maneuver 249
 - McRoberts’ maneuver 247–248, 248

- shoulder dystocia (*cont'd*)
 - somersault maneuver 249–250, 250
 - suprapubic pressure 248–249, 249
 - precautionary measures 245–246
 - warning signs 246
- shoulder mini-massage 338–339
- shower use during labor 333, 333
 - see also* hydrotherapy
- side-lying positions 278, 279–288, 279–280
 - contraindications 282–284
 - effects of 280
 - indications 282
 - prolonged active phase of first stage 138–139, 138, 139, 140
 - second stage dystocia 182–183, 182, 195, 196
 - side-lying release 120–121, 140, 140, 285–287, 286
 - contraindications 288
 - effects of 287
 - indications 287–288
 - side-prone lunge 139, 139, 199
- sitting positions 149, 207
 - leaning forward 108, 179, 180, 197, 290–292, 291
 - contraindications 292
 - indications 291
 - upright 108, 195, 202, 289–290, 290
 - contraindications 290
 - indications 289–290
 - see also* semisitting positions
- skin-to-skin contact (SSC) 221–222, 225
- slow dancing 186, 200, 318–319, 319, 360
 - advantages and disadvantages 319
 - with electronic fetal monitoring 38
- somersault maneuver 249–250, 250
- Spinning Babies approach 16–17
- squatting positions 179, 181, 197, 300–308, 301
 - effects of 301–302
 - half-squatting, lunging, and swaying 304–306, 305
 - indications and contraindications 302
 - lap squatting 306–308, 307
 - supported squat 183, 184, 196, 197, 302–304, 303
- stair climbing 200, 317–318, 318
- stations of descent 81–83, 81
 - assessment of 82–83, 83
- sterile water injections 364
 - injection sites 189, 253
 - intrapartum pain relief 252–253
 - procedure 253–254
- straining physiological effects
 - on fetus 172
 - on woman 172
- subcutaneous sterile water injections *see* sterile water injections

- suffering
 - psycho-emotional measures 24–31
 - before labor 24–26
 - during labor 26–27
 - trauma-informed care 27–31, **29–30**
 - versus pain 21–22
- supine hypotension 142, 145, 172, 201
- supine positions 200–201, 308–311, 308
 - effects of 308–309
 - indications and contraindications 309
 - prolonged active phase of first stage 142, 142
 - second stage dystocia 200–202, 201, 202
 - sheet “pull-to-push” 309–310, 309
 - see also* exaggerated lithotomy position
- supported squat positions 183, 184, 196, 197, 302–304, 303
 - effect of 304
 - indications and contraindications 304
- suprapubic pressure 248–249, 249
- synclitism 116
 - see also* asynclitism
- tend-and-befriend response 19–20, 19
- therapeutic rest 232–233
- third stage of labor 211–213, 216
 - active management 218–221
 - physiologic (expectant) management 217–218, 217
- three-part foot massage 341–343, 341, 342
- touch therapy 337
 - see also* massage
- transcutaneous nerve stimulation (TENS) 189, 362–364, 362
 - effectiveness 364
 - indications and contraindications 363
- trauma histories 27–28
- trauma-informed care (TIC) 28–31, **29–30**
- traumatic births 28–31
- UK maternity care practice 5
- ultrasound, fetal position assessment 129–130, 133–134
- umbilical cord
 - controlled cord traction (CCT) 218
 - delayed clamping and cutting 214–216
 - milking 216
 - tight nuchal cord management 216
- urinary retention 266
- urine assessment during labor 67
- US maternity care practice 5
- uterine bimanual compression 220, 220
- uterine lactic acidosis 149–150
- uterine massage 218–219
- uterotonics, prophylactic administration 219

388 *Index*

- vaginal examination 77–79, 87
 - descent assessment 81–83, 83
 - fetal position assessment 83–85, 84
 - indications for 77
 - performance of 77–79, 78
- vital signs during labor 69
- vomiting during labor 67

- walking 317–318, 318, 360
- weak contractions *see* inadequate contractions
- wireless telemetry 40–42, 41, 42

- zipper cervix 80
 - management 233–234