



---

## INDEX

---

### A

- Abdominal pain, emergency department visit with diagnosis of, 336–337, 339
- ACA. *See* Affordable Care Act
- Accidents: cerebrovascular, African Americans death rates with, 30; death, African Americans death rates with, 30; proportionate mortality ratio with, 81; road traffic, 271, 272, 273
- Accountable Care Organizations (ACOs), 200–201
- Accuracy, 152
- ACOs. *See* Accountable Care Organizations
- Acquired Immunodeficiency Syndrome (AIDS): azidothymidine for treatment of, 97; cases in U. S. of, 49, 57; by state, 58; DALYs caused by, 271, 272
- Act for Relief of Sick and Disabled Sailors, 5
- Active surveillance, 187
- Activities of daily living (ADL), 223
- Acute myocardial infarction (AMI): CMS incentives for improving quality care for, 243; mortality measures for, 224
- Acute upper respiratory infections, emergency department visit with diagnosis of, 336–337
- Adams, John (President), 5
- ADL. *See* Activities of daily living
- Administrative claims data, quality of care measurement with, 243
- Adnexa, ICD-10-CM codes for, 24
- Adult day care services, long-term care services utilization for, 309, 311
- Advance Payment ACO Model, 200–201
- Affordable Care Act (ACA), 10, 17; ACOs established by, 200; changes to OPPTS created by, 196; operating margin decrease from, 282; payment outliers

- cap made by, 198; preventive services emphasized by, 254; SNF prospective payment system in, 197
- Age differences: adult day care services utilization by, 309, 311; cancer deaths rates in U.S. with, 28; disease patterns correlated with, 26; doctor visits in U. S. by, 61; East Bank County, 304–306; East Bank Regional Hospital cancer incidence rates by, 297; East Bank Regional Hospital diabetes prevalence by, 298; East Bank Regional Hospital discharges by, 288, 290; East Bank Regional Hospital heart disease prevalence by, 297; East Bank Regional Hospital stroke prevalence by, 297; emergency department visits by, 26, 322–331; home health services utilization by, 309, 312, 316; Hospice utilization by, 309, 313, 316; hospital stay in East Bank Regional Hospital by, 285; insurance coverage trends with, 27, 294; Lakeshore County, 304, 306–307; length of hospital stay by, 291, 292; long-term care services utilization by, 309, 311–318; Medicaid coverage for Bank Regional Hospital by, 295; Medicare coverage for Bank Regional Hospital by, 294; nursing home utilization by, 309, 314, 316; office visits by, 345; outpatient department visits in U.S. by, 119; patient population by, 346; residential care community utilization by, 309, 314; service area population by, 283; tuberculosis cases by, 28
- Agency for Healthcare Research and Quality (AHRQ), 248
- Age-specific mortality rate, 75–76
- AHRQ. *See* Agency for Healthcare Research and Quality
- AIDS. *See* Acquired Immunodeficiency Syndrome
- Alcohol: liver dysfunction with, 106; lung cancer incidence with long-term use of, 120–121
- Allocation, 50; nonrandom, 96–97; random, 96
- Alternative Payment Model (APM), 257
- Alzheimer's disease, cause-specific mortality rate for, 74
- AMA. *See* American Medical Association
- Ambulatory patient classification (APCs), 9
- American College of Surgeons National Cancer Database, 187
- American Medical Association (AMA): standardized reporting attempted by, 193
- AMI. *See* Acute myocardial infarction
- Analytic scoring model, best practice assessment applied to, 239
- Analytic study designs, 90, 98, 115
- Anemia, emergency department visit with diagnosis of, 339
- Anthrax: cases in U. S. of, 49; as notifiable infectious disease, 48
- APCs. *See* Ambulatory patient classification
- APM. *See* Alternative Payment Model
- Arboviral diseases, as notifiable infectious disease, 48
- Area under the curve (AUC), 167
- Arteriography, East Bank Regional Hospital discharge following, 289, 290
- Arteriosclerotic heart disease, whites higher death rates with, 30

- Arthritis: gastrointestinal complications in patients with, 275; validity of screening tests for, 156
- Arthropathies, emergency department visit with diagnosis of, 336
- Asian and pacific islanders: cancer deaths rates in U.S. for, 31; crude death rate by, 74; frequency of new cancer cases in U.S. for, 31; hospital stay in East Bank Regional Hospital for, 285
- Assessment: best practices, 238–240; CAHPS, 230–231; HCAHPS, 221; health status, 225–227; IRF-PAI, 274; OASIS, 274; outcomes, 220–224; patient satisfaction, 227–231; PG-SGA, 212–213; quality of care functions in, 222
- Association: between risk factors, health, and disease, 103; alcohol use and liver dysfunction, 106; long-term alcohol use and lung cancer incidence, 120–121; measuring in prospective studies, 104–108; measuring in retrospective studies, 108–110; smoking and CHD, 106, 108, 110
- Asthma: emergency department visit with diagnosis of, 336; prevalence rate for, 69–70
- Attack rate, 178
- Attributable risk, 105–106
- AUC. *See* Area under the curve
- Average length-of-stay days, quality of care measurement with, 242
- Azidothymidine (AZT): AIDS treatment with, 97
- B**
- Babesiosis, as notifiable infectious disease, 48
- Balanced Budget Act of 1997, 9
- Baldrige Award, 249–250
- Behavioral disorders, ICD-10-CM codes for, 24
- Behavioral Risk Factor Surveillance Survey (BRFSS), 41–42; prevalence of women having mammogram, 70–71; prevalence of women having Papanicolaou test, 70–71
- Bell's Palsy, 3
- Benchmarking: best practices adoption from, 234; data as component for, 236; defined, 234; by external organizations, 237; future planning with, 235; process of, 235; stages of, 235–237; as team effort, 236
- Best practices: assessment for, 238–240; population health outcomes, 237–240
- Bias, 97
- Bills of Mortality* (Graunt), 22
- Bioterrorism, 4
- Birth certificates, 45–46, 77, 85
- Births, cesarean section rate, 110–112
- Blacks/African Americans: asthma prevalence rate for, 70; births among Medicaid beneficiaries for, 111; cancer deaths rates in U.S. for, 31; cardiovascular mortality among women of, 32; crude death rate by, 74; death rates with, 30, 33; doctor visits in U. S. by, 61; emergency department visits in U.S. for, 31, 62; end-stage renal disease risk for, 36; frequency of new cancer cases in U.S. for, 32; homicide rates for, 33; hospital stay in East Bank Regional Hospital for, 285; hypertension among, 32; infant mortality rate in U.S. for, 77; invasive carcinoma of cervix for woman of, 133–134; long-term care services

- utilization by, 309; neonatal mortality rate in U.S. for, 79; outpatient visits for, 132; post-neonatal mortality rate in U.S. for, 80; prevalence of women having mammogram for, 71
  - Blinding, clinical trials with, 97
  - Blood diseases: bloodborne pathogens, HAIs with, 185–186; ICD-10-CM codes for, 24
  - Bogalusa Heart Study, 4
  - Botulism: cases in U. S. of, 49; as notifiable infectious disease, 48
  - Breast cancer, retrospective studies on, 103
  - BRFSS. *See* Behavioral Risk Factor Surveillance Survey
  - Brucellosis: cases in U. S. of, 49; as notifiable infectious disease, 48
  - Burden of disease analysis, 269–272, 276; DALYs with, 270–272
- C**
- CABG. *See* Coronary artery bypass graft
  - CAHPS. *See* Consumer Assessment of Health Plan Survey
  - CAHPS® Database, 231
  - CAHPS Database Chartbook, 231
  - CAHPS Surgical Care Survey, 231
  - CAHs. *See* Critical care hospitals
  - Campylobacteriosis, as notifiable infectious disease, 48
  - Cancer: American College of Surgeons National Cancer Database, 187; breast, retrospective studies on, 103; cervix, 133–134; colorectal, validity of screening tests for, 157; deaths rates in U.S. by age for, 28; deaths rates in U.S. by race for, 31; East Bank Regional Hospital incidence rates for, 296, 297; frequency of new cases in U.S. by race for, 32; incidence locations for, 38; incidence rate in California for, 68; inpatient deaths in U.S. hospitals from, 42; lung: cost effectiveness analysis, 263; death caused by, 273; incidence with long-term alcohol use, 120–121; smoking with risk of, 120–121; as notifiable infectious disease, 48; socioeconomic status with prognosis for, 33–34
  - CAP. *See* Community-acquired pneumonia
  - Capitation, 194
  - Carbon monoxide poisoning, as notifiable infectious disease, 48
  - Cardiac catheterization, East Bank Regional Hospital discharge following, 289, 290
  - Carotid endarterectomy, East Bank Regional Hospital discharge following, 289, 290
  - Carve-outs, 169–170
  - Case-control study designs, 103
  - Case-fatality rate, 76–77; direct standardization example with, 126–129; myocardial infarction with, 126–129; quality of care measurement with, 242
  - Case-mix: analysis, 50; hospital's internal data as, 206
  - Case-mix groups (CMGs), 196, 198
  - Causation: exposure with, 93; risk with, 91–93; types of relationships with, 92
  - Cause and effect diagrams, quality control with, 245
  - Cause-specific mortality rate, 73–75
  - CDC. *See* Centers for Disease Control and Prevention

- CDC WONDER, 44
- Cellulitis, emergency department visit with diagnosis of, 336–337, 339
- Census Bureau, U.S., 123–124
- Centers for Disease Control and Prevention (CDC): death and disease data from, 44; periodic reports to, 44
- Centers for Medicare and Medicaid Services (CMS): coding system for inpatient procedures, 23; economic incentives for improving quality by, 243; Hospital Quality Initiative of, 221, 243; mortality measures for AMI from, 224; outcomes assessment for, 221, 231; Pneumonia National Project of, 224
- Cerebrovascular accidents, African Americans death rates with, 30
- Cerebrovascular disease: cause-specific mortality rate for, 74; emergency department visit with diagnosis of, 339
- Cesarean section rates: among Medicaid beneficiaries, 110–113; relative risk by age with, 112; relative risk by locale with, 112
- Chancroid: cases in U. S. of, 49; as notifiable infectious disease, 48
- CHD. *See* Coronary heart disease
- Checklists: AHRQ, 248; quality control with, 245, 247–248; WHO Checklist, 247
- Chest pain, emergency department visit with diagnosis of, 336–337, 339
- Chickenpox. *See* Varicella
- Child Health Insurance Program (CHIP), 9
- Children: cost-utility of intervention for obesity in, 256; health insurance program for, 9; health policy for, 9; health status indicated by growth in, 35
- CHIP. *See* Child Health Insurance Program
- Chlamydia Trachomatis*: cases in U. S. of, 49; incidence rate in U. S. for, 68; as notifiable infectious disease, 48
- Cholera, 4; cases in U. S. of, 49; as notifiable infectious disease, 48
- Chromosomal abnormalities, ICD-10-CM codes for, 24
- Chronic bronchitis, emergency department visit with diagnosis of, 336, 339
- Chronic disease management, 169–170
- Chronic liver disease, potential years of life lost with, 83
- Chronic Liver Disease Questionnaire (CLDQ): construct validity of, 211
- Chronic lower respiratory diseases: cause-specific mortality rate for, 74; potential years of life lost with, 83
- Chronic obstructive pulmonary disease, DALYs caused by, 271, 272
- Circulatory system diseases: ICD-10-CM codes for, 24; proportionate mortality ratio with, 81
- Cirrhosis, potential years of life lost with, 83
- Claims data, 223
- CLDQ. *See* Chronic Liver Disease Questionnaire
- Clinical effectiveness, 152
- Clinical Risk Groups (CRGs), 273
- Clinical trials: bias avoidance in, 97; blinding for, 97; controlled, 95; defined, 95; evidence for testing study

- hypotheses with, 94; framework of randomized controlled, 96; need for, 94–95; nonrandom allocation with, 96–97; Phase I, 95; Phase II, 95; Phase III, 95; random allocation with, 96; types of, 94, 95; uncontrolled, 95
- Clostridium difficile*, HAI caused by, 183
- Cluster sampling, 114, 204, 205
- CMGs. *See* Case-mix groups
- CMS. *See* Centers for Medicare and Medicaid Services
- Coccidioidomycosis: cases in U. S. of, 49; as notifiable infectious disease, 48
- Cohort studies. *See* Prospective studies
- Colorectal cancer, validity of screening tests for, 157
- Commission on Professional and Hospital Activities (CPHA), 46
- Common-source epidemics, 178
- Communities, managing health care for, 12–14
- Community-acquired pneumonia (CAP): CMS incentives for improving quality care for, 243
- Component cause, 92
- Confounding, 118
- Confounding factor, 118
- Confounding variable, 118
- Congenital malformations, ICD-10-CM codes for, 24
- Congestive heart failure, patients readmissions incidence rate for, 67
- Connective tissue diseases, ICD-10-CM codes for, 24
- Construct validity, 211, 213
- Consumer Assessment of Health Plan Survey (CAHPS), 230–231
- Consumer Satisfaction Survey, 229
- Content validity, 211
- Control charts: LTBI monitoring with, 245–246; patient falls in, 246; quality control with, 245–247
- Contusion, emergency department visit with diagnosis of, 336–337
- Convergent validity, 211
- COOP charts, 226
- Coronary artery bypass graft (CABG): CMS incentives for improving quality care for, 243
- Coronary heart disease (CHD): smoking with, 106, 108, 110
- Correlational analysis, 213
- Cost analysis, 259–260, 275
- Cost-benefit analysis, 261–262, 275
- Cost-effectiveness analysis, 275; cost-benefit distinction from, 261; diagnostic interventions illustration of, 262–263; lung cancer screening illustration of, 263
- Cost-effectiveness ratio, 268, 274, 279
- Cost-utility analysis, 275; disability-adjusted life years used by, 264; QALYs used by, 264; treatment approaches in, 265
- Counts, 57–58; doctor visits in U. S., 61; emergency department visits in U. S., 62
- CPHA. *See* Commission on Professional and Hospital Activities
- CPT. *See* Current Procedural Terminology
- CRGs. *See* Clinical Risk Groups
- Criterion validity, 212–213
- Critical care hospitals (CAHs), 196
- Cronbach's alpha, 214
- Crossing the Quality Chasm: A New Health System for the 21st Century* (Bloom, 2002), 201, 222

- Cross-sectional studies designs: examples of, 100–101; problem associated with, 100; purpose of, 100; risk factor exposure in, 99–100
- Crude birth rate, 85
- Crude death rate, 73–74
- Cryptosporidiosis: cases in U. S. of, 49; as notifiable infectious disease, 48
- CT scan, East Bank Regional Hospital discharge following, 289, 290
- Cumulative incidence, 65
- Current Procedural Terminology (CPT), 193; RBRVS with, 198–199
- Cyclosporiasis: cases in U. S. of, 49; as notifiable infectious disease, 48
- D**
- DALYs. *See* Disability-adjusted life years
- Data analysis: analytical analysis for, 206–207; case-mix in, 206; descriptive analysis for, 206; epidemiologic data in, 207; market research, 205–207; types of data in, 205–206
- Death rate: African American, 30; age-specific mortality rate, 75–76; cancer in U.S. by age, 28; cancer in U.S. by race, 31; case-fatality rate, 76–77; cause-specific mortality rate, 73–75; crude, 73–74; fetal death rate, 78; infant mortality rates, 77–78; neonatal mortality rates, 78, 79; post-neonatal mortality rates, 78, 80; proportionate mortality ratio, 79–81; race and ethnicity based, 32–33; sex based, 29
- Demand estimation: econometric models for, 210; population-based models for, 210; source of data for, 209; traditional utilization projections for, 210
- Demography, founder of, 22
- Dengue viral infections, as notifiable infectious disease, 48
- Department of Health and Human Services, U.S., periodic reports to, 44
- Dependent variable, 90; independent variable relationship with, 91
- Descriptive data: BRFSS for, 41–42; health essential information from, 19–53; important to health care administrators, 20; medical records for, 46; payment source as, 42–43; person variables in: age, 26–29; marital status, 33, 34; race or ethnicity, 30–33; sex, 29–30; socioeconomic status, 33–36; place variables for, 36–39; population pyramids for, 36, 37; reportable diseases in, 47–51; sources of, 43–46; time as variable for, 39–42; two forms of, 25; vital statistics for, 45–46; YRBSS for, 42
- Descriptive studies designs, 99
- Diabetes: cause-specific mortality rate for, 74; East Bank Regional Hospital prevalence of, 298, 299; Medicare coverage for, 254; mellitus, emergency department visit with diagnosis of, 339; mental health status among, 85; potential years of life lost with, 83; prevalence ratios for, 64
- Diagnosis-related groups (DRGs), 9, 10, 194–195
- Diagnostic test, impact of, 162–168
- Diagnostic ultrasound, East Bank Regional Hospital discharge following, 289, 290
- Diarrheal diseases: DALYs caused by, 271, 272; death caused by, 273

- Digestive system diseases, ICD-10-CM codes for, 24
- Diphtheria: cases in U. S. of, 49; as notifiable infectious disease, 48
- Direct causal relationships, 92
- Direct method of standardization: case-fatality rate example for, 126–129; outpatient department visits by sex using, 126; perspectives to understand, 124–125; steps of, 123; U.S. population from Census Bureau as example of, 123–124
- Direct rating, 264–265
- Direct transmission, 177
- Disability-adjusted life years (DALYs), 186; cost-utility analysis using, 264; disease burden is expressed in, 270–272; leading causes of, 271, 272
- Discharge status, quality of care measurement with, 242
- Discharges, 287–289
- Discounting, 265–269; community outreach program benefits with, 268; formula for, 266; in-hospital program benefits with, 268; present value of costs for, 267
- Discriminant validity, 211
- Disease: definitions associated with, 21–22; ICD-10-CM codes for, 24; international classification of, 22–24; measurement as indication of health status, 21; prevention, medical management with, 150–151; screening, 152
- DRGs. *See* Diagnosis-related groups
- E**
- Ear diseases, ICD-10-CM codes for, 24
- East Bank County, 308, 309–319; population characteristics of, 304–306
- East Bank Regional Hospital: age of hospital stay in, 285; background information on, 282, 342; board recommendations for reconfiguration and expansion of, 282; cancer incidence rates for, 296, 297; case-fatality rates for MI in, 126–129; chronic conditions for, 356–358; cost analysis illustrated in, 260; cost effectiveness analysis illustrated in, 263; description of, 292–293; diabetes prevalence for, 298, 299; discharges for, 287–290; by age, 90, 288; per procedure, 289, 290; by sex, 288, 289; heart disease prevalence for, 297; hospital care utilization for, 284–287; one hospital stay per year percent for, 286; percent occupancy rate by hospital in, 292, 293; two or more hospital stay per year percent for, 286, 287; insurance coverage in, 294, 349–351; length of hospital stay for, 289–292; Medicaid coverage in, 295; Medicare coverage in, 295; payment sources for, 293–295; physician practices for, 341–358; population characteristics for, 342–343; preventive care visits for, 355–356; race of hospital stay in, 285; reason for visit for, 351–354; resource utilization for, 343–346; risk adjustment in, 137–139; service area population for, 283–284; service area population health for, 295–301; sex of hospital stay in, 285; stroke prevalence for, 297; time since last visit for, 346–348
- Ecological studies designs, 101–102
- Econometric models, 210



- Economic analysis, 253–279; burden of disease analysis, 269–272, 276; cost analysis, 259–260, 275; cost-benefit analysis, 261–262, 275; cost-effectiveness analysis, 262–263, 275; cost-utility analysis, 264–265, 275; discounting, 265–269; economic evaluation, 257–259; examples of, 272–275; sensitivity analysis, 269
- Economic evaluation: four steps in, 258–259; importance to health care managers of, 258; outcomes of, 258; questions underlying, 257–258
- Educational status: age-adjusted diabetes prevalence ratios by, 64; East Bank Regional Hospital cancer incidence rates by, 297; East Bank Regional Hospital heart disease prevalence by, 297; East Bank Regional Hospital stroke prevalence by, 297; health insurance coverage in U.S. for, 35–36
- Efficacy, 152
- Efficiency, 152
- Ehrlichiosis: cases in U. S. of, 49; as notifiable infectious disease, 48
- Emergency department services, 321–340; age of patient visiting for, 322–331; patient disposition for, 337–340; payment sources for, 331–332; primary diagnosis at arrival to, 336–337; race of patient visiting for, 326–327, 330–331; sex of patient visiting for, 324–325, 328, 329; time analysis for, 332–335; utilization of, 322–331; wait time spent in, 333–334
- Employment status, mental disorders trends for, 35
- Encephalitis/meningitis, cases in U. S. of, 49
- Endocrine diseases, ICD-10-CM codes for, 24
- Endoscopy: large intestine, East Bank Regional Hospital discharge following, 289, 290; small intestine, 289, 290
- Enteritis, emergency department visit with diagnosis of, 339
- Enterohemorrhagic *Escherichia coli*, cases in U. S. of, 49
- EPH. *See* Expected probability of mortality
- Epidemic curve, 39
- Epidemics, 178; time period with, 39. *See also* Infectious epidemics
- Epidemiology: data for, 15; defined, 2, 16; experimental, 7; focus of, 6–8; history and introduction to, 2–5, 16; infectious disease, 176–182; infectious epidemics in, 178–182; managerial, 5–6, 15; observational, 6–7; philosophical framework for, 5–6; population-based perspective of, 5, 12; primary prevention in, 7–8; role of, 14–15; social, 12; subpopulations in, 11–12; 21st century, 4; uses of, 6–8
- EQ-5D, 226
- Error, 152
- Escherichia coli*: enterohemorrhagic, cases in U. S. of, 49; HAI caused by, 182
- Ethnicity. *See* Race and ethnicity
- Evidence-based practices, 237–238
- Expanding hospital care services, 281–301; discharges with, 287–290; hospital care utilization with, 284–287; hospital description with, 292–293; length of hospital stay with, 289–292; payment sources with, 293–295; service

area population health with, 295–301;  
 service area population with, 283–284  
 Expected probability of mortality (EPH),  
 167–168  
 Experimental epidemiology, 7  
 Experimental study designs: array of, 94;  
 framework of, 93–94  
 Exposure, 93  
 Eye diseases, ICD-10-CM codes for, 24

### F

Face validity, 210  
 False negatives (FN), 153  
 False positives (FP), 153  
 Farr, William, 3  
 Fee-for-service (FFS), 220  
 Fertility rate, 85–86  
 Fetal death rate, 78  
 FFS. *See* Fee-for-service  
 Flow charts, quality control with, 245  
 FN. *See* False negatives  
 Food-borne infection outbreak, as  
 notifiable infectious disease, 48  
 Forecasting, demand estimation, 209–210  
 FP. *See* False positives  
 Fracastorius, Girolamo, 3  
 Fractures, emergency department visit  
 with diagnosis of, 336–337, 339  
 Framingham Heart Study, 4  
 Frequency, disease patterns, 30  
 Functional Status Questionnaire, 226

### G

Gallbladder removal, East Bank Regional  
 Hospital discharge following, 289, 290  
 Gastrointestinal hemorrhage, emergency  
 department visit with diagnosis of, 339

GBD. *See* Global Burden of Disease  
 Study  
 General population surveillance, 187  
 Genitourinary system diseases: ICD-10-CM  
 codes for, 24; proportionate mortality  
 ratio with, 81  
 Geographic Practice Cost Index  
 (GPCI), 199  
 Germ theory, 3  
 GHAA. *See* Group Health Association of  
 America  
 Giardiasis, as notifiable infectious  
 disease, 48  
 Global Burden of Disease Study (GBD),  
 186, 270  
 Gonorrhea: cases in U. S. of, 49;  
 incidence rate in U. S. for, 68; as  
 notifiable infectious disease, 48  
 Gossett, William, 107  
 GPCI. *See* Geographic Practice Cost Index  
 Graunt, John, 22  
 Group Health Association of America  
 (GHAA), 229

### H

HAIs. *See* Health care–associated  
 infections  
 Hansen disease (Leprosy): cases in U. S.  
 of, 49; as notifiable infectious  
 disease, 48  
 Hantavirus pulmonary syndrome: cases in  
 U. S. of, 49; as notifiable infectious  
 disease, 48  
 Harvard University School of Public  
 Health, burden of disease study  
 by, 270  
 Hawaiian: doctor visits in U. S. by, 61; HIV  
 cases, 60

- HCAHPS. *See* Hospital Consumer Assessment of Healthcare Providers System
- Headache, emergency department visit with diagnosis of, 336
- Health: definitions associated with, 21–22; description of, 19–53
- Health care: community-based nature of, 13; cost analysis for, 259–260, 275; cost-benefit analysis for, 261–262, 275; cost-effectiveness analysis for, 262–263, 275; cost-utility analysis for, 264–265, 275; economic analysis for population, 253–279; economic evaluation, importance to, 258; managing for populations and communities, 12–14; policy for, 254–255; reform, 8–11; system, marketing of, 203
- Health Care and Education Reconciliation Act of 2010, 10
- Health care planning, 5, 8, 14, 16, 21, 41, 56, 65, 100, 118, 192, 214, 254
- Health care–associated infections (HAIs), 176, 188; bloodborne pathogens are responsible for, 185–186; calculating infection rates for, 184–185; causes of, 182–183; high costs associated with outbreaks of, 183; occurrence of, 182
- Health insurance: children, program for, 9; coverage by educational status in U.S. for, 35–36; coverage by socioeconomic status in U.S. for, 35–36; coverage trends by age for, 27; East Bank Regional Hospital coverage with, 294, 349–351; emergency department payments from, 331–332; payment source with, 43; physician practices, coverage with, 349–351
- Health Maintenance Organization Act, 8, 194
- Health maintenance organizations (HMOs), 8–9; capitation with, 194; carve-outs with, 169–170; outcomes to evaluate potential providers for, 233
- Health measurements, 55–87; counts in, 57–58; health-related quality of life in, 84–85; morbidity in, 65–72; mortality in, 73–83; potential years of life lost in, 80–83; proportions in, 63–65; quantification of disease in population as basic to, 56; rates in, 58–62; crude birth rate, 85; death rate, 28–33, 73–80; fertility rate, 85–86; incidence rate, 65–68; period prevalence rate, 72; prevalence rate, 69–72; survival rate, 83; ratios in, 62–64
- Health risk behaviors, 42. *See also* Risk factors
- Health status: assessment for, 225–227; descriptive disease measurement indicating, 21; dimensions of, 24; factors in determination of, 225; growth in children with socioeconomic status for, 35; measurement of, 24–25; mental among diabetics, 85; quality of life with, 25
- Health-related quality of life (HRQL), 84–85; patient satisfaction with, 208
- Heart disease: arteriosclerotic, whites higher death rates with, 30; cause-specific mortality rate for, 74; coronary, smoking with, 106, 108, 110; East Bank Regional Hospital prevalence of, 297; emergency department visit with diagnosis of, 339; hypertensive, African Americans death

- rates with, 30; inpatient deaths in U.S. hospitals from, 42; ischemic, DALYs caused by, 271, 272; ischemic, death caused by, 273; potential years of life lost with, 83
- Heart failure (HF): CMS incentives for improving quality care for, 243
- Heart surgery, East Bank Regional Hospital discharge following, 289, 290
- HEDIS® Effectiveness of Care measures, 232
- Hemolytic uremic syndrome: cases in U. S. of, 49; as notifiable infectious disease, 48
- Hemophilus Influenza*: cases in U. S. of, 49; as notifiable infectious disease, 48
- Hepatitis, as notifiable infectious disease, 48
- Hepatitis A: cases in U. S. of, 49; incidence rate in U. S. for, 68
- Hepatitis B: cases in U. S. of, 49; HAI with, 185–186; incidence rate in U. S. for, 68
- Hepatitis C: cases in U. S. of, 49; HAI caused by, 182–183
- HF. *See* Heart failure
- HH-PPS. *See* Home Health Prospective Payment System
- HHRGs. *See* Home Health Resource Groups
- Hip replacement: CMS incentives for improving quality care for, 243; East Bank Regional Hospital discharge following, 289, 290
- Hippocrates, 3
- Hispanic and Latinos: crude death rate by, 74; doctor visits in U. S. by, 61; emergency department visits in U.S. for, 31, 62; hospital stay in East Bank Regional Hospital for, 285; long-term care services utilization by, 309; prevalence of women having mammogram for, 71
- Histograms, quality control with, 245
- HIV. *See* Human immunodeficiency virus
- HMOs. *See* Health maintenance organizations
- Home Health Prospective Payment System (HH-PPS), 197–198
- Home Health Resource Groups (HHRGs), 198
- Home health services, long-term care services utilization for, 309, 312, 316
- Homicide, African Americans death rates with, 30
- HOS. *See* Medicare Health Outcomes Survey
- Hospice, long-term care services utilization for, 309, 313, 316
- Hospital care utilization, 284–287; by age, 285; length of hospital stay with, 289–292; percent of days hospital stay per year in, 286, 287; by race, 285; by sex, 285
- Hospital Compare, 221–222
- Hospital Consumer Assessment of Healthcare Providers System (HCAHPS), 221
- Hospital Quality Initiative, 221
- Hospital service area, 13, 216, 295, 322
- Hospital standardized mortality ratio (HSMR), 135–136
- Hospital value-based purchasing (HVBP), 233
- Hospitals: East Bank Regional Hospital, 90, 126–129, 137–139, 260, 263,

- 282–301, 341–358; Riverview Hospital, 321–340; Riverview Medical Center Hospital, 137–139; Shoreline Regional Hospital, 292, 293; Westbank Hospital, 292, 293; Westside Memorial Hospital, 126–129, 292–293
- HRQL. *See* Health-related quality of life
- HSMR. *See* Hospital standardized mortality ratio
- Human immunodeficiency virus (HIV):  
cause-specific mortality rate for infection with, 74; DALYs caused by, 271, 272; ethnicity ratios of case in U. S. with, 63; HAIs with, 185–186; myocardial infection with, 113; as notifiable infectious disease, 48; rates of cases in U. S. of, 60
- HVBP. *See* Hospital value-based purchasing
- Hypertension, African Americans with, 32
- Hypertensive heart disease, African Americans death rates with, 30
- I**
- ICD. *See* International Classification of Diseases
- ICD-9-CM, 23
- ICD-10-CM, 23–24; disease category codes, 24
- ICD-10-PCS, 23
- Immunogenicity, 177
- Incidence density, 65–66
- Incidence rate: calculation example of, 66–67; cancer in California, 68; congestive heart failure patients readmissions, 67; cumulative incidence and incidence density with, 65–66; defined, 65; denominator of, 65; incident cases with, 66; infectious diseases in U. S., 68; numerator of, 65; prevalence rate related to, 70, 72
- Incident cases, 66, 102
- Independent variable, dependent variable relationship with, 91
- Indirect causal relationships, 92
- Indirect method of standardization:  
HSMR with, 135–136; invasive carcinoma of cervix example using, 133–134; outpatient visits example using, 131–133; process of, 134; SMR with, 130, 132–135, 140, 148; steps in, 129–130
- Indirect transmission, 177
- Infant mortality rate, 77–78
- Infection control procedures, 15
- Infection rates, HAIs with, 184–185
- Infectious diseases, ICD-10-CM codes for, 24
- Infectious epidemics: attack rate for, 178; defined, 178; effect of, 178; point-source exposure with, 178–179; propagated epidemics with, 179; SARS, 180–181; West Nile virus, 181; Zika virus, 181–182
- Infectious epidemics management, 175–190; epidemiology with, 176–182; HAIs with, 176, 182–186, 188; surveillance for, 186–188
- Infectivity, 177
- Influenza, 4; cause-specific mortality rate for, 74; *Hemophilus Influenza*, 48, 49; potential years of life lost with, 83
- Influenza-associated pediatric mortality, as notifiable infectious disease, 48
- Inpatient rehabilitation entities (IREs), 196

Inpatient Rehabilitation Facility  
 Prospective Payment System (IRF-PPS),  
 196–197

Inpatient Rehabilitation Facility-Patient  
 Assessment Instrument (IRF-PAI), 274

Institute of Medicine (IOM), 222

Integrated health care system, 203

Internal consistency, 213–214

International Classification of Diseases  
 (ICD): basis for, 23; clinical modifica-  
 tion 9, 23; clinical modification 10,  
 23–24; disease data collection by, 22;  
 international comparability with,  
 22; origin of, 22; and related  
 conditions, 3

Invasive pneumococcal diseases, as  
 notifiable infectious disease, 48

IOM. *See* Institute of Medicine

IREs. *See* Inpatient rehabilitation entities

IRF-PAI. *See* Inpatient Rehabilitation  
 Facility-Patient Assessment Instrument

IRF-PPS. *See* Inpatient Rehabilitation  
 Facility Prospective Payment System

Ischemic heart disease: DALYs caused by,  
 271, 272; death caused by, 273

## J

Jefferson county: crude rate for MI  
 complications in, 145; expected MI  
 complications for, 147; population  
 distribution for, 146

Joint Commission, 240–241

## K

Kappa statistic, 162

Kidney disease, inpatient deaths in U.S.  
 hospitals from, 42

Knee replacement: CMS incentives for  
 improving quality care for, 243; East  
 Bank Regional Hospital discharge  
 following, 289, 290

Knowledge About Older Patients-Quiz  
 (KOP-Q), 214

## L

Lacroix, Francois Bossier de, 22

Lakeshore County, 309–319; population  
 characteristics of, 304, 306–308

Latent tuberculosis infection (LTBI):  
 control charts as tool to monitor  
 improvement of, 245–246

Lead, elevated blood levels, as notifiable  
 infectious disease, 48

Leading causes of DALYs, 271, 272

Legionellosis: cases in U. S. of, 49; as  
 notifiable infectious disease, 48

Legionnaire's disease, 179

Length of hospital stay, 289–292

Leprosy. *See* Hansen disease

Leptospirosis, as notifiable infectious  
 disease, 48

Leukemia: deaths rates in U.S. by race for,  
 31; frequency of new cases in U.S. by  
 race for, 32; whites higher death rates  
 with, 30

Lifetime prevalence rate, 69

Likelihood ratios, 170; dichotomous  
 results in tests with, 163; parallel testing  
 with, 166; polychotomous results in  
 tests with, 164–165; pretest odds  
 with, 165

Lind, James, 3

Listeriosis: cases in U. S. of, 49; as  
 notifiable infectious disease, 48

- Liver disease, chronic: CLDQ, 211;  
potential years of life lost with, 83
- Liver dysfunction, alcohol use  
with, 106
- Long-term care: population  
characteristics of, 304–308; spending  
in U.S. for, 303; staffing for, 319;  
utilization of services for, 308–318;  
Westside Elderly Care, Inc. example  
of, 308
- Lower respiratory infections: DALYs  
caused by, 271, 272; death caused  
by, 273
- LTBI. *See* Latent tuberculosis infection
- Lung cancer: cost effectiveness analysis  
illustrated in screening, 263; death  
caused by, 273; incidence with  
long-term alcohol use, 120–121;  
smoking with risk of, 120–121
- Lyme disease: cases in U. S. of, 49;  
incidence rate in U. S. for, 68; as  
notifiable infectious disease, 48
- Lymphomas: deaths rates in U.S. by race  
for, 31; frequency of new cases in U.S.  
by race for, 32
- M**
- Magnetic resonance imaging, East Bank  
Regional Hospital discharge following,  
289, 290
- Malaria: cases in U. S. of, 49; DALYs  
caused by, 271, 272; as notifiable  
infectious disease, 48
- Malcolm Baldrige National Quality  
Program, 249–250
- Mammogram, prevalence of women  
having, 70–71
- Managed care contracting, 169, 233
- Managerial epidemiology: definition of,  
5; evolutionary pressures on, 6;  
external forces leading to, 5–6;  
relationship of data to aspects of, 15
- Managing chronic diseases, 169–170
- Managing health outcomes:  
benchmarking for, 234–237; benefits  
of, 232; best practices for, 237–240;  
practice guidelines for, 237–238; root  
cause analysis for, 240–241; uses of  
outcomes assessment in, 232–234
- Marginal cost-effectiveness ratio, 274, 279
- Marine Hospital Service, 5
- Marital status: chronic disease risk factors  
associated with, 34; morbidity and  
mortality associated with, 33
- Market research: data analysis for,  
205–207; demand estimation with,  
209–210; patient satisfaction with,  
207–209; reliability of, 210–214;  
sampling for, 204–205; validity of,  
210–214
- Marketing, in context of health care:  
defined, 202; health care system in,  
203; role of, 203
- Masking, 97
- Mastoid process diseases, circulatory  
system diseases, ICD-10-CM codes  
for, 24
- Matching: caveats with, 122; classifications  
for, 121–122
- MDS. *See* Minimum Data Sets
- Measles. *See* Rubella
- Medicaid: cesarean section rates among  
beneficiaries of, 110–113; East Bank  
Regional Hospital payment from, 295;  
emergency department payments  
from, 331–332; long-term care

- services utilization by, 309; Oregon, 255–256
- Medicaid/SCHIP, payment source with, 43
- Medical management, 149–173; chronic disease in, 169–170; clinical effectiveness as part of, 152; likelihood ratios with, 163–166, 170; practice analysis with, 168–169; prevention of disease as component of, 150–151; reliability with, 160–162; test information impact on, 162–168; validity with, 152–160
- Medical Outcomes Study (MOS), 225
- Medical records, 46; outcomes assessment with, 223; quality of care measurement with, 243
- Medicare: capitation rates adjustment for, 272–275; diabetes coverage, 254; East Bank Regional Hospital payment from, 295; emergency department payments from, 331–332; managed care, 232; outpatient prescription medication with, 10; Part D coverage, 10, 254; payment source with, 43; reimbursements, 282; resource-based relative value scale with, 9
- Medicare Advantage: hospital admissions with, 257; mortality with, 257
- Medicare Health Outcomes Survey (HOS), 232
- Medicare Prescription Drug, Improvement, and Modernization Act of 2003, 10
- Medicare severity diagnosis-related groups (MS-DRGs), 10, 195
- Medicare Shared Savings Program, 200
- Meningococcal disease: cases in U. S. of, 49; incidence rate in U. S. for, 68; as notifiable infectious disease, 48
- Mental disorders, ICD-10-CM codes for, 24
- Merida's Department of Human Resources, 227
- Merit-Based Incentive Payment System (MIPS), 257
- Mesothelioma: deaths rates in U.S. by race for, 31; frequency of new cases in U.S. by race for, 32
- Metabolic diseases, ICD-10-CM codes for, 24
- MI. *See* Myocardial infarction
- Minimum Data Sets (MDS), 274
- MIPS. *See* Merit-Based Incentive Payment System
- Misoprostol prophylaxis cost effectiveness, 274–275
- MMWR. *See* Mortality and Morbidity Weekly Report
- Modified determinism, theory of, 92
- Monitoring health outcomes, 231–232
- Morbidity: incidence rate for, 65–68; marital status associated with, 33; measurement of, 65–72; MMWR, 44; period prevalence rate for, 72; prevalence rate for, 69–72; rates based on sex, 29; SMR with, 130, 132–135, 140, 148
- Mortality: AMI and measures of, 224; *Bills of Mortality*, 22; cardiovascular African Americans women, 32; death rate for, 73–80; expected probability of, 167–168; HSMR, 135–136; influenza-associated pediatric, 48; marital status associated with, 33; measurement of, 73–83; Medicare



- Advantage with, 257; MMWR, 44;  
 potential years of life lost for, 80–83;  
 SMR with, 130, 132–135, 140, 148;  
 standardized rate of, 124
- Mortality and Morbidity Weekly Report  
 (MMWR), 44
- Mortality Prediction Model III, 167
- Mortality rate: age-specific, 75–76;  
 cause-specific, 73–75; infant, 77–78;  
 neonatal, 78, 79; quality of care  
 measurement with, 242. *See also*  
 Death rate
- Mortality ratio, proportionate, 79–81
- MOS. *See* Medical Outcomes Study
- MOS short-form 36 (SF-36), 225
- MS-DRGs. *See* Medicare severity  
 diagnosis-related groups
- Multistage sampling, 114, 204, 205
- Mumps: cases in U. S. of, 49; as notifiable  
 infectious disease, 48
- Musculoskeletal system diseases:  
 ICD-10-CM codes for, 24;  
 proportionate mortality ratio with, 81
- Myelomas: deaths rates in U.S. by race for,  
 31; frequency of new cases in U.S. by  
 race for, 32
- Myocardial infarction (MI): case-fatality  
 rate with, 126–129; crude rate for  
 complications, 145; expected complica-  
 tions for, 147; HIV infected with, 113.  
*See also* Acute myocardial infarction
- N**
- National Ambulatory Medical Care  
 Survey, 43
- National Center for Health Statistics  
 (NCHS), 44; benchmark infection rate  
 data from, 184
- National Diabetes Collaborative Registry  
 (NDCR), 187–188
- National Health and Nutritional  
 Examination Survey (NHANES), 44
- National Health Interview Survey (NHIS),  
 27, 34, 44
- National Hospital Ambulatory Medical  
 Care Survey (NHAMCS), 118–119;  
 data, 26, 61; Outpatient Department  
 Summary, 131
- National Hospital Discharge Survey, 37,  
 38, 44, 46
- National Quarantine Act of 1878, 5
- NCHS. *See* National Center for Health  
 Statistics
- NDCR. *See* National Diabetes  
 Collaborative Registry
- Necessary, but not sufficient  
 relationship, 92
- Necessary and sufficient relationship, 92
- Neither sufficient, nor necessary  
 relationship, 93
- Neonatal intensive care unit (NICU):  
 clinical process improvement  
 intervention for, 244
- Neonatal mortality rate, 78, 79
- Neoplasms: age-specific mortality rate for,  
 75; cause-specific mortality rate for, 74;  
 emergency department visit with  
 diagnosis of, 339; ICD-10-CM codes  
 for, 24; potential years of life lost  
 with, 83
- Nephritis, cause-specific mortality rate  
 for, 74
- Nervous system diseases, ICD-10-CM  
 codes for, 24
- Neurodevelopmental disorders,  
 ICD-10-CM codes for, 24

New York State Cardiac Advisory Committee, 237  
 NHAMCS. *See* National Hospital Ambulatory Medical Care Survey  
 NHANES. *See* National Health and Nutritional Examination Survey  
 NHIS. *See* National Health Interview Survey  
 NICU. *See* Neonatal intensive care unit  
 Norovirus, cost-utility of intervention for, 256  
 Notifiable diseases, 47–51  
 Nottingham Health Profile, 226  
 Nursing home, long-term care services utilization for, 309, 314, 316  
 Nutritional diseases, ICD-10-CM codes for, 24

## O

OASIS. *See* Outcome and Assessment Information Set  
 Obesity, cost-utility of childhood intervention for, 256  
 Observational epidemiology, 6–7  
 Observational study designs, 98–99  
 Occupation, risk factor exposures with, 29  
 Odds ratio, 109; pretest odds and likelihood ratios with, 165; smoking and CHD in analysis of, 110  
 OHP. *See* Oregon Health Plan  
 OHP-Plus, 255  
 OHP-Standard, 255  
 Older People in Acute Care Survey-United States (OPACS-US), 214  
 Open wound, emergency department visit with diagnosis of, 336–337  
 OPPS. *See* Outpatient Prospective Payment System

OPS surveys. *See* Outpatient Satisfaction surveys  
 Oregon Health Plan (OHP), 255  
 Oregon Medicaid proposal, 255  
 Otitis media, emergency department visit with diagnosis of, 336  
 Outbreaks, 178. *See also* Infectious epidemics  
 Outcome and Assessment Information Set (OASIS), 274  
 Outcome variable. *See* Dependent variable  
 Outcomes assessment, 220–224; activities of daily living in, 223; claims data in, 223; economic evaluation for, 258; medical records in, 223; monitoring, 231–232; objectives of, 221; process measures for, 222–223; quality of care comparison with, 221–222; uses of, 232–234  
 Outpatient department visits, 119, 126, 131, 142–143  
 Outpatient Prospective Payment System (OPPS), 9, 195–196  
 Outpatient Satisfaction (OPS) surveys, 229

## P

PACE. *See* Program for All-inclusive Care for the Elderly  
 Pacific islanders. *See* Asian and pacific islanders  
 PAN26, 214  
 Papanicolaou test, prevalence of women having, 70–71  
 Parallel testing, 166  
 Parasitic diseases, ICD-10-CM codes for, 24  
 Pareto charts, quality control with, 245, 248

- Partial hip replacement, East Bank  
 Regional Hospital discharge following, 289, 290
- Pathogenicity, 176–177
- Patient falls, control chart for, 246
- Patient Protection and Affordable Care Act, 10
- Patient satisfaction, 207–209; assessment for, 227–231; control chart, 230; quality of care measurement with, 242
- Patient Satisfaction Questionnaire (PSQ), 228
- Patient-Generated Global Assessment (PG-SGA), 212–213
- Pay-for-performance, 201, 233
- Payment per member per month (PMPM), 194
- Payment sources, 42–43, 293–295
- Pediatric population, 183
- Performance improvement, 250
- Period prevalence rate, 72
- Person variables, 20, 26
- Personal protection equipment, 15
- Pertussis: cases in U. S. of, 49; incidence rate in U. S. for, 68; as notifiable infectious disease, 48
- Pesticide-related illness and injury, as notifiable infectious disease, 48
- PG-SGA. *See* Patient-Generated Global Assessment
- Phase I clinical trials, 95
- Phase II clinical trials, 95
- Phase III clinical trials, 95
- Phthisis, Hippocrates' descriptions of, 3
- Physician practices, 341–358; chronic conditions with, 356–358; insurance coverage with, 349–351; population characteristics with, 342–343; preventive care visits with, 355–356; reason for visit with, 351–354; resource utilization with, 343–346; time since last visit with, 346–348
- Pioneer ACO Model, 200–201
- Place diseases, 37
- Place variables: cancer incidence locations, 38; etiologic factors of disease understood by, 36; important for health impact understanding, 39; political boundaries as, 38; short-stay U.S. hospital discharges with, 37–38; socioeconomic status with, 38; U.S. regional, 37–38
- Plague: cases in U. S. of, 49; as notifiable infectious disease, 48
- Planning. *See* Health care planning
- PMPM. *See* Payment per member per month
- PMR. *See* Proportionate mortality ratio
- Pneumonia, 4; community-acquired, 243; emergency department visit with diagnosis of, 339; inpatient deaths in U.S. hospitals from, 42
- Pneumonia National Project, 224
- Point prevalence rate, 69
- Point-source exposure, 178–179
- Poisoning: carbon monoxide, 48; ICD-10-CM codes for, 24
- Policy, 254
- Policy analysis, 254
- Poliomyelitis, as notifiable infectious disease, 48
- Poliovirus infection, as notifiable infectious disease, 48
- Population distribution: Jefferson county, 146; percent of, 284



- Population health outcomes, 219–251;  
 assessing, 220–224; uses of, 232–234;  
 benchmarking for, 234–237; best  
 practices for, 237–240; health status  
 assessment for, 225–227; Malcolm  
 Baldrige National Quality Program for,  
 249–250; managing, 232–241;  
 monitoring, 231–232; patient  
 satisfaction and patient expectations  
 assessment for, 227–231; practice  
 guidelines for, 237–238; quality of  
 health care for, 241–251; rapid cycle  
 improvement process for, 243–244;  
 root cause analysis for, 240–241;  
 statistical process control for, 244–249
- Population health studies, 89–116;  
 analytic study design example, 90–91;  
 application of, 110–113; designs for:  
 clinical trials, 94–98; cross-sectional  
 studies, 99–101; descriptive studies, 99;  
 ecological studies, 101–102;  
 experimental, 93–94; observational,  
 98–99; prospective studies, 102; retro-  
 spective studies, 102–103; measuring  
 association in prospective, 104–108;  
 measuring association in retrospective,  
 108–110; risk and causation in, 91–93;  
 risk factor associated with, 103;  
 sampling in, 113–114
- Population pyramids, 36, 37
- Population-based models, 210
- Populations: characteristics by age of, 283;  
 commonalities with, 2; concept of,  
 11–12; defining, 11; economic analysis  
 of health care for, 253–279; incidence  
 rate in, 67; managing health care for,  
 12–14; percent distribution of, 284;  
 service area, 283–284; service area  
 health for, 295–301; subpopulations,  
 11–12; U.S. Census Bureau data for,  
 123–124
- Post-neonatal mortality rate, 78, 80
- Potential years of life lost (PYLL), 270;  
 calculation method: before age 75  
 calculation method, 82–83; age group  
 calculation method, 81–82;  
 individual-level calculation method,  
 80–82; defined, 80
- PPS. *See* Prospective payment system
- Practical efficiency, 122
- Practice analysis, 168–169
- Practice guidelines, population health  
 outcomes, 237–238
- Precision. *See* Reliability
- Pregnancy, ICD-10-CM codes for, 24
- Preterm birth complications, DALYs  
 caused by, 271, 272
- Pretest odds, likelihood ratios with, 165
- Prevalence, validity with, 155–156
- Prevalence rate: asthma, 69–70;  
 incidence rate related to, 70, 72; life-  
 time, 69; period, 72; point, 69; women  
 having mammogram, 70–71; women  
 having Papanicolaou test, 70–71
- Prevalence studies. *See* Cross-sectional  
 studies designs
- Preventive medicine, 7–8
- Primary prevention, 7–8, 150–151
- Private insurance: emergency department  
 payments from, 331–332; payment  
 source with, 43. *See also* Health  
 insurance
- Probabilistic inference, 153–154
- Process measures, 222–223
- Professional standards review  
 organizations (PSROs), 223

- Program for All-inclusive Care for the Elderly (PACE), 273
- Propagated epidemics, 179
- Proportionate mortality ratio (PMR), 79–81
- Proportions, 63–65; home health and hospice care agencies by ownership and location, 64
- Proportions analysis, smoking and CHD association in, 108
- Prospective payment system (PPS), 9, 220; advent of, 192; creation of, 194; DRGs with, 194–195. *See also* Inpatient Rehabilitation Facility Prospective Payment System; Outpatient Prospective Payment System
- Prospective studies: design of, 102; measuring association in, 104–108; myocardial infection among HIV infected, 113; odds ratio calculated for, 109
- Psittacosis, cases in U. S. of, 49
- PSQ. *See* Patient Satisfaction Questionnaire
- PSROs. *See* Professional standards review organizations
- Psychoses, emergency department visit with diagnosis of, 339
- Public Health Service, U.S., 5; periodic reports to, 44
- Public policy, 254
- PYLL. *See* Potential years of life lost
- Pyrexia, emergency department visit with diagnosis of, 336
- Q
- QALYs. *See* Quality-adjusted life years
- Q-fever: cases in U. S. of, 49; as notifiable infectious disease, 48
- QIO. *See* Quality improvement organizations
- QLI. *See* Quality of Life Index
- QLQ-C30. *See* Quality of Life Questionnaire
- QoL. *See* Quality of life
- QPP. *See* Quality Payment Program
- Quality improvement organizations (QIO), 223–224
- Quality of care: cause and effect diagrams for, 245; checklists for, 245, 247–248; comparison of, 221–222; control charts for, 245–247; data tools of, 245; defined, 242–243; flow charts for, 245; functions in assessment of, 222; histograms for, 245; Malcolm Baldrige National Quality Program for, 249–250; measurement parameters for, 242; Pareto charts for, 245, 248; population health outcomes with, 241–251; rapid cycle improvement process for, 243–244; scatter diagrams for, 245; statistical process control for, 244–249
- Quality of life (QoL), 84–85; health status with, 25
- Quality of Life Index (QLI), 226
- Quality of Life Questionnaire (QLQ-C30), 214
- Quality Payment Program (QPP), 257
- Quality-adjusted life years (QALYs): cost-utility analysis using, 264

## R

- Rabies: cases in U. S. of, 49; as notifiable infectious disease, 48
- Race and ethnicity: asthma prevalence rate for, 70; births among Medicaid

- beneficiaries for, 111; cancer deaths rates in U.S. by, 31; crude death rate by, 74; death rates for, 32–33; disease patterns according to, 30; doctor visits in U. S. by, 61; East Bank County, 304–305; East Bank Regional Hospital cancer incidence rates by, 296, 297; East Bank Regional Hospital diabetes prevalence by, 298; East Bank Regional Hospital heart disease prevalence by, 297; East Bank Regional Hospital stroke prevalence by, 297; emergency department visits by, 31, 62, 326–327, 330–331; frequency of new cancer cases in U.S. by, 32; HIV cases in U. S., 63; hospital stay in East Bank Regional Hospital by, 285; infant mortality rate in U.S. by, 77; insurance coverage for Bank Regional Hospital by, 294; Lakeshore County, 306–307; long-term care services utilization by, 309; Medicaid coverage for Bank Regional Hospital by, 295; Medicare coverage for Bank Regional Hospital by, 294; neonatal mortality rate in U.S. by, 79; office visits by, 347; patient population by, 346; post-neonatal mortality rate in U.S. by, 80
- Random error, 152
- Randomization, 114
- Rapid cycle improvement process, 243–244
- RASS. *See* Richmond Agitation Sedation Scale
- Rates, 118; age-specific mortality, 75–76; case-fatality, 76–77; cause-specific mortality, 73–75; counts differ from, 58; crude birth, 85; crude death, 73–74; death rate, 28–33, 73–80; defined, 59; denominators of, 59; doctor visits in U. S., 61; emergency department visits in U. S., 62; expressed as percentage, 59; fertility, 85–86; fetal death, 78; HIV case, U. S., 60; homicide, 33; incidence rate, 65–68, 70, 72; infant mortality, 77–78; infection, 184–185; lifetime prevalence, 69; mortality, 73–79, 242; neonatal mortality, 78, 79; numerator of, 59; period prevalence rate, 72; point prevalence, 69; post-neonatal mortality, 78, 80; prevalence rate, 69–72; survival rate, 83; using for comparison, importance of, 61–62
- Ratios, 62–63; age-adjusted diabetes prevalence, 64; cost-effectiveness, 268, 274, 279; HIV case, U. S., 63; hospital standardized mortality, 135–136; likelihood, 163–166, 170; marginal cost-effectiveness, 274, 279; odds, 109, 110, 165; proportionate mortality, 79–81; SMR, 130, 132–135, 140, 148
- RBRVS. *See* Resource-based relative value scale
- Readmission rates, quality of care measurement with, 242
- Receiver-operating characteristic (ROC) curve, 166–168
- Reimbursement methods: accountable care organizations for, 200–201; capitation as, 194; current procedural terminology with, 193; diagnosis-related groups with, 194–195; emergency department visits, 331–332; HH-PPS as, 197–198; hospital

- value-based purchasing, 233; IRF-PPS as, 196–197; Medicare, 282; outpatient prospective payment system as, 9, 195–196; pay-for-performance, 201, 233; prospective payment system as, 9, 192, 194–195, 220; RBRVS as, 198–199; SNF prospective payment system as, 197
- Relative risk: calculation of, 105; cesarean section rates by age with, 112; cesarean section rates by locale with, 112; contingency table explaining, 105; defined, 104; range of, 104; smoking and CHD in analysis of, 106
- Relative value units (RVUs), 198–199
- Reliability: calculation of, 161–162; contingency table for, 161; defined, 160; market research, 210–214
- Reliability index, 161–162
- Repeatability. *See* Reliability
- Reportable diseases, 47–51; analytical analysis for, 51; epidemiologic data for, 51; external data for, 50; internal data for, 50; primary data for, 50; research data for, 50; secondary data for, 50; two types of, 48
- Reproducibility. *See* Reliability
- Residential care community, long-term care services utilization for, 309, 314
- Resource-based relative value scale (RBRVS), 9, 198–199
- Respiratory failure, inpatient deaths in U.S. hospitals from, 42
- Respiratory infections: lower, DALYs caused by, 271, 272; lower, death caused by, 273
- Respiratory system diseases: chronic lower respiratory diseases, 74, 83; ICD-10-CM codes for, 24. *See also* Severe acute respiratory syndrome
- Retrospective studies: breast cancer example of, 103; design of, 102–103; measuring association in, 108–110; odds ratio in, 109–110
- Rheumatism, emergency department visit with diagnosis of, 336
- Richmond Agitation Sedation Scale (RASS), 166
- Risk: causation with, 91–93; defined, 91; theory of modified determinism with, 92
- Risk adjustment, 136–139; rationale for, 137; recovery distribution with, 138, 139; severity of illness distribution with, 138, 139; stratum-specific rates with, 138, 139
- Risk factors, 91–92; cross-sectional studies with exposure to, 99–100; health and disease associated with, 103; marital status associated with chronic disease, 34; observational study designs with, 98–99; occupation with exposures as, 29; smoking and lung cancer, 120–121
- Risk-adjusted capitation rates, 273
- Riverview Hospital: age of patient visiting for, 322–331; background information on, 321–322; emergency department utilization in, 322–331; patient disposition for, 337–340; payment sources for, 331–332; percent occupancy rate for, 292, 293; primary diagnosis at arrival to, 336–337; race of patient visiting for, 326–327, 330–331; sex of patient visiting for, 324–325, 328, 329; time analysis for, 332–335; wait time spent in, 333–334



Riverview Medical Center Hospital, risk adjustment in, 137–139  
 Road traffic accidents: DALYs caused by, 271, 272; death caused by, 273  
 ROC curve. *See* Receiver-operating characteristic curve  
 Rocky Mountain spotted fever, cases in U. S. of, 49  
 Root cause analysis, 240–241; components of, 241  
 Rubella (Measles): cases in U. S. of, 49; death caused by, 273; as notifiable infectious disease, 48  
 Rubeola, as notifiable infectious disease, 48  
 RVUs. *See* Relative value units

## S

Salmonellosis: cases in U. S. of, 49; as notifiable infectious disease, 48  
 Sampling, 113–114; cluster, 114, 204, 205; design, 114; multistage, 114, 204, 205; randomization with, 114  
 Sanitary reform movement, 2  
 SARS. *See* Severe acute respiratory syndrome  
 Scatter diagrams, quality control with, 245  
 SCHIP. *See* State Children's Health Insurance Program  
 Screening, 152  
 Scurvy, 3  
 Seasonal epidemics, chickenpox, 39–40  
 Secondary attack rate, 178  
 Secondary prevention, 8, 150, 151  
 Secular trends, 40–41  
 SEER registries, 187  
 Self-pay: emergency department payments from, 331–332; payment source with, 43  
 Sensitivity, 154–155, 159  
 Sensitivity analysis, 269  
 Sentinel surveillance, 187  
 Septicemia, inpatient deaths in U.S. hospitals from, 42  
 Service area population: characteristics by age of, 283; health, 295–301; percent distribution of, 284  
 Severe acute respiratory syndrome (SARS): hospital workers infected by, 14–15; as notifiable infectious disease, 48; symptoms of, 180; transmission of, 180; worldwide impact of, 180–181  
 Severity: disease patterns, 30; illness distribution with risk adjustment for, 138, 139; MS-DRGs, 10, 195  
 Sex differences: age-adjusted diabetes prevalence ratios by, 64; asthma prevalence rate for, 70; crude death rate by, 74; death rates based on, 29; doctor visits in U. S. by, 61; East Bank County, 304–305; East Bank Regional Hospital cancer incidence rates by, 296, 297; East Bank Regional Hospital diabetes prevalence by, 298; East Bank Regional Hospital discharges by, 288, 289; East Bank Regional Hospital heart disease prevalence by, 297; East Bank Regional Hospital stroke prevalence by, 297; emergency department visits by, 62, 324–325, 328, 329; hospital stay in East Bank Regional Hospital by, 285; injury-related emergency room visit rates by, 30; insurance coverage for Bank Regional Hospital by, 294;



- Lakeshore County, 304, 306; length of hospital stay by, 291; long-term care services utilization by, 309; Medicaid coverage for Bank Regional Hospital by, 295; Medicare coverage for Bank Regional Hospital by, 294; morbidity rates based on, 29; outpatient department visits by, 119, 126
- SF-8 survey, 226
- SF-12 survey, 226
- SF-36. *See* MOS short-form 36
- Shiga-toxin producing *Escherichia aureus*, as notifiable infectious disease, 48
- Shigellosis: cases in U. S. of, 49; incidence rate in U. S. for, 68; as notifiable infectious disease, 48
- Shoreline Regional Hospital, percent occupancy rate for, 292, 293
- Sickness Impact Profile, 226
- SIDS. *See* Sudden infant death syndrome
- Skilling nursing facility (SNF):  
prospective payment system of, 197
- Skin diseases: ICD-10-CM codes for, 24; proportionate mortality ratio with, 81
- Smallpox, 5; as notifiable infectious disease, 48
- Smoking: coronary heart disease with, 106, 108, 110; lung cancer risk with, 120–121; stratification of status of, 121
- SMR. *See* Standardized morbidity or mortality ratio
- SNF. *See* Skilling nursing facility
- Snow, John, 4
- Social epidemiology: aim of, 12; defined, 12
- Socioeconomic status: disease incidence related to, 36; East Bank Regional Hospital diabetes prevalence by, 298; end-stage renal disease risk for, 36; health insurance coverage in U.S. for, 35–36; health status indicated by growth in children for, 35; mental disorders trends for, 35; place with, 38; prognosis for cancer patients across, 33–34; uninsured percentages for persons by, 34–35
- Specificity, 154–155, 159
- Spinal disorders, emergency department visit with diagnosis of, 336–337
- Spotted fever rickettsiosis, as notifiable infectious disease, 48
- Sprains, emergency department visit with diagnosis of, 336
- Standard reference gamble, 264, 265
- Standardization of information, 117–148; described, 118–120; direct method of, 123–129; indirect method of, 129–136; matching with, 121–122; risk adjustment with, 136–139; stratification with, 120–121
- Standardized morbidity or mortality ratio (SMR), 140, 148; defined, 130; invasive carcinoma of cervix with, 134–135; outpatient visits with, 132–133
- Staphylococcus aureus*, HAI caused by, 182
- State Children's Health Insurance Program (SCHIP), 9
- Statistical process control: data tools of, 245; development of, 244; quality of care with, 244–249; Six sigma technologies for, 246–249
- Stratification, 120–121
- Stratified sampling, 204–205
- Stratum-specific rates, risk adjustment with, 138, 139

- Streptococcal disease, cases in U. S. of, 49
- Streptococcal toxic shock syndrome: cases in U. S. of, 49; as notifiable infectious disease, 48
- Streptococcus Pneumoniae*, cases in U. S. of, 49
- Stroke: DALYs caused by, 271, 272; East Bank Regional Hospital prevalence of, 297; inpatient deaths in U.S. hospitals from, 42
- Student's t-distribution, 107
- Study designs: clinical trials, 94–98; cross-sectional studies, 99–101; descriptive studies, 99; ecological studies, 101–102; experimental, 93–94; observational, 98–99; prospective studies, 102; retrospective studies, 102–103; sampling with, 113–114
- Study efficiency, 122
- Subcutaneous tissue diseases, ICD-10-CM codes for, 24
- Sudden infant death syndrome (SIDS): post-neonatal mortality rates with, 78
- Sufficient, but not necessary relationship, 93
- Suicide, whites higher death rates with, 30
- Surgical Safety Checklist, 247
- Surveillance, infectious epidemics management with, 186–188
- Survival rate, 83
- Syncope, emergency department visit with diagnosis of, 339
- Sydenham, Thomas, 3
- Syphilis: African Americans death rates with, 30; cases in U. S. of, 49; incidence rate in U. S. for, 68; as notifiable infectious disease, 48
- Systematic error, 152
- ## T
- T-distribution, 107
- Tertiary prevention, 150, 151
- Tetanus: cases in U. S. of, 49; Hippocrates' descriptions of, 3; as notifiable infectious disease, 48
- Time patterns: chickenpox with, 39–40; disease trends explained with, 39–41; emergency department utilization with, 40–41; epidemics defined by, 39; foodborne outbreak explained with, 40; inpatient deaths from U.S. hospitals with, 41, 42; seasonal epidemics with, 39–40; secular trends with, 40–41
- Time trade-off methods, 264, 265
- TN. *See* True negatives
- Total hip replacement, East Bank Regional Hospital discharge following, 289, 290
- Total knee replacement, 289, 290
- Toxic shock syndrome: cases in U. S. of, 49; as notifiable infectious disease, 48
- TP. *See* True positives
- Traditional utilization projections, 210
- Transmission mechanisms: direct transmission, 177; indirect transmission, 177
- Trichinosis: cases in U. S. of, 49; as notifiable infectious disease, 48
- True negatives (TN), 153
- True positives (TP), 153
- T-test, 107

- Tuberculosis, 4; African Americans death rates with, 30; cases by age for, 28; cases in U. S. of, 49; death caused by, 273; incidence rate in U. S. for, 68; as notifiable infectious disease, 48
- Tularemia, cases in U. S. of, 49
- Typhoid fever: cases in U. S. of, 49; as notifiable infectious disease, 48
- Typhus, Hippocrates' descriptions of, 3
- U**
- UCR costs. *See* Usual, customary, and reasonable costs
- UHC. *See* University Health Consortium
- Universal precautions, 15
- University Health Consortium (UHC), 167–168
- Urinary tract infection, emergency department visit with diagnosis of, 336, 339
- Usual, customary, and reasonable (UCR) costs, 198
- V**
- Validity: arthritis screening tests with, 156; colorectal cancer screening tests with, 157; contingency table for, 153; defined, 152; managerial aspects of, 158–160; market research, 210–214; parameters for, 153, 156, 157; predictive, 157, 159; prevalence with, 155–156; sensitivity and specificity with, 154–155, 159
- Varicella (Chickenpox): cases in U. S. of, 49; as notifiable infectious disease, 48; seasonal epidemic of, 39–40
- Veterans RAND 12-item Health Survey (VR-12), 232
- Vibriosis, as notifiable infectious disease, 48
- Virulence, 177
- Visit Specific Satisfaction (VSS), 229
- Vital statistics, 45–46
- Vitamin C deficiency, 3
- VR-12. *See* Veterans RAND 12-item Health Survey
- VSS. *See* Visit Specific Satisfaction
- W**
- Waterborne disease outbreak, as notifiable infectious disease, 48
- WEC, Inc. *See* Westside Elderly Care, Inc.
- WEC-EB. *See* Westside Elderly Care—East Bank
- WEC-LC. *See* Westside Elderly Care—Lakeshore County
- West Nile virus, 181
- Westbank Hospital, percent occupancy rate for, 292, 293
- Westside Elderly Care, Inc. (WEC, Inc.): background information on, 304, 308; population characteristics of, 304–308; staffing for, 319; utilization of services for, 308–318
- Westside Elderly Care—East Bank (WEC-EB), 304, 308
- Westside Elderly Care—Lakeshore County (WEC-LC), 304, 308
- Westside Memorial Hospital: case-fatality rates for MI in, 126–129; percent occupancy rate for, 292, 293
- Whites: asthma prevalence rate for, 70; births among Medicaid beneficiaries for, 111; cancer deaths rates in U.S. for, 31; crude death rate by, 74; death rates

- with, 30; doctor visits in U. S. by, 61;  
 emergency department visits in U.S.  
 for, 31, 62; end-stage renal disease risk  
 for, 36; frequency of new cancer cases  
 in U.S. for, 32; hospital stay in East  
 Bank Regional Hospital for, 285; infant  
 mortality rate in U.S. for, 77; long-term  
 care services utilization by, 309;  
 neonatal mortality rate in U.S. for, 79;  
 outpatient visits for, 132; post-neonatal  
 mortality rate in U.S. for, 80;  
 prevalence of women having  
 mammogram for, 71  
 WHO. *See* World Health Organization  
 Workers' compensation: emergency  
 department payments from, 331–332;  
 payment source with, 43  
 World Bank, burden of disease study  
 by, 270  
 World Health Organization (WHO):  
 burden of disease study by, 270; disease  
 classification of, 22; Surgical Safety  
 Checklist of, 247
- X**
- X<sup>2</sup> distribution, 107
- Y**
- Years of Life Lost (YLLs), 271  
 Years of Life Lost due to Disability  
 (YLLDs), 271  
 Yellow fever, 5; as notifiable infectious  
 disease, 48  
 YLLDs. *See* Years of Life Lost due to  
 Disability  
 YLLs. *See* Years of Life Lost  
 Youth Risk Behavior Surveillance System  
 (YRBSS), 42
- Z**
- Zika virus: cause of, 181; infants, effects  
 of, 15; as notifiable infectious disease,  
 48; pregnant woman with, 182;  
 symptoms of, 182