

Index

- 2log likelihood *see* likelihood ratio statistic
- a priori* approach to probability 26
- addition rule 26
- adequacy of fit 97, 98, 167
- adjusted odds ratio 88
- adjusted R^2 84
- administrative censoring 134
- aggregate level analysis, clustered data 127, 128
- agreement
 - limits of 119
- agreement, assessing
 - categorical variables 118, 120
 - measurement variability and error 118
 - numerical variables 118–21
 - reliability 118
- allocation
 - bias 42, 104
 - clinical trials 40
 - random *see* randomization
 - systematic 40
 - treatment 40
- alpha (α) 52, 108
- alternative hypothesis, defining the 50
- Altman's nomogram 108–9, 149
- analysis of covariance 84–5
- analysis of variance (ANOVA) 63
 - F -ratio 80
 - Kruskal-Wallis test 64
 - one-way *see* one-way analysis of variance
 - repeated measures 125
 - table 78–9
- ANOVA *see* analysis of variance
- antilog 30
- area under the curve (AUC) 26, 27, 125
- arithmetic mean 22, 23
 - see also* mean
- ascertainment bias 102
- ASCII files 16
- assessment bias 102
 - clinical trials 42
- association
 - in contingency table 72, 73–4
 - in correlation 75
- assumptions, checking
 - linearity 107
 - Normal distribution 106
 - reasons for 106
 - sensitivity analysis 107
 - variances, equality of 106
 - when assumptions are not satisfied 107
- attrition bias 102
- automatic selection procedures 99
- average 22
 - arithmetic mean 22, 23
 - geometric mean 22, 23
 - median 22, 23
 - mode 22, 23
 - weighted mean 22, 23
- back-transformations 30
- backwards selection 99
- bar charts 20
 - clustered 21
 - segmented 20, 21
- Bartlett's test 106
 - one-way ANOVA 64
- Bayes theorem 136–7
- Bayesian methods 136
 - diagnostic tests 136–7
 - disadvantages 137
- beta (β) 52, 80
- between-subject/group variabilities 25, 63, 118
- bias 37, 102
 - cohort studies 45
 - confounding *see* confounding
 - evidence-based medicine 122
 - information 102–3
 - missing data 18
 - on-treatment analyses 43
 - selection 102
- bimodal distributions 21
- binary (dichotomous) variables 14, 122
 - explanatory 78, 85, 98–101
 - in logistic regression 88–90
 - in meta-analysis 130
 - in multiple regression 84
- binary outcomes and logistic regression 88
- Binomial distribution 29
 - confidence interval for the proportion 34
 - Normal approximation 28
 - single proportion test 66
- bioequivalence trials 51
- Bland and Altman diagram 119
- blinding/masking
 - clinical trials 42
 - evidence-based medicine 122
- blocked randomization 40
- blocking 38
- Bonferroni correction 53
- bootstrapping 35
 - prognostic scores 139
- box (box-and-whisker) plot 21, 24
- Brier score 138
- British Standards Institution repeatability coefficient 119
- calibration, prognostic scores 139
- carry-over effects 38
- case-control studies 36, 47
 - advantages 48
 - analysis 48
 - cases, selection of 47
 - controls, selection of 47
 - disadvantages 48
 - matched 47–8
 - risk factors, identification of 47
 - unmatched 48
- cases 47
 - incident 47
 - prevalent 47
- categorical data (variables) 14, 66–8
 - coding 16, 17
- categorical (qualitative) data 14
 - agreement, assessing 118, 120
 - data entry 16
 - diagrammatic display 20
 - error checking 18
 - logistic regression 90
 - more than two categories 72–4
 - multiple linear regression 84
- causal modelling 105
- causal pathway and confounding 105
- causality, assessing 37
- causality, in observational studies 36
- censored data 15, 133
 - left- 133
 - right- 133
 - survival analysis 133, 134
- censuses 37
- Central Limit Theorem 32
- central tendency bias 102
- centring, linear regression 81
- channelling bias 104
- checking assumptions *see* assumptions, checking
- Chi-square for covariates *see* model Chi-square test
- Chi-squared (χ^2) distribution 28, 143
- Chi-squared (χ^2) test
 - in 2×2 tables 69
 - for covariates (model Chi-square) 88, 97
 - independent groups 69
 - in $r \times c$ table 72
 - for trend in proportions 72–3, 74
 - for two proportions
 - independent data 69
 - paired data 57, 70–1
- CI *see* confidence intervals
- classification table 89
- clinical cohorts 46
- clinical heterogeneity 131
- clinical trials 36, 40–3
 - avoiding bias in 37
 - blinding/masking 42
 - CONSORT Statement 40, 41–2
 - cross-over design 38, 39
 - endpoints, primary and secondary 40
 - ethical issues 40
 - evidence-based medicine 122
 - inclusion/exclusion criteria 41
 - informed consent 40
 - intention-to-treat analysis 42, 122
 - patient issues 43
 - phase I/II/III 46
 - placebo in 40
 - protocol 41–2, 43
 - sequential 42
 - size 42
 - subgroup analyses 40
 - treatment allocation 40–2
 - treatment comparisons 40
- cluster randomization
 - clinical trials 40–2
 - observation, unit of 37
- cluster-specific models 127
- clustered bar charts 21
- clustered data 124
 - appropriate analyses 124–5
 - displaying 124

- inappropriate analyses 124
- rates 92
- regression methods 127–9
- Cochran’s Chi-squared test for homogeneity 132
- Cochrane Collaboration 130
- coding
 - data 16, 17
 - missing values 16, 17
- coefficient
 - correlation *see* correlation coefficient
 - intraclass correlation 118, 128
 - logistic regression 87
 - partial 84
 - regression 78
 - repeatability/reproducibility 118, 120
 - of variation 25
- Cohen’s kappa 118
 - weighted 118
- cohort studies 36, 44–6
 - advantages 45
 - analysis of 44–5
 - clinical cohorts 46
 - confounding 104
 - disadvantages 45
 - dynamic 44
 - fixed 44
 - follow-up of individuals 44
 - historical 44
 - management of study 45–6
 - outcomes and exposures, information on 44
 - selection of cohorts 44
- collinearity
 - logistic regression 89
 - multiple linear regression 85
- column charts *see* bar charts
- competing risks 134–5
- complementary events 26
- complete randomized studies 38
- composite endpoints 39
- computer output 150–62
- computer packages, data entry 16–17
- conditional logistic regression 90
- conditional probability 26, 136
- confidence intervals (CI) 34, 35, 122
 - 95% 80
 - for correlation coefficient 75, 76
 - for difference in two means 60, 62
 - for difference in two medians 61
 - for difference in two proportions
 - independent groups 69, 70
 - paired groups 70, 71
 - interpretation of 34–5, 54, 60, 122
 - for mean 34, 35
 - for mean difference 57–58, 59
 - for median 145
 - for median difference 57, 58–9
 - in meta-analysis 130–1, 132
 - multiplier for calculation of 142, 143
 - for proportion 34–5
 - for regression coefficient 80
 - for relative risk 45, 46
 - for slope of regression line 80, 81
 - versus hypothesis testing 51
- confidence limits 34
- confounding 103
 - and causal pathway 104–5
 - clinical trials 40
 - cohort studies 45
 - dealing with 103–4
 - in non-randomized studies 104
 - time-varying 105
- connecting lines in diagrams, use of 21
- consent, informed
 - clinical trials 43
 - cohort studies 46
- CONSORT Statement 40, 41–2, 112
- contingency tables
 - 2×2 69
 - $2 \times k$ 72
 - $r \times c$ 72
- continuity correction
 - sign test 55
 - single proportion test 66
- continuous data 14
- diagrammatic display 20–1
- continuous probability distributions 26–7, 28
- controls 37, 47
 - clinical trials 40
 - see also* case–control studies
- convenience factor 92
- convenience sample 32
- Cook’s distance 84
- correlation 75–7
 - linear 75–6
- correlation coefficient
 - assumptions 75
 - confidence interval for 76, 77
 - hypothesis test of 76, 77
 - intraclass 118, 128
 - misuse of 76
 - non-parametric 76
 - Pearson 75, 77, 147
 - Spearman’s rank 76, 77, 147
 - square of (r^2) 75
- counts 29
- covariance, analysis of 85
- covariate, multiple linear regression 84
- Cox proportional hazards regression model 90, 134, 135
- critical appraisal 122
- cross-over studies 38, 39
- cross-sectional studies 36, 37
 - repeated 36, 37
- cross-sectional time series models, clustered data 127
- cross-validation 139
- c* statistic 116, 117
- cumulative meta-analyses 131
- cut-off values 116
- data
 - categorical *see* categorical (qualitative) data
 - censored 15
 - clustered 92, 124–6, 127–9
 - coding 16, 17
 - derived 14–15
 - describing 22–5
 - diagrammatic display *see* diagrammatic display of data
 - distinguishing between types 14
 - dredging 53
 - error checking 18–19
 - graphical display 20–21, 124
 - missing 16, 18
 - numerical *see* numerical (quantitative) data
 - paired 57
 - range checking 18
 - and statistics 14
 - summarizing 22, 24
 - transformation 30–1
 - see also specific types*
- data entry 16
- Data Safety and Monitoring Committee (DSMC) 40
- dates
 - data entry problems 16
 - error checking 18
- deciles 24
- decision making in hypothesis testing 52
- Declaration of Helsinki 43
- degrees of freedom (df) 34–5
 - Chi-squared distribution 28
 - F*-distribution 28
 - t*-distribution 28
- delimiters 16
- dependent (outcome, response) variables 78
 - binary 105
 - categorical 90
- derived data 14–15
- design *see* study designs
- deviance *see* likelihood ratio statistic
- df see* degrees of freedom
- diagnostic tests 115–16, 136–7
 - in Bayesian framework 136–7
- diagrammatic display of data 20
 - clustered data 124
 - connecting lines 21
 - one variable 20–1
 - outliers, identifying 21
 - two variables 21
- diagrammatic display of results 112
- dichotomous variables 14
- discrete data 14
- discrete probability distributions 26, 28–9
- discriminant analysis 138
- disease register 46
- dispersion *see* spread
- distribution
 - bimodal 20
 - continuous probability 28
 - discrete probability 29
 - empirical frequency 20
 - frequency 20, 21
 - probability 26
 - sampling 32–3
 - skewed 20
 - Standard Normal 27
 - symmetrical 20
 - theoretical 26–9
 - uniform 20
 - unimodal 20
 - see also specific types*
- distribution-free tests *see* non-parametric tests
- dot plots 20, 21
- double-blind trials 42
- dummy (indicator) variables
 - multiple linear regression 84
 - sensitivity analysis 107
- Duncan’s test 63
- dynamic cohort studies 44
- ecological fallacy 102–3
 - clustered data 125
 - meta-regression 103, 131

- ecological studies 37, 102–3
- effect (of interest) 52
 - in evidence-based medicine 122
 - in meta-analysis 130–1
 - importance of 122
 - power of test and 52
 - in sample size calculation 108
- effect modifiers 45
- efficacy 40
- empirical frequency distributions 20, 26
 - displaying 20–1
- endpoints of studies
 - choosing 39
 - clinical trials 40
 - primary 40
 - secondary 40
- epidemiological studies 36
- EQUATOR Network 112
- equivalence range 51
- equivalence trials 51
- error
 - residual 78
 - Type I 52
 - Type II 52
 - typing 18
 - variation 53
- estimated relative risk 45
- ethical issues
 - clinical trials 43
 - cohort studies 46
 - power of a test 52
- ethics committees
 - clinical trials 43
 - cohort studies 46
- even numbers 22
- evidence, levels of 122
- evidence-based medicine (EBM) 122
 - approach 122–3
 - systematic reviews 130
- exclusion criteria, clinical trials 43
- expected frequency 69
- experimental studies 36, 37
- experimental units 38
- explanatory (independent, predictor) variables
 - 78, 85, 98–101
 - in logistic regression 88–89
 - in multiple regression 84–5
 - nominal 84, 99
 - numerical 103
 - ordinal 98
- Exponential model, survival analysis 134
- exposure variable 45, 102–5
- external validation 139
- extra-Binomial variation 89–90, 94
- extra-Poisson variation 93–4

- factorial experiments 38–9
- Fagan's nomogram 136, 137
- false negative rate 89
- false positive rate 89
- F*-distribution 28
 - one-way ANOVA 63
 - statistical table 142, 144
- Fisher's exact test 69, 72
- fitted value 70
- fixed cohort studies 44
- fixed effect model 130

- follow-up
 - cohort studies 44, 45
 - loss to *see* loss to follow-up
 - person-years of 92
- forest plots 131
- forms 16
 - multiple 16
- forwards selection 98
- frailty models 134
- F*-ratio 80
- free format 16
- frequency 20, 69, 72
 - displaying 20–1
 - observed and expected 69
 - shape 21
 - table of 115
- frequency matching, case–control studies 47
- frequentist approach to probability 26, 136
- F*-test *see* variance-ratio test
- funding bias 102
- funnel plots 131

- G see* model Chi-square test
- Galton, Sir Francis 79
- Gaussian distribution *see* Normal distribution
- generalized estimating equations (GEE) 128
 - example 129
- generalized linear models (GLMs) 96
 - adequacy of fit, assessing 96–7
 - likelihood and maximum likelihood estimation 96
 - regression diagnostics 97
 - types 96
- geometric mean 22, 23
- G*-estimation 105
- gold standard tests 115
- Gompertz model 134
- goodness of fit
 - linear regression 79, 80
 - logistic regression 89
 - multiple linear regression 85
- Gossett, W. S. ('Student') 28
- gradient, of regression line 78
- graphical display
 - of data *see* diagrammatic display of data
 - of results 112

- Harrell's *c* statistic 138–9
- hazard 134
- hazard ratios 134
- healthy entrant effect 102
 - cohort studies 44
- Helsinki, Declaration of 43
- heterogeneity
 - clinical 131
 - statistical 130
 - of variance 106, 130
- hierarchical models, clustered data 127
- histograms 20–1
 - logistic regression 89
- historical cohorts 44
- homogeneity, of variance 130
- homoscedasticity 106
- Hosmer–Lemeshow test 139
 - logistic regression 89
 - prognostic score 139
- hypothesis, null and alternative 50

- hypothesis testing 50–1
 - categorical data 66
 - consequences of 52
 - for correlation coefficient 75–6
 - errors in 52–3
 - likelihood ratio 88
 - meta-analysis 130
 - more than two categories 72
 - more than two means 63
 - multiple 53
 - non-parametric 51
 - presenting results 112
 - in regression 80–1
 - single mean 54–5
 - single proportion 66
 - two means
 - related groups 57–59
 - unrelated groups 60–2
 - two proportions
 - independent groups 69
 - related groups 69–70
 - two variances 105, 106
 - versus confidence intervals 52
 - Wald 88
- I*² 130
- identity link, generalized linear models 96
- inappropriate analyses 124
- incidence rate 92
- incidence rate ratio *see* relative rates
- incident cases 47
- inclusion criteria, clinical trials 43
- independent variables *see* explanatory variables
- indicator (dummy) variables
 - multiple linear regression 84
 - sensitivity analysis 107
- inferences 32
- influence plots 131
- influential observations
 - linear regression 80
 - logistic regression 90
 - multiple linear regression 85, 107
- information bias 102–3
- informative censoring 134
- informed consent
 - clinical trials 43
 - cohort studies 46
- intention-to-treat (ITT) analyses
 - clinical trials 43
 - evidence-based medicine 122
- interaction 38, 99–100
- interdecile range 24
- interim analyses
 - clinical trials 40
 - hypothesis testing 53
- intermediate variables and confounding 105
- internal–external cross-validation 139
- internal pilot studies 108
- internal validation 139
- interpolation, unpaired *t*-test 61
- interquartile range 24
- inter-subject/group variation 25, 63, 118
- interval estimates 32, 34
- intraclass correlation coefficient (ICC) 118, 119
 - clustered data 128, 129
- intra-subject/group variation 25, 63, 118

- jackknifing 35

- Kaplan–Meier curves 133
- Kappa κ
 - Cohen's 118, 120
 - weighted 118
- k*-fold cross-validation 139
- Kolmogorov–Smirnov test 106
- Kruskal–Wallis test 64
 - clustered data 124, 126
- lead-time bias 102
- least squares *see* method of least squares
- leave-one-out cross-validation 139
- left-censored data 133
- Lehr's formula 109
- Levene's test 106
 - one-way ANOVA 63
- leverage 85
- lifetable approach, survival analysis 133
- likelihood 96
 - 2log 89, 97
 - Bayes theorem 136
- likelihood ratio (LR) 97
 - Bayes theorem 137
 - diagnostic tests 116, 117
- likelihood ratio statistic (LRS) 97
 - logistic regression 88, 89
 - Poisson regression 93
- likelihood ratio test
 - clustered data 128
 - logistic regression 88–9
- Likert scales, central tendency bias 102
- limits of agreement 119
- Lin's concordance correlation coefficient 119
- linear regression 78
 - ANOVA table 78–9
 - assumptions 78, 79, 80
 - and clustered data 124
 - goodness of fit, assessing 80
 - improving the interpretation of the model 81
 - method of least squares 78
 - multiple 84–7
 - multivariable 84
 - outliers and influential points 80
 - regression line 78, 80–1
 - regression to the mean 79
 - simple 78, 82
 - theory 78
- linearity
 - checking assumptions 107
 - transformations 30, 31
- link function, generalized linear models 96
- location, measures of 22
- log log plots 134
- logarithmic transformation 30
- logistic regression 88
 - adequacy of model, assessing 89–90
 - conditional 90
 - diagnostics 90
 - equation 88
 - explanatory variables 88–9
 - multinomial (polycotomous) 90
 - odds ratio and relative risk, comparing 90
 - ordinal 90
 - prognostic scores 138
- logit (logistic) transformation 31
- Lognormal distribution 28, 29
 - transformations 30
- log-rank test 134
- longitudinal studies 36, 37
 - clustered data 127
- loss to follow-up
 - bias 102
 - cohort studies 44, 45
 - sample size adjustment 109
- Mann–Whitney *U* test 60
- Mantel–Haenszel procedure 69, 103
- marginal structural models 105
- marginal total 69
- masking *see* blinding/masking
- matching
 - case–control studies 47–8, 90
 - conditional logistic regression 90
 - confounders 103, 104
- mathematical models 96
- maximum likelihood estimation (MLE) 96
 - clustered data 127
- McNemar's test 70
 - confounding 103
- mean 23
 - confidence interval for difference in two 60, 62
 - difference 57, 60, 62
 - weighted 22, 23
- mean Briar score 138
- mean square 63
- measurement bias 102
- measurement error 118
- measurement variability 118
- measures
 - of location 22–23
 - of spread 24–25
- median 22, 23, 24
 - difference between two 57–8, 61
 - ranks for confidence intervals for the 142, 145
 - survival time 133
 - test for a single 54–6
- Medline 122
- meta-analysis 130
 - advantages and disadvantages 131
 - sensitivity analysis 107, 131
 - statistical approach 130–1
- meta-regression 131
 - ecological fallacy 103, 131
- method agreement 119
- method of least squares (ordinary least squares, OLS) 96
 - clustered data 127, 129
 - linear regression 78
- misclassification bias 102
- missing data 16, 18
- missing values
 - coding 16, 19
 - handling 18, 19
- mixed models, clustered data 127
- mode 22, 23
- model Chi-square test 97
 - logistic regression 89
 - Poisson regression 93
- models
 - Cox proportional hazards regression 90, 134, 135
 - Exponential 134
 - fixed effect 130
 - generalized linear 96–7
 - Gompertz 134
 - hierarchical 127
 - logistic regression 88
- multi-level 78, 84–6
- multivariable 94
- over-fitted 99
- Poisson regression 93–4, 134
- random effects 128
- random intercepts 129
- random slopes 128
- regression 134
 - statistical 88–9, 96, 102–4
 - univariable 99
 - Weibull 134
- mortality rates 92
- multicentre studies 37
- multi-coded variables 16
- multi-collinearity 100
- multilevel models, clustered data 127
- multiple forms per patient 16
- multiple hypothesis testing 53
- multiple linear regression 84
 - analysis 85
 - analysis of covariance 84–5
 - assumptions 84
 - categorical explanatory variables 84
 - explanatory variables, choice of 85
 - outliers and influential points 85, 107
- multiplication rule 26
- multivariable regression
 - confounding 103, 104
 - see also* logistic regression; multiple linear regression; Poisson regression
- multivariate analysis 84
 - hypothesis testing 53
- mutually exclusive (categories) 69
- negative controls 40
- negative predictive value 116
- negatively skewed distributions (NPV) 21
 - square transformation 31
- nested models 97
- nominal data 14
 - multinomial logistic regression 90
 - multiple linear regression 84
- nominal significance level 53
- non-inferiority trials 51
- non-parametric (distribution-free, rank) tests 51, 107
 - for more than two independent groups 64
 - for single median 54–6
 - for Spearman correlation coefficient 76
 - for two independent groups 61–2
 - for two paired groups 57–8
- Normal (Gaussian) distribution 27, 28
 - approximation to Binomial distribution 28, 66
 - approximation to Poisson distribution 29
 - in calculation of confidence interval 34
- Normal plot 106
- normal range *see* reference intervals
- null hypothesis, defining the 50
- number of patients needed to treat (NNT) 122
- numerical (quantitative) data 14
 - agreement, assessing 118–21
 - data entry 16
 - diagrammatic display 20–1
 - error checking 18
 - more than two groups 63–5
 - single group 54–6
 - two related groups 57–9
 - two unrelated groups 60–2
- numerical results, presenting 112

- observation 14
 - defining the unit of 36–7
- observational databases 46
- observational studies 36, 37
- observed frequency 69
- observer bias 102
- odds (of disease)
 - adjusted 88
 - case–control studies 48
 - logistic regression 88, 90
 - posterior 137
 - prior 137
- offsets, Poisson regression 93
- one-sample *t*-test 54
- one-tailed tests 50
- one-way analysis of variance 63–4
 - and clustered data 124
- on-treatment analyses, clinical trials 43
- ordinal data 14
 - multiple linear regression 84
 - ordinal logistic regression 90
- ordinal logistic regression 90
- ordinary least squares (OLS) *see* method of least squares
- outliers 18
 - checking for 18
 - handling 18
 - identifying using graphical methods 21
 - linear regression 80
 - logistic regression 90
 - multiple linear regression 85
- overdispersion
 - logistic regression 89
 - Poisson regression 93–4
- over-fitted models 99
- overview *see* meta-analysis

- paired data 57
 - categorical 69, 70
 - numerical 57–8
- paired *t*-test 57
 - Altman’s nomogram 109
 - confounding 103
 - repeated measures ANOVA 125
- pairwise matching 47
- panel models, clustered data 127
- parallel studies 38, 39
- parameters
 - point estimates 32
 - probability distributions 26
- parametric tests 51
- partial likelihood, survival analysis 134
- partial regression coefficients 84
- Pearson’s correlation coefficient 75–6
 - intraclass correlation coefficient 119
 - statistical table 142, 147
- percentages 14
- percentiles 24
 - ranges derived from 24–5
- pie charts 20
- pilot studies 108
- placebos 14
- point estimates 32
- point prevalence 37
- Poisson distribution 29, 92
 - transformations 30
- Poisson regression 92
 - coefficients 93
 - computer output 93
 - equation 92–3
 - extra-Poisson variation 93–4
 - groups, entering data for 93
 - offset, use of an 93
 - variables that change over time 93
- polychotomous linear regression 90
- polynomial regression 99
- population 14, 32
- positive controls 40
- positive predictive value (PPV) 116, 137
- positively skewed distributions 21
 - logarithmic transformation 30
- posterior odds 136, 137
- post-hoc* comparisons 63
- post-test probability 137
- power 52–3
 - curves 52–3
 - sample size 52, 53, 108, 109–10
 - statement 109
- precision 32–3, 112, 122
 - in systematic reviews 130–1
- predictive efficiency, indices of 89
- predictive values
 - negative 116
 - positive 116
- predictor variables *see* explanatory variables
- preferences, analysing 66
- pre-test probability 137
- prevalence 115, 137
- prevalent cases 47
- primary endpoint of trial 40
- prior odds 137
- prior (pretest) probability, Bayes theorem 136, 137
- probability 26
 - addition rule 26
 - a priori* 26
 - Bayesian approach 136, 137
 - complementary 26
 - conditional 26, 136
 - discrete 26, 29
 - frequentist 26, 136
 - multiplication rule 26
 - posterior (post-test) 136, 137
 - prior (pre-test) 136, 137
 - subjective 26
 - survival 133
- probability density function 26
 - Normal distribution 26–7
- probability distributions 26–9
 - continuous 26–7, 28
 - discrete 28–9
 - theory 26
- prognostic index 138
- prognostic scores 139
 - developing 139
 - performance assessment 138–9
 - reasons for 138
- propensity score approach, confounding 103–4
- proportion(s)
 - confidence interval for difference in two 69–71
 - logit transformation of 31
 - sampling distribution 32–3
 - sign test for 66–7
 - standard error of 33
 - test for single 66–8
- test for trend in 72–4
- test for two
 - independent groups 69
 - related groups 70–1
- proportional hazards regression model (Cox) 134, 135
- prospective studies
 - cohort studies 44
 - longitudinal studies 36, 37
- protocol 42–3
 - deviations 43
- protocols, clinical trials 43
- publication bias 102
 - meta-analysis 131
- P*-value 50–1
 - explanation 50–1
 - obtaining the 50
 - post-hoc* adjustment of 53
 - using the 50–1

- qualitative data *see* categorical (qualitative) data
- quality, of studies in meta-analysis 131
- quantitative data *see* numerical (quantitative) data
- quartiles 24
- questionnaires 16
- quick formulae in sample size estimation 108
- QUORUM Statement 112
- quota sampling 32
- quotients 14

- r^2 75
- R^2 79, 85
 - adjusted 85
 - linear regression 79, 80
- $r \times c$ contingency table 72
- random effects
 - meta-analysis 130–1
 - meta-regression 131
 - models, clustered data 127–8, 129
- random intercepts linear two-level model 127, 129
- random measurement error 118
- random numbers table 142, 148
- random samples 32
- random slopes model 128, 129
- random variables 26
- random variation 38
- randomization 122
 - blocked 40
 - cluster 40
 - stratified 40
- randomization, clinical trials 40–2
- randomized controlled trials (RCTs) 40
 - and confounding 104
 - evidence-based medicine 122
- randomized studies, complete 38
- range 24, 25
 - checking 18
 - derived from percentiles 24–5
 - interdecile 24
 - interquartile 24
 - normal *see* reference intervals
 - presenting results 112
 - reference (normal) 24, 26–7, 115
- rank correlation coefficient *see* Spearman’s rank correlation coefficient
- rank (non-parametric) tests 51, 107
- ranks for confidence interval for the median 142, 145
- rate ratio *see* relative rates

- rates 14, 92
 - features 92
 - relative 92
 - and risk, comparison between 92
- ratio 14
 - F-ratio 80
 - hazard 134
 - incidence rate 92
 - likelihood 96, 116, 137
 - odds 48, 88, 92
 - rate 92
- recall bias
 - case-control studies 48
 - cohort studies 45
- receiver operating characteristic (ROC) curve 116
 - area under the (AUROC) 116, 138
 - logistic regression 89
 - prognostic scores 138, 140
- reciprocal transformation 31
- reference categories 84
- reference intervals (normal range) 24, 115
 - calculation 115
 - Normal distribution 27
- reference range 24
- regression
 - clustered data 127–9
 - Cox 90, 134, 135
 - diagnostics 90, 97
 - dilution bias 102
 - linear *see* linear regression
 - logistic 88–90, 156
 - methods 125
 - models, survival data 133
 - Poisson 92–5
 - polynomial 99
 - presenting results in 112–13
 - simple 78, 80–82
 - to mean 79
- regression coefficients 78
 - linear 78
 - logistic 88
 - partial 84
 - Poisson 93
- regression line 78
 - goodness of fit 78, 79, 80
 - prediction from 80–82
- relative frequency distributions 20
 - displaying 20–1
- relative hazards 134
- relative rates 92
 - Poisson regression 93
- relative risk (RR) 45
 - cohort studies 45
 - in meta-analysis 45
 - logistic regression 88, 90
 - odds ratio as estimate of 88
- reliability 118
 - indices of 119, 121
- repeatability 118–19
- repeated cross-sectional studies 36, 37
- repeated measures
 - analysis of variance 125
 - data 124
 - models, clustered data 127
- replication 38
- reporting bias 102
- reproducibility 118–19
- rescaling, linear regression 81
- residual error 78
- residual variance/residual mean square 63, 79
- residual variation 80
- residuals 78
 - linear regression 78, 80
 - multiple linear regression 85
- response bias 102
- response rates 38
- response variables *see* dependent variables
- restricted/blocked randomization 40
- results, presenting 112–13
- retrospective studies
 - cohort studies 44
 - longitudinal studies 36, 37
- right-censored data 133
- risk factors
 - case-control studies 47
 - cohort studies 44–5
- risk scores *see* prognostic scores
- risks, competing 134–5
- robust analysis 107
- robust standard errors, clustered data 127, 128
- robustness 107
 - one-sample *t*-test 54
 - one-way ANOVA 64
- ROC curves *see* receiver operating characteristic curves
- sample 14, 32
 - convenience 32
 - random 32
 - representative 32
 - statistic 32
 - training 139, 140
 - validation 139, 140
 - size 38, 52, 106–10
- sample size
 - Altman's nomogram 108–9, 110, 111, 149
 - bias 102
 - calculations 108–11
 - importance 108
 - power of a test 52, 53, 108, 109–10
 - quick formulae 109
 - requirements 108
 - study design 38
- sample statistic 32
- sampling
 - error 32
 - distribution 33
 - frame 32
 - point estimates 32
 - quota 32
 - reasons for 32
 - standard deviation vs standard error of the mean 32
 - standard errors 32
 - systematic 32
 - variation 32
- sampling distribution
 - of the mean 32
 - of the proportion 33
- saturated models 97
- scale parameter, Poisson regression 93
- scaling, linear regression 81
- scatter diagrams 20, 21
 - correlation analysis 75
 - linear regression 78, 80
- Scheffe's test 63
- scores 14
- screening 79
- SD *see* standard deviation
- SE *see* standard error
- secondary endpoints of trials 40
- segmented bar charts 20, 21
- selection bias 102
 - cohort studies 45
- sensitivity analysis 107
 - logistic regression 89
 - meta-analysis 131
 - multiple linear regression 84
 - outliers 18
- sensitivity of a test 115–16
- sequential trials 42
- Shapiro–Wilk test 106
- shrinkage estimates 127
- sign test 55
 - paired data 57–58
 - for a proportion 66–7
- significance level 51
 - multiple hypothesis testing 53
 - and power 52–3
 - and sample size 108
- significance testing *see* hypothesis testing
- significant result 50
- Simpson's (reverse) paradox 69
 - confounding 103
- single-blind trials 42
- single-coded variables 16
- single proportion test 66
- skewed to the left distributions 21
 - square transformation 31
- skewed to the right distributions 21
 - logarithmic transformation 30
- software, data entry 16–17
- Spearman's rank correlation coefficient 76
 - statistical table 142, 147
- specificity 115–16, 117
 - logistic regression 89
- spread 24
 - range 24–5
 - standard deviation 25
 - variation within- and between-subjects 25
- square root transformation 30
- square transformation 31
- standard deviation (SD) 25, 32
 - SD vs SEM 32
- standard error
 - of the mean (SEM) 32, 112
 - presenting results 112
 - of the proportion 33
 - robust 127, 128, 129
- Standard Normal distribution 27
 - statistical tables 142, 143, 144
- standardized difference 108
- Standardized Normal Deviate (SND) 27
- statistic
 - sample 32
 - test 50
- statistical heterogeneity 130
- statistical homogeneity 130
- statistical packages, data entry 16–17
- statistical tables 142–8
- statistics 14
 - probability distributions 26
- stem-and-leaf plots 21
- stepwise selection 99

- stratification 38
 - stratified randomization 40
- STROBE Statement 112
- Student's *t*-distribution *see t*-distribution
- study designs 36, 38
 - bias 37
 - blocking (stratification) 38
 - causality, assessing 37
 - controls 37
 - cross-over 38, 39
 - cross-sectional studies 37
 - endpoints, choosing 39
 - experimental studies 36
 - factorial experiments 38–9
 - longitudinal studies 37
 - multicentre studies 37
 - observational studies 36
 - parallel 38, 39
 - replication 38
 - sample size 38
 - unit of observation, defining the 36–7
 - variation 38
- subgroups
 - clinical trials 40
 - confounding 103, 104
 - hypothesis testing 53
- subjective approach to probability 26
- summary measures 124–5
 - aggregate level analysis 127
 - of location 22–3
 - of spread 24–5
- surrogate marker endpoints 39
- surveys 37
- survival analysis 94, 133
 - censored data 133
 - comparing survival data 134
 - Cox proportional hazards regression 134
 - displaying survival data 133
 - Kaplan-Meier log-rank test 134
 - problems 134–5
 - summarizing survival data 134
- survival curves 133
- survival probability 133
- survivorship bias 102
- symmetrical distributions 21
- systematic allocation, clinical trials 40
- systematic measurement error 118
- systematic reviews 130
 - meta-analysis 130–2
- systematic sampling 32
- tables
 - 2 × 2 table 69
 - contingency of frequencies 115
 - presenting results as 112
 - statistical 142–8
- t*-distribution 28
 - confidence interval for the mean 34
 - statistical table 142, 143
- test statistic, obtaining the 50
- text files 16
- text format 16
- times, data entry problems 16
- time-varying confounding 105
- training sample 139
- transformations
 - method 30
 - reasons for 30, 107
 - typical 30–1
- treatment allocation 40–2
- treatment effect 52
- trend, Chi-squared test for 72–3
- trial size 43
- trials *see* clinical trials
- true negatives 115
- true positives 115
- t*-test 34
 - one-sample 54
 - for partial regression coefficients 84
 - paired 57
 - unpaired (two-sample) 60, 109
- two-level structure, clustered data 124, 127
- two-sample *t*-test
 - see* unpaired *t*-tests and Wilcoxon rank sum test
- two-tailed tests 50
- Type I errors 52
 - clustered data 124
 - sample size 108
- Type II errors 52
 - sample size 108
- typing errors 18, 19
- unbiased estimate 32
- underdispersion
 - likelihood ratio statistic 97
 - logistic regression 90
 - Poisson regression 94
- uniform distributions 21
- unimodal distributions 21
- unit, experimental 38
- unpaired *t*-test, *see* two-sample *t*-test
- validation, prognostic scores 139
- validation sample 139
- variability
 - between-subject 109
 - sample size calculation and 109
 - within-subject 24
- variables 14
 - random 26, 38–9
- variance 24–5
 - Binomial 29
 - heterogeneity of 106, 130
 - homogeneity of 106, 130
 - Poisson 29
 - residual 63, 79
 - stabilizing 30, 31
 - testing for equality of two 64, 106, 107, 153
- variance-ratio test (*F*-test) 106
 - multiple linear regression 85
- variation 38
 - between-group 63
 - between-subject 24
 - coefficient of 24
 - explained 80
 - extra-Poisson 93–4
 - over time 93
 - random 38
 - unexplained (residual) 80
 - within-group 63
 - within-subject 25
- Wald test 88
 - logistic regression 88, 89
 - Poisson regression 93
- washout period 38
- Weibull model 134
- weighted kappa 118
- weighted mean 22, 23
- Wilcoxon rank sum (two-sample) test
 - 60–1
 - statistical tables 142, 146
- Wilcoxon signed ranks test 55, 57–8
 - statistical table 142, 145
- within-subject/group variation 25, 63, 118
- zero cell count, logistic regression 89
- z*-test 54
- z*-value 55, 56, 142