

Index

- 3 Rs, (essence of coping in active labor), 26, 160, 216
- abdomen, pendulous, 24, 118–9, 143, 320
- abdominal lifting, 119, 140, 144, 203, 318–20, 357
- precautions for, 119, 144, 319
- abdominal shape, as a sign of OP, 52, 54
- abdominal “jiggling”, 319–20, precautions for, 320
- abdominal stroking, 104, 140, 317–8, 366
- abuse, survivors of, needs of, during labor, 112–3, 161, 164–5, 216
- vaginal exam with, 69
- active labor, prolonged, definitions of, 22, 125–6
- management of, 126–8
- possible causes, 127–33
- active management of labor, 122, 126, 229–32
- acupressure, 46, 154, 345–7
- avoiding before term, 346
- acupuncture, 154, 267–9, 347
- ambulation, *see* walking
- American College of Nurse-Midwives, 88
- American College of Obstetricians and Gynecologists, 3, 37
- artificial rupture of membranes (AROM), 17, 18, 107, 127–8, 140, 156, 165, 246
- as a possible factor in dystocia, 134–5, 246
- need for more study, 135
- risks, 135, 246
- Association of Women’s Health, Obstetric and Neonatal Nurses, 88
- assessments, caput, 80, 82
- cervix, prenatally, 64–6
- cervix in labor, 66–8, 70–2, 177
- see also* cervix
- contractions, 81–4
- see also* contractions
- descent, abdominal assessment of, 73–5

The Labor Progress Handbook: Early Interventions to Prevent and Treat Dystocia.

Edited by Penny Simkin, Ruth Ancheta

© 2011 by Penny Simkin and Ruth Ancheta; illustrations copyright Ruth Ancheta

- assessments (*cont'd*)
- vaginal assessment of,
 - 72–3, 75, 80
 - dilation, 70, 126–7, 177
 - effacement, 65, 70
 - fetal well-being, 16, 28, 86–94,
 - 153, 247, 311
 - fetal position, prenatally, 52–64
 - fetal position, in labor, 66,
 - 69, 76–9, 133–4
 - accuracy of clinical
 - assessments vs.
 - ultrasound during
 - labor, 79
 - fetal weight, 57, 64,
 - as an estimation, 64
 - prediction equation, 64
 - flexion, 79
 - maternal well-being, 84–6
 - membranes, 71
 - molding, 80
 - pelvis, 80,
 - presenting part, 72
 - progress in first stage, 94–6, 125
 - progress in second stage, 96,
 - 174, 185, 186–8
 - synclitism/asynclitism, 76–9
 - woman's emotional state, 85–6,
 - 160–6, 112–3, 215
 - urge to push, 146
 - see also* emotional dystocia
 - and maternal coping,
 - signs of
- asymmetrical dilation, *see* cervical lip
- asymmetrical positions and movements, 141–2, 192, 196, 203, 283–4, 296, 313–4
see also lunge
- asynclitism, 24, 35, 76–79,
 - 116–120
 - in latent labor, 116–20
 - in active labor, 127–8, 130–1
 - and AROM, 134–5
 - and cervical lip, 250
 - maternal positions for, 117–8,
 - 136–42, 200, 203,
 - 284, 288–94,
 - 296–306
 - maternal movement for, 144,
 - 311–316, 319–324
 - in pre-labor and latent labor,
 - 115–8
 - repositioning *see* manual or digital rotation of the fetal head
 - signs of, 76–7, 79
 - in second stage, 200–203
 - see also* malposition
- augmentation, *see* contractions, inadequate
- Baby-Friendly Initiative, 236–7
- back pain,
 - measures to alleviate, 103–6,
 - 114, 119–21, 136–45,
 - 188, 191, 354–61
 - in pre-labor and latent labor,
 - 103–6, 114, 119–21
 - in active labor, 136–45
 - in second stage, 188, 191
 - as a clue to progress, 94
 - as a clue to malposition,
 - CPD, 129, 20
 - nuchal hand and, 204
 - TENS for, 114, 136, 368–71
 - see also* malposition; occiput posterior (OP) *and* occiput transverse

- bearing down, 174, 176–7, 374–6
 delayed, 183–5
 diffuse, 180–1, 216–7, 375
 directed, 13, 178–9, 182, 203, 258, 375–6
 directed by clinician, 258
 epidural, effect on, 182–5
 epidural, helping women push with, 185–7
 fear as an impediment to, 216–9
 “holding back”, 217
 “laboring down”, *see* bearing down, delayed
 with a malpositioned fetus, 190–203
 see also malposition, maternal positions and movements for
 premature urge *see* urge to push
 relaxing the perineum and, 218–9
 self-directed, 181, 374–5
 spontaneous, 17, 19, 67, 96, 176–7, 179–81, 186, 188, 203, 250, 253, 257–8, 271, 374
 verbal support for, 253–4,
 “trial pushes”, 146
see also Quick Epidural Index; second stage, urge to push *and* Valsalva Maneuver
- bed rest in late pregnancy, 114, 336
 “Belly Mapping”, 60–3
 birth ball, 32–3, 38, 137, 156–7, 286, 289, 291, 312, 320, 324, 367
 precautions for, 368
 birth plan or preference list, 31–2, 238
 birth rope, 301–2
 precautions for, 302
 birth sling, 194, 300–1
 precautions for, 302
 bladder, emptying,
 before examinations and procedures, 57, 68, 248
 every hour or two, to reduce pain and facilitate progress, 34, 261
 breast crawl, 236
 breastfeeding, 12, 17, 19, 20, 27, 28, 226–8, 230, 234–8
 Bishop score, various interpretations of, 65–6
 breathing,
 in latent labor, 110, 114
 in first stage, 114, 159, 165–6
 when to begin, 110
 naturalistic, 328,
 in second stage, 177, 216, 218
 see also bearing down
 supporting and teaching, 371–3
 through contractions in second stage, 183–5
- caput, 72–3, 80, 96, 135, 200, 323
 cardinal movements, 35, 108–9
 effect of epidural on, 149, 183–5

- care plans (flowcharts),
 - little or no progress in labor, 7
 - anxiety or distress in labor, 165
 - prolonged prelabor or latent phase, 107
 - prolonged active phase, 128
 - occiput posterior/asynclitism in active first stage, 140
 - premature urge to push, 146
- complete dilation, with no epidural, 180
- diffuse pushing with no epidural, 182
- complete dilation with epidural, 183
- occiput posterior/asynclitism in second stage, 203
- carpal tunnel syndrome, 292, 342
- catecholamines, 27–30, 112, 158, 162, 301, 333, 351,
- cephalopelvic disproportion (CPD), 22, 24, 94, 127, 129,
 - clues to, 76, 80, 132
 - and AROM, 134, 246
 - and dorsal positions, 211–4
 - caused or made worse by malposition, 129, 132–3
 - suspected in active labor, 22, 135–6
 - maternal positions for, 146–7
 - suspected in second stage, 177–8, 188, 190–1, 205–14
 - maternal positions and movements for, 188, 190–1, 205–14
 - see also* assessments, fetal weight
- cervical dystocia, 24, 65, 71, 245
- cervical lip, 71, 129, 146, 156–8, 250–1, 363,
 - and asynclitism, 250
 - interventions for, 250–1
 - manual reduction of, 250–1
 - positions for, 293–5,
 - possible other interventions, 158
- cervix,
 - assessing prenatally, 64–6
 - assessing during labor, 66–8, 70–2, 177
 - difficult to find, 71
 - dilation, 32–3, 64–8, 70–2, 95–6, 102, 177
 - effacement, 65, 70–1, 95, 102, 106, 108, 112–3, 245, 248,
 - as a factor in dystocia, 24, 65, 71, 245
 - position of, 108–9
 - remodeling of, 65, 111
 - see also* cervix, ripening
 - retraction of, 81, 175–6
 - rigid os (stenosis), 24, 71, 129
 - manual stretching of, 245
 - ripening of, 65–6, 106, 108, 113
 - scarred, 24, 113
 - swollen, 71, 146, 156–7, 292, 294–5
 - “zipper” cervix, 71
 - manual stretching, 250

- cesarean birth, 3, 11, 13, 17,
 - 22–3, 37, 64, 73,
 - 107, 110, 126, 128,
 - 134–5, 140, 149,
 - 151–2, 162, 165,
 - 177, 203, 205, 217,
 - 219, 246–8, 250,
 - 257, 270, 328, 349
- potential to reduce cesarean rate using:
 - longer time frame before diagnosing dystocia, 22–23
 - sterile water injections, 270
- clinical interventions, low-technology, 242–72
- intermediate-level interventions, 243
- for use by clinical personnel only, 243
- tertiary care, referral or consultation, 243
- see also* doulas, role during clinical interventions
- prelabor and latent labor interventions, 244
- nipple stimulation, 244
- management of cervical stenosis or “zipper” cervix, 245–6
- therapeutic rest, 244–5
- active labor and second stage interventions, 245–51
- artificial rupture of the membranes (AROM), 246
- digital or manual rotation of the fetal head, 246–50
- manual reduction of a persistent cervical lip, 250–1
- fostering normality in second stage labor,
 - guiding women through crowning of fetal head, 254–5
 - hand skills to protect the perineum, 255–6
 - maternal birth positions 253–4
 - perineal management in second stage, 252–3
 - prenatal perineal massage, 251–2
 - differentiating perineal massage from other interventions, 256–7
- when progress in second stage remains inadequate, 257–8
- duration of second stage labor, 257–8
- directing bearing down efforts, 258
- hand maneuvers and anticipatory management for intrapartum problems
- shoulder dystocia, 258–65
 - precautionary measures, 260–1
 - shoulder dystocia maneuvers, 261–5
 - warning signs, 261
- nuchal cord, somersault maneuver, 265–7

- clinical interventions, low-technology (*cont'd*)
 - non-pharmacologic and minimally invasive pain relief, 267–271
 - acupuncture, 267–9
 - nitrous oxide, 271
 - subcutaneous sterile water injections, 269–71
 - topical anesthetic on perineum, 271
 - cold,
 - for hemorrhoids, in second stage, 332
 - for pain relief, 33, 136, 140, 198, 283, 330–2, 363–5
 - precautions when using, 331–2
 - to reduce swollen cervix, 330
 - consultation and referral, 16, 93, 93, 243
 - continuous support, 328, 347–50
 - from doulas, 347–9
 - from nurses and midwives, 349–50
 - contractions,
 - assessing, 81–4, 125, 74–6, 146–153
 - inadequate, 24, 146–51, 215
 - and dehydration, 150–1,
 - and exhaustion, 151
 - and immobility, 147–8
 - and medication, 149
 - and uterine lactic acidosis, 151–3
 - also see* emotional dystocia
 - irregular or coupling
 - contraction pattern, as a sign of possible malposition, 115, 116, 100, 120, 132,
 - in pre-labor or latent phase, 95
 - non-pharmacological ways to stimulate, 24, 45–6
 - acupressure, 46, 154
 - breast or nipple stimulation, 45–6, 153,
 - heat on fundus, 46
 - hydration, 45, 150
 - hydrotherapy, 45, 154–6
 - movement and positioning, 45, 153
 - prevent/alleviate fear, 24, 29, 31–3
 - touch, 45
 - non-progressing, 102
 - patterns in latent and active second stage, 174–5, 177
 - pre-labor, 102, 106–7
 - progressing, 102
 - tetanic, 153, 177,
- cooperation between patients and caregivers, 5–6, 30–3, 60–3, 66–9, 86, 112–3, 151–2, 230, 238
 - coping with labor pain, *see* maternal coping
 - cord
 - compression, 91, 94, 179, 246
 - positions to help prevent or reduce. 280, 291, 293

- clamping and cutting,
 - 228–30, 232, 237,
- nuchal, and somersault
 - maneuver, 229
- counterpressure,
 - on the back, 197, 293, 317,
 - 354–5
- Cook's counterpressure on
 - isical tuberosities,
 - 359–61
- Cood's perilabial
 - counterpressure
 - 361–3
- cultural factors, 4, 7–8, 158–9,
- 162–3, 166, 332,
- 337, 353
- non-clinical factors influencing
 - care, 9–11, 23
- deflexed head and CPD, *see*
 - flexion of fetal head
 - on chest
- dehydration
 - see* hydration and dehydration
- “difficult” patients, 164
- diffuse pushing *see* bearing
 - down
- dilation,
 - as a feature of normal labor,
 - 21–2, 95
 - assessing, 70, 126–7, 177
 - asymmetrical, 250
 - when cervix is scarred, 15, 92
 - and definitions of dystocia,
 - 21–3
 - and definitions of labor onset,
 - 102–3
 - and hydrotherapy, 154, 336–7
 - one of many ways to progress
 - in labor, 108–9
 - in pre-labor and latent phase,
 - 106–7
 - as a sign of active labor, 125
 - as a sign of readiness to push,
 - 176–7
- double hip squeeze, 195–7, 355–7
 - distinguished from pelvic
 - press, 195
- doulas, 10, 12, 52, 243, 328,
- 347–50
 - role during clinical
 - interventions, 234,
 - 238, 243, 246, 248,
 - 254, 263, 271,
- drinking in labor, *see* hydration
 - and dehydration
- drive angle, 35–6, 213, 307, 309
- drug-induced rest, 102, 107,
- 115
- dystocia,
 - and cesarean rates, 3, 23,
 - 135, 152, 250,
 - etiologies of,
 - cervical, 24, 65, 71, 111–2,
 - 129, 245
 - emotional, 4, 24, 112–3,
 - 129, 215–9
 - extrinsic, 4,
 - fetal, 4, 24, 112, 127, 204,
 - iatrogenic, 4, 24, 111, 127,
 - 147–9, 182–5
 - intrinsic, 4
 - maternal, 4, 24, 129, 150–3
 - pelvic, 4, 24,
 - unknown, 129, 153–6
 - uterine, 4, 24, 82, 129,
 - see also* contractions,
 - inadequate
 - various definitions of, 21–3,
 - 101–3, 125–7, 174–8

- early interventions,
 and their place in the care of
 laboring women, 4,
 6–7, 8, 14, 127, 129,
 188, 200, 278
see also, interventions/actions,
 primary, secondary
 and tertiary
- eating in labor, 34, 84, 85
also see hydration and
 dehydration
- effacement,
 assessing, 65, 70–1, 95,
 AROM done after 100%
 effaced, 246
 when cervix is scarred,
 112–3
 and definitions of labor onset,
 102
 one of many ways to progress
 in labor, 108,
 in pre-labor and latent phase,
 95
also see Bishop Score
- emotional dystocia, 4, 24,
 85–6, 112–3, 129,
 160–8, 215–9
 assessment, 162, 216, 351–2
 common fears, 162, 216–7
 as distinguished from
 “holding back”, 217
 effects on fetus, 29
 physiology of, 29, 112
 predisposing factors, 112,
 163, 216–7
 preventing, 31–2, 85–6,
 110–11, 349, 352–4
 ways to help, 132, 163–8,
 217–9
see also abuse, survivors of
- environment, effect on labor
 progress, 4, 28, 31,
 34, 86, 158, 226,
 230, 253, 350–1,
 353
see also emotional dystocia
- epidural, 128, 182–6 *see also*
 Quick Epidural
 Index
 adjusting dose, 185
 advantages of, 128, 146, 165,
 327
 bearing down with, 183–4,
 care plan for, 183
 delayed bearing down, 183
 possible benefits of delayed
 bearing down plus
 lateral position, 184
 effect on fetal position, 129,
 149, 156, 182–5
 effect on progress, 129, 149,
 182–5, 189
 maternal mobility, variations
 in, 184
 maternal positions that may
 be usable with
 epidurals, 138–9,
 147–9, 182–5, 187,
 305–6, 308–10
 non-pharmacological ways to
 augment
 contractions (to
 avoid increased pain
 associated with
 synthetic oxytocin
 regimens), 23, 45–6,
 153–4, 156, 337
- precautions,
 avoiding hydrotherapy
 336–7

- avoiding pelvic press, 242
 - protecting joint and nerve injuries when using lithotomy and semi-sitting positions, 214, 310
 - when using heat and cold, 329–32
- removing time limit in second stage, 185–8
- sheet pull, 306
- trying other measures first, 153–4, 156, 337
- upon request, 149
- used more in recent years, 22
- exhaustion, 24, 84, 109, 112, 114, 128–9, 151, 160, 162, 179, 215–6, 244,
- “failure to progress”, 3, 22, 125, 250
- fear, 24–5, 28–32, 69, 112, 129, 160–3, 167, 216–8, 278, 327, 351–2, 354
- fetal ejection reflex, 28–30,
- fetal influence on labor, 23
- fetal attitude *see* flexion of fetal head on chest
- fetal monitoring, 37–44,
 - audible vs. silent, 352
 - auscultation, 24, 37, 87–91
 - comparison with EFM, 37
 - brief, when using abdominal lifting or abdominal “jiggling”, 99, 119, 144, 319, 321
 - and cesarean and instrumental delivery rates, 37
 - continuous, 7, 17, 19, 37–40, 42–4, 86–7, 89–91, 111, 248,
 - explaining, 163, 357
 - as feedback to aid bearing down in second stage, 186
 - and hydrotherapy, 41–4, 335,
 - intermittent, 41, 87, 313, 335,
 - risks, 24, 37
 - telemetry, 42–4, 335,
 - and TENS, 369–70
 - three-tiered system, 91–3
 - without immobilizing the woman, 35–44
- fetus,
 - pre-term, 65, 66, 90, 93, 228, 235
 - post-term, 66, 93
 - cervical ripening, role in pre- and post-term births, need for study, 66
 - “fight or flight” response, 28–30
 - using heat to reduce, 328
 - see also* “tend and befriend” response
 - flexion of fetal head on chest, 35, 79, 96, 109, 142, 183, 200,
 - after epidural, 183,
 - effect of maternal dorsal and semi-sitting positions, 142–3
 - maternal positions to promote, 289–90,
- fourth stage, *see* third and fourth stage

- fundal pressure,
distinguished from
suprapubic pressure,
264
- gestational age, 93–4, 179
- gravity, 8, 19, 35, 104, 114,
117–20, 131, 136,
138, 139, 142–3,
145, 156, 178, 181,
188, 191, 230, 254,
265, 289–3
- as an aid to progress, 117–20,
131, 138–9, 145, 156,
158, 178, 206–7, 212,
278, 280–3
- negative (“anti-gravity”)
positions, 156, 178
- neutral positions, 156,
158, 214, 278, 280,
- positive, 181, 191, 193
- and shoulder dystocia, 265
- and protecting the perineum,
254
- and birth of the placenta, 230
- head compression, 135, 179
- heat,
to augment contractions, 46
for pain relief, 328–30, 363–5
precautions when using, 329,
365
to reduce “fight or flight”
response, 328
- hemorrhoids, 280, 283, 293–5,
330, 332
- hopelessness, 112, 114, 216
- hydration and dehydration, 24,
34, 45, 84, 85, 128–9,
146, 150–1, 215
- hydrotherapy, 140, 155–6, 327,
332–7, 365
for anxiety and distress, 164
depth of bath, 336
and fetal monitoring, 41–4,
335
for labor augmentation, 154,
337
for pain relief, 154, 199, 337
precautions, 336–7
leaving water periodically,
335
timing of, 154, 336–7
when not to use, 336–7
- hypertension, 111, 114, 147,
283, 286, 336
- hypotension, supine, 44, 142,
147, 179, 212, 214,
278, 280, 283, 306,
308, 309
- iatrogenic dystocia, *see* dystocia,
etiologies of,
iatrogenic
see also hydration and
dehydration; fetal
monitoring,
continuous
and positions, maternal,
dorsal, problems
with
- ice massage, 346
- induction of labor, 17–9, 24,
65, 107, 113, 153,
350
breast stimulation for, 153
acupressure for, 346
see also contractions,
inadequate *and*
oxytocin, synthetic

- informed consent, 10, 38, 134, 248,
- instrumental deliveries *see*
operative deliveries
- interventions/actions,
primary, secondary
(intermediate) and
tertiary, 6–7, 242
- intra dermal sterile water
injections, *see* sterile
water injections
- knee press, 136, 198, 199, 357–9
- labor and birth, normal,
definitions and descriptions
of, 16–21
active, 95–6, 125–6
latent phase, 95, 102–3,
106
second stage, 96
fostering, 251–7
movement as a component, 20
psychosocial outcomes of, 20
work, strenuous, as a
component, 20
see also dystocia, various
definitions of
- labor onset, various definitions
of, 102–3
- “Laboring down,” *see* bearing
down, delayed
- latent phase of first stage,
definitions of, 102–3
- latent phase of second stage,
174–7
- lactic acidosis, uterine, 24,
151–3
- Leopold’s maneuvers, 52, 57–9
- lordosis, lumbar, 129
- lunge, 128, 187, 195–6, 203,
210, 302–4
instructions, 141–2, 313–4
precautions for, 313–4
side-lying, 139–40, 283–4
see also movements, maternal
- macrosomia, 64, 94, 113, 129,
132, 135, 205,
and AROM, 134, 246
see also cephalopelvic
disproportion
(CPD) and assessing
fetal weight
- malposition,
and AROM, 134
as a cause of dystocia, 4, 24,
35, 52, 112, 129–33,
188–9
clues to, 52–63, 134
and contraction patterns, 133
and maternal dorsal positions,
35–6, 188
see also positions, maternal,
dorsal, problems with
- and epidurals, 149, 182–5
- maternal positions and
movements for, 35
in pre-labor and latent
phase, 115–20
in active labor, 136–48
in second stage, 188–95
and operative delivery
rates, 129, 149
preventing *see* Optimal Fetal
Positioning
see also back pain; occiput
posterior; occiput
transverse and
asynclitism

- malpractice, 11
- manual reduction of a persistent cervical lip, 158
- manual or digital rotation of the fetal head (manual repositioning), 145, 246–50, 319
- massage, *see* touch and massage
see also ice massage, perineal massage
- maternal coping, 25–6, 67, 95, 110–13, 126, 129, 158–60, 166, 215–6, 225–7, 236, 323–4, 351, 353, 37–2,
coping scale, 26
signs of coping,
in first stage, 95, 158–60, 166, *see also*, 3Rs
in second stage, 215–6,
in third and fourth stages, 225–7, 236
- maternal distress, 24, 46, 85
preventing, 30–5, 112–3, 161–2, 350–4
see also emotional dystocia; hormones of labor
- maternal movements, *see* movements, maternal
- maternal positions, *see* positions, maternal
- maternity care practices,
comparison of US/UK/
Canada, 8–11, 32, 149, 247,
- meconium, 86, 93–4, 228, 244
- medication, 27, 128, 146, 149, 164–5, 337, 349, 371,
benefits, 27, 128, 146, 164–5
and intensity of contractions, 149
and malposition,
see also epidural, effect on fetal position
and maternal positions that may be usable with epidurals
as a possible contraindication to hydrotherapy, 337
narcotics, 128, 140, 149, 165
nitrous oxide, 271
used less by women with doulas, 349
used less by women using TENS, 371
topical anesthetic on perineum, 271
see also epidural; drug-induced rest, *and* the Quick Epidural Index
- midwifery model of care, 20, 23, 126–7, 158, 230
- moaning, *see* vocalization
- molding, 15, 73, 80, 96, 108–9, 135, 200, 205
- monitoring, fetal, *see* fetal monitoring
- movements, maternal,
as an aid to progress, 35, 42, 45, 200, 202, 311–24
as a comfort measure, 136–44, 311–24
see also 3Rs of labor coping
effect on oxygen supply to fetus, 35, 311,
effect on maternal pelvis 4, 24, 35

- and fetal monitoring *see*
 monitoring, fetal
- positions and movements for
 suspected malposition or
 CPD in active labor,
 134–43
- suspected malposition or
 CPD in second
 stage, 188–95
- specific movements,
 abdominal “jiggling” with a
 rebozo 320–1
- precautions when using,
 321
- abdominal lifting, 318–20
- precautions when using,
 319
- abdominal stroking, 317–8
- lunge, 313–4
- other rhythmic movements,
 323–4
- pelvic press, 321–3
- precautions for, 323
- pelvic rocking, 311–2
- slow dancing, 316
- walking and stair climbing,
 316–7
- see also* optimal fetal
 positioning
- neocortex, role in labor
 progress, 29, 34, 158
- newborn, *see* third and fourth
 stage labor
- nipple stimulation
see contractions, inadequate,
 non-pharmacological
 ways to stimulate
- nuchal hand, 204
- see also* back pain
- observation, 5–6, 25, 30, 60, 67,
 92, 129, 160, 163–4,
 176, 179, 203, 204,
 228, 238, 351,
- occiput posterior (OP), 24, 35,
 53, 127
- in active labor, 129–32
- in second stage, 188–9
- and AROM, 134–5
- and cervical lip, 250
- determining direction of,
 54–63, 133–4,
- disadvantages of supine and
 semi-sitting
 positions for, 143,
 188, 212–3
- see also* positions, maternal,
 dorsal, problems
 with
- increased incidence with
 epidurals, 149
- and long pre-labor or latent
 phase, 112
- maternal positions for active
 labor, 136–41,
- maternal positions for second
 stage, 190–5
- and premature urge to push,
 144
- preventing, 103–6
- re-positioning *see* abdominal
 lifting; abdominal
 stroking *and* manual
 repositioning
- and side-lying positions,
 138–9
- signs of, 54–9, 132–3
- see also* back pain;
 malpositions *and*
 positions, maternal

- occiput transverse (OT),
 persistent, 78, 132–3
 in active labor, 132–3
 and AROM, 134–5
 deep arrest and epidurals,
 183–5
 deep arrest and pelvic press,
 321–3
 identifying, 168–9, 201 (if
 both OT and
 asynclitic)
 see also assessments, fetal
 position
 maternal positions for,
 in active labor, 136–9
 in second stage, 190–5
see also positions, maternal
 and rotation
- Optimal Fetal Positioning, 62,
 103, 136
- oxytocin
 endogenous, 27–9, 45, 153,
 176, 184–5, 226,
 228, 236, 246, 267,
 333, 335
 synthetic, 13, 17, 37, 22–4,
 87, 107, 111, 126–9,
 133–5, 140, 147,
 152–4, 177, 165,
 177, 180, 183, 205,
 219, 231, 244, 337,
 reduced need for among
 women using water
 immersion? 154
 trying nipple stimulation,
 cautiously, to
 augment labor, in
 low-risk women,
 before giving oxytocin,
 244–5
- when there is uterine lactic
 acidosis, 151–3
also see third and fourth
 stages, placenta,
 uterotonic
 medications
- pain, 16, 24–7, 84, 95, 346
 as a factor in progress, 4, 278
 excessive, reasons for, 4,
 34–5, 111–3, 332
 intensity scale and coping
 scale, 26
 pain vs. suffering, 24–5, 27
 patience, 22–3, 85, 126–7, 164,
 205
 pelvic dystocia, 24, 127–8, 205
 pelvic floor damage, 213
 pelvic press, 195–6, 203, 321–3,
 precautions when using, 323
 distinguished from double
 hip squeeze, 195,
 356
 pelvic rocking, 311–3
also see movements, maternal
- pelvis,
 effect of movement on, 4, 24,
 35
 see also maternal
 movements
 effect of position on, 24,
 35–6, 139, 141, 192,
 195, 200
 late pregnancy changes in,
 35–6
- perineal lacerations, 204
 perineal massage, 251–2
 differentiated from other
 interventions 256–7
 perineum, supporting, 256,

- policies, effect on labor
 progress, 4, 8, 9, 34,
 38, 111, 151, 156,
- placenta, *see* third and fourth
 stages/ third stage/
 placenta
- positions, maternal,
 as an aid to progress, 35, 118,
 120, 203
- categories of, 278
- asymmetrical, 278
- dorsal positions, 278
- and drive angle, 35–6,
 213, 307, 309
- and instrumental
 deliveries, 211–3
- problems with, 35–36,
 142–3, 211–3, 278
- forward-leaning, to
 reposition fetus or
 reduce back pain,
 278
- in active labor, 137, 145
- in late pregnancy, 103–6
- in pre-labor and latent
 phase, 117, 120
- in second stage, 190
- gravity negative (“anti-
 gravity”), 156, 294
- gravity neutral, 278
- gravity positive, (“gravity-
 enhancing”), 278
- horizontal, 278
- for tired women, 152
- specific positions:
- asymmetrical, 278
- asymmetrical upright,
 296–7
- asymmetrical horizontal
 (side-lying lunge), 284
- dangle and birth sling,
 299–301
- exaggerated lithotomy
 (McRoberts’) use of,
 309–10
- precautions for, 310
- hands and knees, 309–10
- knee-chest positions,
 open, 293–5
- and hemorrhoids, 294
- closed, 295–6
- kneeling leaning forward
 with support, 290–2
- roll-over sequence, 148, 282
- semi-sitting, 285
- sheet pull, 229–31
- side-lying, (lateral and
 semi-prone),
 279–81
- which side to lie on if
 fetus is
 malpositioned, 281
- side-lying lunge, 284
- sitting
- leaning forward, 287–8
- upright, 286–7
- squatting positions,
 297–302
- precautions for half-
 squatting, 303–4
- precautions for lap
 squatting, 305–6
- precautions for
 supported squatting,
 301–2
- standing leaning forward,
 289
- supine, 306
- see also* dorsal positions,
 problems with

- positions, maternal (*cont'd*)
 Why focus on maternal position? 35–36
- post-traumatic stress disorder (PTSD), 25, 30
- pre-labor and latent phase, prolonged, 87
 definitions of, 102
 “false” labor, 106, 113
 general suggestions for, 103–7, 108–11, 115–21
see also contractions, irregular or coupling
 management styles for, 102–3, 106–7, 113–121
 “over-reacting” to labor, 110
 possible causes, 111–3
 therapeutic rest, 244
- premature rupture of the membranes (PROM) 132,
- previous back injury, 119
- previous trauma, effect on labor progress *see* emotional dystocia
- psychosocial comfort measures, 30–5, 112–3, 161–2, 350–4
see also emotional dystocia and vaginal exams, instructions for
- pushing *see* bearing down
 “pushing positions” as distinguished from “delivery positions”, 185–7, 211–3
- relaxation, 31, 33, 110, 114, 154, 216, 218, 248, 249, 252, 267, 278, 328,
 as one of the 3 Rs, 75–6, 118–9, 143, 320, 371–3
- respect, 4, 27, 30, 32–3, 165–6, 263, 352, 371
- resting the uterus, 151–2
- restriction to bed, 13, 111, 114, 184,
 minimizing, 38–44, 111, 114, 336
- rhythm, 25–6, 159–60, 216, 327–8, 354, 371–3
 as one of 3Rs, 75–6, 118–9, 143, 320
- ritual, as one of 3rs, 25–6, 159–60, 216, 371–2
- rollover sequence, 148, 282
- rotation,
 encouraging in pre- and early labor, 103–4, 119–20
 after an epidural, 149, 183–4
 after AROM with a malpositioned fetus, 134–5
 in latent phase of second stage, 176
 lack of, as a factor in dystocia, 127, 129, 132
 manual and digital rotation, 246–50
 positions to encourage, 103–6, 136–42, 190–5

- six ways to progress in labor, 87–9
- Royal College of Obstetricians and Gynaecologists, 18, 37
- Royal College of Midwives, 18
- second stage,
 - definition of, 146
 - epidural during, 156–61
 - fear as an impediment to progress, 188–92
 - management styles for, 146–7, 150–62
 - phases of, 147–51
 - positions for, *see* positions, maternal
 - prolonged latent phase of, 150
 - prolonged, possible causes, 162, 188,
 - pushing during latent phase, 149–50
 - time limits on, 159, 161–2
 - Valsalva maneuver during, 178–81, 256, 374
 - verbal support during, 253
 - see also* bearing down
- sheet pull, 308–9
- short waist, 119, 129
 - see also* abdominal lifting
- shoulder dystocia, 94, 214–5, 258–65
 - precautionary measures, 260
 - warning signs, 261
 - maneuvers, 261–5
 - see also* supra-pubic pressure, distinguished from fundal pressure
- six ways to progress in labor, 32, 108–9
 - see also* movements, maternal, and positions, maternal
- slow dancing, 32, 38, 195–6, 211, 316–7, 366,
- Society of Obstetricians and Gynecologists of Canada (SOGC), 17, 37, 88, 247,
- station, assessing,
 - see* assessments, descent
- sterile water injections for back pain, 136, 140, 199, 269–71, 371
- supine positions *see* dorsal positions
- suprapubic pressure, 261–2
 - distinguished from fundal pressure, 264
- survivors of abuse, *see* abuse, survivors of
- “tend and befriend” response, 30
- third and fourth stages, 225–238
 - overview, 225–6
 - mutual regulation by mother and newborn, 225–7, 236
 - skin-to-skin contact, 230, 234–8
- third stage:
 - definition, 225
 - family integration, 225–7, 235
 - hemorrhage, 231–4
 - hormones of, 225–7

- third and fourth stages (*cont'd*)
- newborn,
 - assessments, 237–8
 - behavior at birth, 225
 - blood volume, 228
 - delayed cord clamping, 228–9
 - early behavior, 225
 - meconium, 228
 - sensory experience, 225
 - suctioning, 227–8
 - transition, 225–8
 - placenta,
 - active management, 229–34
 - expectant management, 229–34
 - guarding the uterus, 231
 - hemorrhage, 229, 231–4
 - uterine massage, 231
 - uterotonic medications, 230–4
 - fourth stage:
 - Baby-friendly breastfeeding practices, 236
 - Ten steps to successful breastfeeding, 237
 - definition, 234
 - keeping mother and baby together, 234–8
 - procedures done with baby in mother's arms, 238
 - routine newborn assessments, 237–8
 - time as an ally, 5, 205
 - touch and massage, 33, 45, 165, 337–45
 - and endogenous oxytocin production, 45
 - mini-massages of shoulders, back, hands, feet, 339–45
 - to reduce anxiety and pain, 338
 - transcutaneous electrical nerve stimulation (TENS)
 - for back pain, 114, 136, 368–71
 - not for use during hydrotherapy, 370
 - trial and error, 5, 23, 76, 129, 130, 134, 141, 188, 194, 282
 - trust, ways to enhance, 5–6, 30–3, 60–3, 66–9, 86, 112–3, 151–2, 230, 238
 - UK/US/Canada, differences in maternity care, 8–11
 - ultrasound predictions of fetal size, 205
 - ultrasound to identify fetal position,
 - before manual or digital rotation, 247
 - during pregnancy, 59, 60, 64, 72, 76
 - during labor, 79, 104, 129, 133, 134, 138, 140,
 - urge to push,
 - absence of, during latent phase of second stage, 174–77
 - bearing down with, 180, 216, 374–5
 - delayed, reasons for, 174, 183–6

- and epidural anesthesia, 183–6
 - see also* Quick Epidural Index
- premature
 - as a sign of possible malposition or CPD, 133, 144–5
 - ways to deal with, 144–6, 294, 361–3
 - ways to enhance, 177, 290, 298
 - see also* second stage, latent phase of
- uterine dystocia, 24, 82, *see also*
 - contractions, inadequate
- uterine massage, 230–3
- vaginal birth after cesarean (VBAC), 3, 217
- vaginal exams,
 - and abuse survivors, 69
 - fear of, 69
 - indications for, 66–7
 - instructions for, 66–72
 - positions for, 68
 - reactions to, in early labor, 69
 - timing of, 66, 67
 - see also* assessments
- Valsalva maneuver, 178–81, 253, 256, 374
 - history of prescribing, 178
 - problems caused by, 178–80
 - alternatives, when women have epidurals, 182–6
- visualization, 31, 33, 114, 163, 165,
- vocalization, 33, 114, 159–60,
- walking and stair climbing, 35, 42, 114, 147, 177, 280, 315–7
 - compared to AROM plus oxytocin, 147
 - see also* movements, maternal
 - woman as key to solution, 5
 - woman-centered care, 10
- World Health Organization (WHO), 4, 16–7, 73–4

